**Zeitschrift:** Archives des sciences [1948-1980]

Herausgeber: Société de Physique et d'Histoire Naturelle de Genève

**Band:** 14 (1961)

Heft: 10: Colloque Ampère

**Artikel:** Measurements of the microwave spectra of meta-fluoro-chloro-benzene

Autor: Rachman, A.

**DOI:** https://doi.org/10.5169/seals-739599

# Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

## **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF:** 30.10.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# Measurements on the Microwave spectra of meta-fluoro-chloro-benzene

# by A. RACHMAN

Faculty of Exact and Natural Sciences, University of Buenos Aires, Argentine \*

#### Sommaire.

On a mesuré le spectre en microondes de la molécule 1-3  $C_6H_4ClF$  avec un spectromètre à modulation Stark d'absorption en gaz. On a utilisé divers klystrons, en couvrant la région dès 11,5 GHz. jusqu'au 26,3 GHz. En comparant le spectre mesuré et le spectre calculé d'après un modèle de la molécule, on donne les premières interprétations et on discute ses fondements.

It has been measured the microwave absorption spectra of the molecule 1-3 C<sub>6</sub>H<sub>4</sub>ClF on the way to continue a systematic study of the substituted benzene ring [1, 2, 3, 4]. The measurements have been made in a conventional 100 KHz Stark effect modulated gas absorption spectrometer with square wave modulation, described in [2]. Several klystrons have been used and the spectra was measured from 11,5 GHz up to 26,3 GHz.

A commercial sample was used, measuring at room temperature. The absorption lines frequencies were measured with markers produced by a frequency standard, driven by a 10 MHz quartz crystal oscillator, which was kept in a thermostatic oven. The frequency standard was calibrated with very well measured lines. In order to determine in unambiguous form the harmonic of the fundamental standard frequency, that fits a line, the arrangement described in [5] for coarse measurements was used. But for stability reasons some improvements have been made. A variable frequency oscillator inserted in the frequency standard was substituted by a signal generator, which feeds with 150 MHz an amplifying-tripler stage of 450 MHz that finally drives a 1N26 crystal multiplier. Tuning slightly both signal generator and tripler stage it is possible to have in general two settings of markers over the line or if one is lacking, a coarse measurement with a wavemeter is possible without any loss of accuracy in the deter-

<sup>\*</sup> This work is being performed at the Institute of physics of the university of Stockholm in use of a fellowship of the university of Buenos Aires.

mination of the harmonic. With this molecule it is possible to repeat frequency measurements of medium and strong lines within the range of  $\pm$  0,2 MGHz, due to the rather large half-width of the lines, that reaches in this case several MGHz.

# INTERPRETATION AND DISCUSSION

The observed spectra is rather complicated (almost 300 lines measured) due to:

- 1) the molecule is heavy, then high rotational J value lines and transitions in low vibrational states are observable,
- 2) the quadrupole electric moment of the chlorine nucleus causes an hyperfine splitting,
- 3) the presence of two isotopes: Cl35 and Cl37, increases the number of observable lines.

Yet it has not been possible to resolve Stark patterns, of low J value lines, then another kind of analysis was chosen. With the internuclear lengths and bond angles given in [2, 3], a planar model of the molecule was built. Following Bragg and Golden's theory [6], a careful study, on the model, of the relative quadrupole splitting was made, which is defined by

$$\frac{E_Q}{eqQ} = f(J, \kappa) \cdot Y(I, J, F)$$

where  $E_Q$  is the quadrupole perturbation energy and eqQ is the quadrupole coupling constant. It was assumed that the field around the Cl-nucleus has rotational symmetry around the C-Cl bond. J is the rotational quantum number,  $\kappa$  the asymmetry parameter and Y(I, J, F) the Casimir's function. Using the Cl<sup>35</sup> isotope, it showed that it is possible "to pick up" two series of  $\mu_b$  type, R-branch rotational lines with high relative intensities [7], somehow equidistants and of low relative quadrupole splitting. These two series are characterized by  $J, K_{-1}$  and  $K_{+1}$  numbers connected in a form that

$$J_{0,J} \to J + 1_{1,J+1}$$
  
$$J_{1,J} \to J + 1_{0,J+1} .$$

They were found and a first interpretation is given in Table I.

	TABLE	I.
1—	-3 C <sub>6</sub> H <sub>4</sub>	Cl³5F.

Observed lines MHz	Interpretation	Observed lines MHz	Interpretation
12.106,9 13.621.0 15.173.5 16.826.8 18.502.4 20.210.9 21.939.0 23.675.2 25.410.4	50.5 - 61.6 $60.6 - 71.7$ $70.7 - 81.8$ $80.8 - 91.9$ $90.9 - 101.10$ $100.10 - 111.11$ $110.11 - 121.12$ $120.12 - 131.13$ $130.13 - 141.14$	12.662.6 14.596.7 16.465.9 18.289.8 20.084.6 21.867.5 23.627.5 25.389.5	$\begin{array}{c} 61.6 \ -\ 70.7 \\ 71.7 \ -\ 80.8 \\ 81.8 \ -\ 90.9 \\ 91.9 \ -\ 100.10 \\ 101.10 \ -\ 110.11 \\ 111.11 \ -\ 120.12 \\ 121.12 \ -\ 130.13 \\ 131.13 \ -\ 140.14 \\ \end{array}$

The observed low *J*-value lines from Table I were found to be unresolvable from the point of view of the quadrupole splitting, like it was predicted by the theory. The calculations were carried out using electronic computing machine methods for rigid asymmetric rotor levels and the assignments were made assuming planarity of the molecule. But it must be noticed that they should be considered as an attempt and not definitive.

The comparison between observed and calculated Stark effect components gives evidence of the correctness of the assignments; the calculations were carried following Golden and Wilson's theory for Stark effect of a rigid asymmetric rotor [8], assuming that the dipole electric moment component on the b-axis is much greater than on the a-axis, using the well known convention for principal inertia moments that  $I_a < I_b < I_c$ . With the energy levels calculated from the supposed molecular model for the four first transitions in both series, it is possible to explain satisfactorily the inversion of the Stark effect on the lines  $8_{0.8}-9_{1.9}$  and  $9_{1.9}-10_{0.10}$ , like is showed in table II.

The measurements shall be continued down to  $8.5~\mathrm{GHz}$  in the X-band to try to find lower J value transitions of both series. Calculations are carried on the way to give accurate values of the rotational constants and the "inertia defect".

I wish to express my thanks to Laborator Gunnar Erlandsson of the university of Stockholm for valuable advice and help in performing this work.

TABLE II.

Transition	Observed Stark ef.	Calculat. Stark ef.
50.5— 61.6	Regular	Regular
60.6— 71.7	Regular	Regular
70.7— 81.8	Regular	Regular
80.8— 91.9	Inverted	Inverted
61.6— 70.7	Inverted	Inverted
71.7— 80.8	Inverted	Inverted
81.8— 90.9	Inverted	Inverted
91.9—100.10	Regular	Regular

## REFERENCES

- 1. McCulloh, K. E. and G. F. Pollnow, J. Chem. Phys., 22, 1144 (1954).
- 2. ERLANDSSON, G., Arkio f. Fysik, 9, 399 (1955).
- 3. Bak., Christensen, Hansen-Nygaard and Tannenbaum, J. Chem. Phys., 26, 134 (1957).
- 4. SELÉN, H., Arkio f. Fysik, 13, 81 (1957).
- 5. Erlandsson, G. and H. Selén, Arkiv f. Fysik, 11, 391 (1956).
- 6. Bragg, J. K. and S. Golden, Phys. Rev., 75, 735 (1949).
- 7. Schwendeman and Laurie, Tables of line Strengths for Rotat. Trans. of Asym. Rot. Molecules, Pergamon Press.
- 8. Golden, S. and E. Bright Wilson, Jr., J. Chem. Phys., 16, 669 (1948).
- 9. Gordy, Smith and Trambarulo, *Microwave Spectroscopy*, John Wiley and sons (1953).
- 10. Townes and Schawlow, Microwave Spectroscopy, Mc-Graw-Hill Book Company (1955).