

<b>Zeitschrift:</b>	Archives des sciences [1948-1980]
<b>Herausgeber:</b>	Société de Physique et d'Histoire Naturelle de Genève
<b>Band:</b>	28 (1975)
<b>Heft:</b>	2
<b>Artikel:</b>	Addendum : evolutionary trends in the Arkhangelskiellaceae (calcareous nannoplankton) of the Upper Cretaceous of Central Oman, SE Arabia
<b>Autor:</b>	Lauer, G.
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-739803">https://doi.org/10.5169/seals-739803</a>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 09.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

**ADDENDUM****EVOLUTIONARY TRENDS IN THE ARKHANGELSKIELLACEAE  
(CALCAREOUS NANNOPLANKTON) OF THE UPPER CRETACEOUS  
OF CENTRAL OMAN, SE ARABIA****G. LAUER<sup>1</sup>**

The study was initiated at the request of the oil company P.D. (Oman) Ltd. to arrive at a detailed chronostratigraphic subdivision of the Late Cretaceous Fiqa Formation in the Oman, SE Arabia. This Formation covers the upper part of the *Globotruncana carinata* to the lower part of the *Globotruncana calcarata* planktonic foraminiferal zones and is therefore of Late Santonian to Late Campanian age. The dark calcareous shales of the Fiqa Formation which can be up to 5000' thick were studied in 25 cored well-intervals distributed over an area of approximately 90.000 km<sup>2</sup>.

**1. SYSTEMATICS**

In the evolution of the family Arkhangelskiellaceae four separate lineages are recognised:

**1. *Aspidolithus* NOËL 1969**

According to the type species *Aspidolithus angustus* NOËL 1969 this organ genus applies to types of Arkhangelskiellaceae with two major distal cycles of elements and three layers in side view. The central area is occupied by a plate. The evolution shows the following morphological trends in the investigated interval:

- a) increase of size
- b) increasing difference in diameter of the three layers (the early types show three layers of the same size; in the later types the distal layer is larger).
- c) reduction in size of central area
- d) reduction in number of perforations

---

<sup>1</sup> P.D. (Oman) Ltd., present address: Koninklijke/Shell Exploratie en Produktie Laboratorium, Volmerlaan 6, Rijswijk (ZH), The Netherlands.

## 2. *Broinsonia* BUKRY 1969

The organ genus is defined by a cross structure in the central area, as indicated by the type species *Broinsonia dentata* BUKRY 1969. The studied time interval does not allow the definition of evolutionary trends. However, a variety of types is recognised, which are combined under the preliminary term of *Broinsonia dentata* group.

## 3. *Arkhangelskiella* VEKSHINA 1959

A large number of morphological variations are condensed in two groups of species. The *Arkhangelskiella* group 1 includes types with four perforations along the short axis of the central area. Six or more perforations are present along the short axis in the central area in the *Arkhangelskiella* group 2. Besides a variety of evolutionary trends, mainly concerning the distribution of the central perforations a clear development of the marginal area is recognisable in both lineages. The early stages have three layers in side view and two cycles on the distal side which indicates an evolutionary connection with *Aspidolithus*. The later stages show four layers in side view and only one cycle at the distal side. The outer cycle has become superimposed by the inner cycle, which finally forms a fourth layer.

## 4. *Gartnerago* BUKRY 1969


The morphology of this organ genus suggests major differences from the three closely related organ genera mentioned before. Within the Santonian-Campanian interval two evolutionary trends are observed:

- a) decrease in size
- b) decrease in number of central perforations. The three types are distinguished by the number of perforations along the short axis of the central area;
  - Gartnerago* sp. 1: 8 perforations
  - Gartnerago* sp. 2: 6 perforations
  - Gartnerago* sp. 3: 4 perforations

## II. STRATIGRAPHY

With these evolutionary lineages a sixfold subdivision has been achieved in the Fiqa Formation of Central Oman. The zonation is defined by morphological criteria which are recognisable under the optical microscope.

On the graph the following evolutionary events are indicated from bottom to top:



Zone F. No lower limit has been defined.

Zone E. First occurrence of the *Arkhangelskiella* group 2. Furthermore only types of Arkhangelskiellaceae showing two major cycles at the distal side are present.

Zone D. First occurrence of *Gartnerago* sp. 2. As a second aspect the first representatives with four layers appear in the *Arkhangelskiella* group 1. Also in this case the absence of a differentiation in the marginal area is indicative.

Zone C. First occurrence of *Aspidolithus* sp. 1, showing three layers of the same size. In addition the first types with four layers appear in the *Arkhangelskiella* group 2. Consequently an undifferentiated marginal area is visible in distal and proximal views.

Zone B. First occurrence of large *Aspidolithus*, showing different diameters of the three layers in side view (*Aspidolithus* sp. 3). In case of proximal or distal views the morphology can be determined by focussing to the appropriate level.

Zone A. First occurrence of large *Aspidolithus* without central perforations, and a very small elongated area (*Aspidolithus* sp. 5).

The frequency of the morphological types and their intermediates is shown in the graph by the lateral extent of the shaded areas. The recognition of the submicroscopical morphology is based on the study of 150 SEM photographs supplied by Prof. Dr. F. R. Allemann and Dr. W. Grün (now at Ö.M.V.A.G., Vienna) from the Geological Institute in Bern (Switzerland). Permission by P.D. (Oman) Ltd., to publish this paper is gratefully acknowledged.

*Authors' addresses :*

Laboratoire de Géologie du Muséum  
61, rue de Buffon  
75005 Paris (France)

Eidg. Technische Hochschule Zürich  
Geologisches Institut  
Soneggstrasse 5  
CH-8006 Zürich (Switzerland)