

**Zeitschrift:** Archives des sciences et compte rendu des séances de la Société  
**Herausgeber:** Société de Physique et d'Histoire Naturelle de Genève  
**Band:** 35 (1982)  
**Heft:** 1

**Artikel:** New and little known species of Aphodius illig. (Coleoptera, Scarabaeidae, Aphodiinae)  
**Autor:** Stebnicka, Z.  
**DOI:** <https://doi.org/10.5169/seals-740550>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 02.05.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

NEW AND LITTLE KNOWN  
SPECIES OF APHODIUS ILLIG.  
(Coleoptera, Scarabaeidae, Aphodiinae)

BY

Z. STEBNICKA \*

With 9 figures

---

ABSTRACT

*Aphodius (Agrilinus) pseudostorkani* from China is described as new and the affinities with related species are discussed. Complementary description for hitherto unknown male of *A. donghariensis* Steb. and explanation of the taxonomic position of two other species are given.

In 1980 I had the chance to study material of the subfamily Aphodiinae from the Petrovitz' collection, kept in the Museum of Natural History in Genève. The examination of the type-specimens of a number of species, necessitates the erection of new one (formerly identified as *Aphodius storkani* Balth.) and determination the synonymies of *A. botulus* Balth. and *A. varians* Duft. Among unidentified specimens from Taiwan, collected by J.-S. Klapperich I found the hitherto unknown male of *A. donghariensis* Steb., described in 1973 from North Korea.

The specimens examined are deposited in the collection of the Museum of Natural History in Genève and secondly, in the collection of the Institute of Systematic and Experimental Zoology (ISEZ), Polish Academy of Sciences in Kraków.

I wish to thank Dr I. Löbl and Dr C. Besuchet for making this material available to study.


***Aphodius (Agrilinus) pseudostorkani* nov. spec. (Figs. 1-3)**

Holotype: male, China, W. Sichuan, Sankiangkou, Balang, 7.VIII.1934, leg. Friedrich (Geneva Museum).

Paratype: male, the same data as holotype (ISEZ).

---

\* Institute of Systematic and Experimental Zoology, Polish Academy of Sciences, 31 016 Kraków, Sławkowska 17, Poland.



FIGS. 1-6. — 1-3: *Aphodius (Agrilinus) pseudostorkani* nov. spec., 1: epipharynx, 2: aedeagus, lateral view, 3: aedeagus, dorsal view. 4-6: *A. (A.) storkani* Balzh., 4: aedeagus, lateral view, 5: aedeagus, dorsal view, 6: epipharynx.

Description. Length 6.0-6.2 mm, greatest width 2.8-3.0 mm. Oblong-oval, convex, moderately shining, piceous, apex of elytra brownish. Head moderately convex, finely reflexed clypeal margin subdentate each side of wide, moderately deep median emargination, sides nearly straight to small, rounded genae; frontal suture with three, very feebly marked transverse tubercles, clypeal surface granulate with indistinct transverse ridge at middle, front moderately punctate. Pronotum convex, anterior angles widely rounded and reflexed, posterior angles obtuse, the slightly

arcuate sides feebly sinuate, base margined; surface with mixed punctures, everywhere with fine, rather evenly distributed punctures separated by one to three diameters and coarse punctures separated by one or two diameters, more close on the sides. Scutellum triangular, finely punctate. Elytra slightly widened behind the middle, convex, humeri finely dentate; striae narrow, moderately deep, striae punctures slightly crenating the inner margins of the weakly convex intervals; surface moderately shining with scattered, minute punctures, sides and apex of elytra finely shagreened. Metasternum shining, midline fine, moderately impressed, the punctures coarse, not closely distributed; abdominal sterna alutaceous, coarsely punctate and piliferous. Middle and hind femora shining with a few coarse punctures bearing yellow hairs. Apical spur of anterior tibia straight, sharply pointed; apical setae of middle and hind tibia short, equal in length; bottom spur of middle tibia short with hooked tip; first segment of posterior tarsus longer than the upper spur and shorter than following three segments combined.

Female unknown.

Affinity: *Aphodius pseudostorkani* n. sp. is very similar to *A. storkani* Balth. described from China — Sichuan (holotype male in Geneva Museum). The differences between the species mentioned are as follows:

<i>Aphodius (Agrilinus) storkani</i> Balth.	<i>A. (A.) pseudostorkani</i> n. sp.
Sides of clypeus toward genae slightly arcuate	Sides of clypeus toward genae nearly straight
Disc of pronotum with equal punctures, moderate in size	Disc of pronotum with mixed, very fine and coarse punctures
Humeri strongly, sharply dentate	Humeri finely, obtusely dentate
Male aedeagus as in figures 4-5	Male aedeagus as in figures 2-3
Epipharynx (Fig. 6): Acanthoparia with moderately long and moderately thick setae nearly the same size as those of chaetoparia; 8-10 bristles of chaetopedia somewhat thicker and shorter than those of chaetoparia; the remaining setae of paria and acroparia moderately long and thin.	Epipharynx (Fig. 1): Acanthoparia with moderately long and moderately thick setae the same size as those of chaetoparia; 20-22 bristles of chaetopedia the same thickness as those of chaetoparia, somewhat shorter; the remaining setae of paria and acroparia moderately long and thin.


***Aphodius (Teuchestes) donghariensis* Steb.**

*Aphodius donghariensis* Stebnicka, 1973: 119-120; 1980: 231.

Material: two males, Taiwan, Fenchihu (1400 m), IV-VI 1977, leg. J.-S. Klap-perich.

Complementary description. Length 5.8-6.0 mm, greatest width 3.0-3.3 mm. Male head wider than in female, scutellum very finely, indistinctly punctate. Apical spur of fore tibia wide, flattened, strongly bent inward and slightly downward; bottom spur of middle tibia short, truncate at apex. Aedeagus as in figures 8-9.

Epipharynx: Anterior part of acanthoparia with a few short and thick bristles, shorter but the same thickness as those of chaetoparia; 15-17 bristles of chaetopedia unequal in length, the remaining setae of paria and acroparia moderately long and moderately thin.



FIGS. 7-9. — *Aphodius (Teuchestes) donghariensis* Steb., 7: epipharynx, 8: aedeagus, lateral view, 9: aedeagus, dorsal view.

***Aphodius (Phalacrothous) botulus* Balth.**

*Aphodius (Paragolius) botulus* Balthasar, 1945: 43; 1964: 281. Locus typicus: Manchuria, Harbin.

*Aphodius (Orodalus) naraensis* Nakane, 1956: 120; BALTHASAR 1964: 201; STEBNICKA 1980: 239-240 (**syn. nov.**). Locus typicus: Japan, Honshu, Nara; (North Korea).

*Aphodius (Amidorus) manschuriensis* Petrovitz, 1958: 138; BALTHASAR 1964: 281.  
Locus typicus: Manchuria, Harbin.

Remarks. I have examined the holotype of *Aphodius botulus* in the Balthasar's collection in Prague and the holotype of *A. manschuriensis* Petr. in Geneva. Type-specimen of *A. naraensis* Nak. was not verified, however, the series of specimens from Korea (STEBNICKA 1980) was determined on the base of single specimen from Japan, compared with holotype.

*A. botulus* has been described by BALTHASAR (1945) in a new subgenus *Paragolius*. The establishment of new subgenus was not justified, since the complex of the features of mentioned species indicates its affiliation to the species-group *Phalacrothous* Motsch. The representatives of these species are difficult to define and to separate in many cases, by reason of great individual variability respecting the shape and punctuation of body and color of elytra.

The male aedeagus of *A. (Phalacrothous) botulus* Balth. (= *A. (Ph.) naraensis* Nak., published drawings: STEBNICKA 1980) is quite similar to this of *A. (Ph.) quadrimaculatus* L.), (published drawings: STEBNICKA 1976).

#### ***Aphodius (Nialus) varians* Duft.**

= *Aphodius (Nialus) rugosopunctatus* Petrovitz,  
1971: 567-568 (syn. nov.).

Material of *rugosopunctatus*. The holotype examined: Turkey, Vilayet Mersin (İçel), Namrun (Geneva Museum). Three specimens: Poland, distr. Warsaw, Swider, 25.VI.1906, leg. Makólski; distr. Kraków, Sikornik hill, 18.VI.1936, leg. Mazur (ISEZ).

Remarks. *Aphodius rugosopunctatus* Petr. was originally described on the base of external morphology of a single male specimen. *A. varians* Duft. is well known, widespread, polymorphic species composed of populations which often are not contiguous. The examined morphology of male copulatory organs and epipharynx is identical in both variants, whilst the upper side of body of «*rugosopunctatus*» individuals shows more pronounced punctuation, as compared to the typical specimens of *varians* Duft. The latter feature, however, seems to fall within the range of individual, very advanced variability of *A. varians* Duft. The specimens similar to «*rugosopunctatus*» are evidently very rare; moreover, it is not excluded that these individuals appear from time to time as mutations in different populations of *A. varians* Duft.

## REFERENCES

- BALTHASAR, V. (1945). Quattuor generis *Aphodius* Ill. subgenera nova. *Čas. čsl. Spol. ent.* 42: 40-44.
- (1964). Monographie der Scarabaeidae und Aphodiidae der palaearktischen und orientalischen Region. Coleoptera: Lamellicornia 3, 652 pp., 224 figs.
- NAKANE, T. (1956). Neue Aphodiinen aus Japan. *Insecta matsum.* 20: 119-124.
- PETROVITZ, R. (1958). Neue asiatische Aphodiusarten (Col., Scarab.). *Ent. Arb. Mus. Georg Frey* 9: 131-139.
- (1971). Ergebnisse zoologischer Sammelreisen in der Türkei. Lamellicornia, Coleoptera. *Annl. naturh. Mus. Wien* 75: 565-589.
- STEBNICKA, Z. (1973). Neue Scarabaeidenarten (Coleoptera) aus der Koreanischen Volksdemokratischen Republik. *Bull. Acad. pol. Sci. Sér. Sci. biol.* 21: 117-121, figs.
- (1976). Keys for the Identification of Polish Insects. Part 19: Coleoptera, No. 28a — Scarabaeidae laparosticti. 139 pp., 500 figs.
- (1980). Scarabaeoidea (Coleoptera) of the Democratic People's Republic of Korea. *Acta zool. cracov.* 24: 191-298, 232 figs.