

Zeitschrift: Archives des sciences et compte rendu des séances de la Société
Herausgeber: Société de Physique et d'Histoire Naturelle de Genève
Band: 41 (1988)
Heft: 3: Archives des sciences

Artikel: New and interesting mites from the Geneva Museum LXVI : two new Scutacarus gros : 1845 species from Sumatra (Indonesia) (Acari, Tarsonemina)
Autor: Mahunka, S. / Mahunka-Papp, L.
DOI: <https://doi.org/10.5169/seals-740401>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 05.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

**NEW AND INTERESTING MITES
FROM THE GENEVA MUSEUM LXVI.
TWO NEW SCUTACARUS GROS, 1845 SPECIES
FROM SUMATRA (INDONESIA) (ACARI: TARSONEMINA)**

BY

S. MAHUNKA * and L. MAHUNKA-PAPP *

With 18 figures

ABSTRACT

New and interesting mites from the Geneva Museum LXVI. Two new Scutacarus Gros, 1845 species from Sumatra (Indonesia) (Acari: Tarsonemina). — Three species are identified from Sumatra, two of them are new for science: *Scutacarus batak* sp. n. and *S. sumatrana* sp. n.

INTRODUCTION

Soil samples taken during the 1985 expedition (see also MAHUNKA 1989) of the Geneva Natural History Museum to South-East Asia (Dr. B. Hauser and Dr. C. Lienhard), yielded some *Scutacaridae* species. Three species have been identified, two of them proved to be new to science.

LIST OF LOCALITIES

Sum-85/30: **SUMATRA** (Sumatera Utara: Simalungun): sur la route en provenance de Pematangsiantar, 18 km avant Prapat, lieu-dit «Ainuli», forêt dans la chaîne montagneuse Bukit Parasat, derrière la station «Holzweg Nr. 2 du Dr. Diehl», forêt primaire autour de l'école

* Zoological Department, Hungarian Natural History Museum, Baross utca 13, H-1088 Budapest, Hungary.

- forestière, 1000 m, prélèvement de sol dans les angles formés par les contreforts de grands arbres; 14.XI.1985, leg. B. H. (B)*
- Sum-85/39: SUMATRA (Sumatera Utara: Deli Serdang): Sibolangit sur la route de Medan à Brastagi, Jardin botanique, prélèvement de sol dans les angles formés par les contreforts de grands arbres, 520 m; 18.XI.1985, leg. B. H. (B)
- Sum-85/47: SUMATRA (Sumatera Utara: Deli Serdang): forêt de *Pinus merkusii* près de la route de Brastagi à Sibolangit, 1400 m; prélèvement de sol sous *Pinus merkusii*; 19.XI.1985, leg. B. H. (B)

LIST OF IDENTIFIED SPECIES

Imparipes burckhardti Mahunka et Mahunka-Papp, 1988

Locality: Sum-85/47: 2 specimens.

Scutacarus batak sp. n.

Locality: Sum-85/39.

Scutacarus sumatranaus sp. n.

Locality: Sum-85-30.

DESCRIPTION OF SPECIES


Scutacarus batak sp. n.

Measurements : Length: 169-180 µm, width: 140-155 µm.

Dorsal side (Fig. 1): Clypeus very large, margin of clypeus very broad anteriorly. Both pairs of setae arising on it, equal in length (42 µm). Among the other setae h_2 short, e minute, f (100 µm) the longest, d (69 µm) and h_1 (64 µm) nearly equal in length. In posteromarginal position three setae observable, but p_2 much shorter and thinner than p_1 , p_3 minute. Setae p_1 resembling a Christmas tree.


Ventral side (Fig. 2): Apodemes of both sternal plates well observable. Posterior border of epimeral region strong, ap. 2 waved, slightly thinner than ap. 3. Anterior margin of posterior sternal plate strongly convex anteriorly but slightly concave medially. Anterolateral spine long and narrow (Fig. 6). Every setae of epimeral region long, and all, excepting 2b, well ciliate. Cilia thin, conspicuously long on setae 3a and 3b. Setae 4a arising very far anterior to setae 4b, their distance scarcely shorter than distance between setae 4b.

*(B): extraction par appareil BERLESE à Pematangsiantar (Sumatra).



FIGS 1-9.

Scutacarus batak sp. n. — 1: body in dorsal view, 2: body in ventral view, 3: leg II, 4: claw of leg I, 5: trichobothrium, 6: posterior sternal plate in lateral view, 7: leg IV, 8: leg III in ventral view, 9: leg I.



FIGS 10-18.

Scutacarus sumatranaus sp. n. — 10: body in dorsal view, 11: body in ventral view, 12: leg IV, 13: posterior part of coxisternal region, 14: trichobothrium, 15: leg III, 16: leg I, 17: claw of leg I, 18: leg II.

L e g s : Claw of leg I (Fig. 9) sitting on a long peduncle, a separate basal scale well observable. Basal tubercle of seta *d* very long, seta *d* slightly shorter than *dT*. Solenidium ψ_1 and ψ_2 nearly equal in length. Tarsi of legs II (Fig. 3) and III (Fig. 8) elongate, more than three times longer than their basal diameter. Setae *ld''* on both joints thickened, spiniform, not ciliate. Trochanter of leg IV (Fig. 7) without dorsal spine, but a well developed spur on its ventral (inner) side present. Femur divided into two parts by a line, seta *dF* strong, erect. Setae *d* on tibiotarsus fine and short, setae *ld'*, the longest of all.

M a t e r i a l e x a m i n e d : Holotype: Sum-85/39; 1 paratype: from the same sample. Holotype: MHNG *, paratype (1213-PT-86): HNHM **.

R e m a r k s : The new species is well characterized by the insertion points of setae *4a-4b* and the shape of leg IV.

Scutacarus sumatranus sp. n.

M e a s u r e m e n t s : Length: 316 μm , width: 257 μm .

D o r s a l s i d e (Fig. 10): Body strongly convex in lateral view. Clypeus large, but not much broader than the other segments. Clypeal margin also wide. Inner pairs of clypeal setae very short (13 μm) and fine, outer pairs much thicker. Ratio of some dorsal setae: $c_1 = d < e$ (22 μm) $< f \equiv c_2 < h_2$ (44 μm) $< h_1$ (49 μm). Setae *p₃* the longest of all (52 μm), *ps₁* well dilate and bilaterally ciliate, pennate; *ps₂* longer than *ps₁*, but simple.

V e n t r a l s i d e (Fig. 11): All apodemes well developed, a very strong transversal band (ap. 5) also existing between the trochanters of legs IV. Posterior margin of this region waved medially (Fig. 13). All epimeral setae — excepting *2b* — very long and with long cilia, but the density and the length of the cilia are quite different, e.g. setae *1a* more densely ciliate than *1b*, or the cilia of setae *4a* much longer than those of *4b*. Setae *4a* arising very near to *4b*.

L e g s : Claw of leg I (Fig. 16) sitting on a long peduncle, its apex long, thin. A well-developed, sharply pointed basal scale (Fig. 17) also observable. All setae of tibiotarsus of leg I very long, seta *dT* (83 μm) longer than seta *d* (62 μm), but setae *ldT* the longest of all (122 μm). Solenidia also long, ψ_1 the thickest. Tarsi of legs II (Fig. 18) and III (Fig. 15) elongate, setae *d* on both tarsi ciliate. Femur of leg IV (Fig. 12) long, its basal margin convex. Tibiotarsus also elongate, nearly three times

* MHNG = deposited in the Muséum d'histoire naturelle, Genève.

** HNHM = deposited in the Hungarian Natural History Museum, Budapest, with identification number of the specimens in the Collection of Arachnida.

longer than its basal diameter. Setae dT much longer than the nearly equally long setae d and $1d'_{2}$.

M a t e r i a l e x a m i n e d : Holotype: Sum-85/30: MHNG.

R e m a r k s : The new species belongs to the *Scutacarus sphaeroideus* Karafiat, 1959 group. It is well distinguished from all members of this group by the waved posterior margin of the coxisternal plate and the shape of the group of setae ps .

REFERENCES

- MAHUNKA, S. 1989. New and interesting mites from the Geneva Museum LXV. Oribatids from Sumatra (Indonesia) I. *Revue suisse Zool.* 96: in print.
- MAHUNKA, S. and L. MAHUNKA-PAPP. 1988. Neue und interessante Milben aus dem Genfer Museum LIX. Hauseripes hungarorum gen. n., sp. n. and some other new Tarsonemina (Acari) from Sabah (East Malaysia). *Revue suisse Zool.* 95: 581-594.