Zeitschrift: Archives des sciences et compte rendu des séances de la Société

Herausgeber: Société de Physique et d'Histoire Naturelle de Genève

Band: 42 (1989)

Heft: 1: Archives des Sciences

Artikel: Polarity: from dipoles to biopolarizations. II. Addenda and indexes

Autor: Turian, Gilbert

DOI: https://doi.org/10.5169/seals-740081

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 21.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

シィー イレロ

POLARITY

FROM DIPOLES TO BIOPOLARIZATIONS

II. ADDENDA AND INDEXES

by

Gilbert Turian

ETHZÜBICH

1 8. Jan. 10.)

BIBLIOTHEK

Ce travail est la suite et le complément de celui intitulé POLARITY, paru en 1989 dans le volume 42, fascicule 1, des *Archives des Sciences*.

La numérotation des pages poursuit celle dudit travail. Nous conseillons donc à nos abonnés de classer ce supplément à la fin du fascicule 1, vol. 42.

This paper supplements the review entitled POLARITY (Arch. Sci. 42: 1-323, 1989) and should be classified at the end of vol. 42, fasc. 1.

Société de Physique et d'Histoire naturelle Genève 1990



POLARITY FROM DIPOLES TO BIOPOLARIZATIONS

II. ADDENDA AND INDEXES

BY

Gilbert TURIAN*

Polarity is a problem of wide interdisciplinary interest that we have attempted to survey in its widest span from its atomic to its embryogenic, plant and animal levels in the Archives of 1989 reprinted as book I (Turian, 1989).

Primeval polarity is bipolar, founded on the separation of two equal but opposite electric charges. Consequently, even apolar molecules are intrinsically electrically polarized but with a symmetrical distribution of their opposite (+ and -) electric charges and therefore they lack in electric polar moment. Similarly, apolar morphological biostructures are examplified by spherical cells (certain eggs, etc.), initially deprived of heterogeneously distributed components, and which being identical with their mirror image can be also considered as achiral.

The whole universe is electrically neutral and, by necessity, contains rigorously equal numbers of opposite electric charges (10⁴⁰ of protons and of electrons, see Souriau in Brack *et al.*, 1989) even though it is filled with electric dipoles from the minute water molecules to giant cosmic dipoles, a basic requirement for its physicochemical and biological functionings. However, in its wider expression, polarity spans not only pure electric and magnetic phenomena but also chemostructural (chiral), biomolecular (cytoskeletal elements) and spatio-temporal developmental processes. Our survey had therefore to encompass them in their whole span from monopoles to multipoles as following:

1) monopoles, electric (+ or -) or magnetic (still elusive north or south isolated poles) as well as homochirals (l- or d-enantiomers) and monopolar, elongating biostructures such as microfilaments (actin), microtubules (tubulins), multinucleate cells such as hyphae and neurites;

^{*} Laboratoire de Microbiologie générale, Sciences III, Université de Genève.

- 2) dipoles, basically electric (+ and -) or magnetic (north and south poles), but also heterochirals (l + d-enantiomers) as well as morphogenetic homodipoles in the twice-budded or -germinated yeasts or fungal spores and heterodipoles in the developing eggs of plants and animals;
- 3) tripoles, electric (+ + as in thunderclouds, see addendum) or morphogenetic as in iris flowers!
- 4) quadrupoles, electric (radio-frequency electric traps and nuclear coupling, see addendum) or morphogenetic as in four- (multi-) budded yeasts and germinated fungal spores;
- 5) multipoles as exhibited by cells such as amoebae or fungal spores outgrowing n (>4) pseudopodia or germ tubes, respectively.

During the second half of 1989 and first trimester of 1990, we have noticed a few omitted significant papers as well as newly published ones, related to dipoles and biopolarities. We have registered them below by following the sequence of the eight preceding chapters and, parallely, added two subject and taxonomic indexes. Their item entries cover the main book (I, 1989) and these first addenda (II, 1990).

I. ATOMIC POLARIZATIONS

A. ORIGINS

A fourth state of matter is the plasma state which is formed when a gas is heated to such high temperatures that it becomes partly or fully ionized: "electrons are torn off the atoms in the gas, leaving a stream of negatively charged free electrons and positively charged ions" (Peratt, 1990). Suggestively, the term plasma proposed in 1932 by Langmuir evokes "the unstable almost lifelike behavior of the ionized material". According to plasma cosmology, "the universe has been and remains a veritable sea of charged particles interlaced with complex magnetic fields and electric currents. Many among the cosmologists therefore conclude with Peratt (1990) that "the universe may have evolved not with the Big Bang but from a vast sea of plasma". However, the theory of primordial explosion and of the shaping of the universe by gravitational rather than by electromagnetic forces keeps strong proponents (Rees, 1990).

In the evolutive perspective from the inert to the living matter, the atom of hydrogen (H) could be viewed as forming "le couple divin" (Turian, 1990) displaying the electron, mobile as a male cell around the passively "courted" proton, as female cell, a most fertile association indeed concretized in the bioenergetics through the ATP-generating redox scale $(2H^+ + 2e^- + O = H_2O)$, see I, IV.B.2.c).

B. SYMMETRY - POLARITY

The whole world appears to be chirally asymmetric from the scale of elementary particles upward. This leads Hegstrom and Kondepudi (1990) to ask the questions, as we did (I) for related polarities "How do the asymmetries arise? Are chiral symmetries at one level linked to those at another, or are they independent?"

Chiral asymmetry must therefore be first studied at the scale of elementary particles. Indeed, there is symmetry within an atom only when it is regarded as governed by the electromagnetic force and its associated property of conservation of parity. The additional weak force (involving W^+ and W^- gauge bosons) gives rise to a violation of parity and consequently an asymmetry between the electrons and the nucleus in the atom (Bouchiat and Pottier, 1984). Chiral asymmetry at the subatomic level is thus fundamentally connected to parity nonconservation. One result of this asymmetry is that nuclear β decay, which is governed by the W force, produces mostly left-handed electrons. Consequently, electrons of matter are polarized with a left helicoidal coil while positrons of antimatter are right-directed. However, such chiral

effects of the electroweak W and Z charges leading to a distinction between left and right chirals are strictly valid only when the electrons are travelling at the high energies near the speed of light (Hegstrom and Kondepudi, 1990).

An important consequence of chiral asymmetry at the *subatomic* level is that it causes a chiral asymmetry at the higher level of *atoms*: under the influence of the Z weak force, the electron orbit becomes a right-handed helix in the vicinity of the nucleus. However, the asymmetric Z force is so small that its effects on the chemical properties of *molecules* has not (yet) been observed (Hegstrom and Kondepudi, 1990). That such a mechanism affecting the production rate of L- and D-amino acids can indeed exist in nonequilibrium chemical systems was shown theoretically by Kondepudi and Nelson (1985, see I).

The problem of equivalence which has been upheld about left and right (see II.D) also arises "with respect to positive and negative electricity" as commented by Weyl (1952) in his book entitled "Symmetry" in which he also discussed relationships between quantum mechanics and symmetry. This author also assumed that "the primary polarity as well as the subsequent *bilateral* symmetry come about by external factors actualizing potentialities inherent in the genetic constitution" (see VII-VIII in I).

As already expressed by Pierre Curie "symmetric systems behave in a symmetric fashion". However, such Curie's principle is contradicted by the occurrence of spontaneous symmetry breaking which occurs when a perfectly symmetric system takes up a state with less symmetry (Field and Richardson, 1989). An example of the phenomenon is the change of form produced by compression of a cylindrical shell initially endowed with a perfectly *circular* symmetry.

The principle of "cosmologie symétrique" has been further discussed by Brack et al. (1989) in relationship with the equivalence between matter (proton + electron) and antimatter (antiproton + antielectron). Among previous books concerned with the principle of symmetry there are those cited by Weyl (1935), namely Jaeger (1917) and Hambidge (Dynamic Symmetry, 1920), completed by Jaeger (1925) and, more recently, those by Nicolle (1950) and Caillois (1973) as well as Hargittai and Hargittai (1986).

C. ELECTRIC BIPOLARIZATION

2) Electric dipoles

The hydrogen atom (H) can be considered as the primordial electric dipole when we consider that its electron or unit of negative charge is probabilistically positioned on a peripheral orbit around the positive proton according to the classical image of a planet circling the sun (Fig. 1B, in I). However, when the atom is placed under

strong stimuli such as a constant magnetic field or exposed to electromagnetic radiation in the form of microwaves, either of these strong stimuli disturbs the orbit of the electron and pushes it into chaotic, unpredictable motion. Eventually, the electron atom is ionized, i.e. its electron has so much energy that the pull of the proton can no longer hold it, and the electron is torn away. According to quantum mechanics, the electron is not considered as a particle orbiting the proton, but as a rather nebulous "wave packet". Ionization high energy will delocalize the wave packet, namely "the electron will become "spread out" over several energy levels", an event corresponding to "the chaos in the classical motion of the electron" (Pool, 1989).

Protons and neutrons, the two types of nucleons, can be examined "by observing electron or muon scattered off them with a large transfer of momentum to one of their constituent particles or partons" (Roberts, 1990). As for the proton, its simplest properties are dependent on the three valence quarks, two "up" (u^+) and one "down" (d^-) (see I, I.C.2), each of which carries a spin of 1/2. These are polarized so that the u^+ quarks contribute 4/3 of the proton's total angular momentum (also 1/2), and the d quark - 1/3. The distribution of polarized quarks can never exceed the distribution of unpolarized quarks (further discussion in Roberts, 1990).

The neutron (1 quark u^+ and 2 quarks d^- , see Cline, 1988) has also an electric-dipole moment, the upper limit of which has been recently measured (Smith *et al.*, 1990). The interest of neutron's electric-dipole moment is that "it would violate the combination of charge conjugation invariance and parity known as CP symmetry. As such, any electric-dipole moment would take the opposite sign for the antineutron, and thus discriminate between matter and antimatter" (Ellis, 1990).

Quantum theory holds that two photons emitted by a particular light source share their similarly oriented polarization. According to Clauser and Freedman's experiments recently recorded by Linden (1990), "a change in one photon did alter the polarization of the other" as if they were not separate objects and thereby obeying to the laws of quantum mechanics also applied to other "wave particles" such as leptons (electrons, etc).

In a search for understanding the charging of storm clouds, and contrarily to previous conclusions from Wilson and Simpson (see Williams, 1988) that electrical structures of thunderclouds were either a positive dipole (Wilson) or a negative dipole (Simpson), their actual structure is tripolar rather than dipolar. The correct explanation for this tripolar structure of thunderclouds is now known to lie in the microphysics of charge transfer between graupel particles (soft hail) and ice crystals (Williams, 1988).

3) Polarized conductivity

In a semiconductor the electrons move through an array of constituent atoms arranged in a crystalline lattice. Electrons move with great ease through gallium

arsenide circuits. This compound is made into bipolar transistor devices by depositing it in three layers: electrons n-type doping, holes p-type base and n-type collector. These compose light-emitting diode of gallium arsenide alloyed with aluminium. Gallium arsenide photodetectors respond faster than silicon ones. They can also detect light by reversing the reaction and the resulting photodetector converts the flash signal to electronic pulses. Such optoelectronic computing systems can be linked by optical fibers which greatly increase the efficiency of the digital computing circuitry (Brodsky, 1990).

D. MAGNETIC POLARIZATION

1) Cosmological level

The sun's magnetic field can affect many aspects of the sun's surface and atmosphere. It oscillates along a 11-year variation of sunspot number. Measurements of sunspot spectra (Zeeman effect's analysis) showed that the strength of the magnetic fields around sunspots is thousands of time stronger than the earth's field. Most spots occurred in paired groupings that resemble giant magnetic dipoles roughly parallel to the solar equator. According to Foukal (1990), the great astronomer Hale already announced in 1924 that this switch in polarity occurred at each activity minimum, in the midst of a 22-year solar magnetic cycle and was a basic feature of the sunspot cycle. The largest areas of single magnetic polarity are the sites of spot formation. These solar magnetic changes may have their effects on the earth's periodic climate changes.

2) Magnetic fields

The discovery of ferroelectric crystals such as barium titanate (BaTiO₃) offered an electrically switchable, two-state device with which one could encode the 1 and 0 states required for the Boolean algebra of binary computer memories. A tetragonal ferroelectric crystal has two polarization states in which the centrally located Ti^{4+} ions are involved through their displacement up or down with respect to the other ions (Ba²⁺ or Pb²⁺, O²⁻). In a crystal of PbTiO³, for example, there would then occur regions in which the polarization is up and regions it is down, called "ferroelectric domains" (Scott and Paz de Araujo, 1989). Most important for memory applications, the polarization of the entire crystal can be switched from up (+1) to down (0) by reversing the applied field. This ferroelectric memory progressively fades when the amount of switched charge decreases with use or by retention failure when the stored charge decreases to a level where the + or - state of polarization cannot be sensed.

All ferroelectric materials display a hysteretic behavior relating polarization and applied field, so that there is a nominal threshold (coercive field) above which the polarization changes sign.

4) Spin polarizations

Dipolar interaction between two nuclear spins depends on size and orientation of the magnetic moment as well as on the distance. In NMR spectroscopy which is based on the Zeeman phenomenon (Ernst et al., 1987), nuclei with a kinetic moment of spin I higher than 1/2 have a quadrupolar (Q) electric moment. The nuclear quadrupolar resonance (NQR) is bound to a nonspherical symmetry in the distribution of electric charges on the nuclear volume. This NQR can only be observed on a limited number of nuclei but is helpful in the study of the electric structures of chemical bonds (Lucken, 1969).

E. LIGHT POLARIZATION

A light ray can be polarized by reflection on a polarizer and the intensity of the reflected ray received on an analyzer varies with its incident angle. The proportion of polarized light in the light ray or the rotation of the polarization plane of light are measured with a polarimeter (Pariselle, 1936).

II. MOLECULAR DIPOLES AND CHIRALS

A. ELECTRIC DIPOLE MOMENTS

Dipolar electric moments and dielectric polarization have been surveyed by Errera (1928, 1935). Further study of the dielectric response of matter to an applied electric field has contributed to the measurement of molecular dipole moments (Price, 1969). The induced dipole moment per unit volume or polarization consists of two components: a *polarizability* one which arises from the distorsion of the electronic distribution of the substance, and an *orientation* component. Farley and McClelland (1990) have demonstrated that even in collisionless molecules, "hot isolated polyatomic molecules can reorient in response to an external field, thereby giving rise to this second component of polarization".

B. MINERAL DIPOLES

1) Dipolar water

Among recent and complementary knowledge about water biophysics and relevant to polarity, mention can be made of Saenger's 1987 review. It mainly concerns the relationships between hydration water and hydrogen bonds. Hydrogen bonding dynamics involves flip-flops and movement of water along the surface of macromolecules. Water would not have its particular properties if the molecules were not associated by hydrogen bonds $O-H \cdot \cdot \cdot O$. If the O-H group is involved in hydrogen bonding it becomes polarized (see II in I). In the association of water molecules to the surface of proteins or nucleic acids, hydrogen bonding of type (water) $O-H \cdot \cdot \cdot Y$ is the main attractive force. When the $O-H \cdot \cdot \cdot O$ bonds all run in the same direction, this is called *homodromic*; it is indicative of the influence of the cooperative effect. When a water molecule donates two hydrogen bonds this gives rise to *heterodromic* situation, where hydrogen bonds are randomly oriented.

According to the idealized structural model for water presented by Finney (1982), the simplest picture of the molecule "assigns partial charges to the two hydrogens and the two lone pairs which are considered to be disposed in an approximately tetrahedral manner. Each molecule is capable of forming four hydrogen bonds to neighbouring molecules" (see also I). Among the three proposed models of the water-water hydrogen bond, the PE model (water molecule electron distribution in terms of an electrical multipole expansion, see Barnes *et al.*, 1979) represents the water molecule electron distribution in an electrical multipole expansion. According

to Finney (1982), "the experimental dipole moment and quantum mechanical quadrupole are used, together with a dipole polarizability to try to handle the cooperative effects". Switching on polarizability in the PE model would therefore affect only the dipole-dipole and dipole-quadrupole energy terms (Finney, 1982).

The local dipolar field of protons of liquid water is averaged out by fast isotropic rotation and translational diffusion, and this gives a single narrow line in the NMR spectrum. In a molecular or biological system which can restrict water motion, causing an anisotropic averaged orientation, the NMR spectrum of the preferentially oriented water molecules can be given by a line pair or doublet. Lenk *et al.* (1980) have reported such NMR doublets spectra due to "structured" water in plant systems.

A typical example of efficient proton translocation across or along the surface membrane is the movement of protons across a cell membrane after their generation in some oxidation process. A high level of proton conductivity is extremely rare in crystalline solids. Thomas and Farrington (1982) have proposed that the proton conduction mechanism in one of the very best crystalline proton conductors so far studied ammonium/hydronium β "-alumina is a useful model mechanism for biological proton transfer. This proton conduction mechanism deduced from an accurate single crystal neutron diffraction study involves a classical Grotthus-type mechanism (see below).

In relationship with bilayer membranes (see IV.B.2) it should be pointed out that "an ion in water is stabilized by the favorable interactions of the water dipoles, the hydration energy. To remove an ion from water and place it in the middle of a membrane is unfavorable because of the loss of this hydration energy". The most successful model for quantifying this is the Born model described in Gennis (1989). In addition to this Born energy, a second component due to the polarization arises at the dielectric interface. An "image energy" results from the "presence of a charge on one side of the interface which causes the dipoles in the medium on the other side to reorient".

Cell water is modified by solvation which arises when water abuts a cell surface. Molecules become restricted in their motions and a greater proportion of them have four (rather than three or fewer) hydrogen bonds with their neighbours. Water modified in this manner is called vicinal (see I and Drost-Hansen and Singleton (1989).

Virtually all of the water in cells is considered to exist as polarized multilayers arising from fixed charges on extended protein surfaces. Cardinal sites exist on these particular proteins, the degree of binding for a given ion being influenced by a number of factors. Clegg (1982) further commented "ATP binding at the cardinal site leads to cooperative alterations and the selective accumulation of K⁺ over Na⁺, and generates the polarized multilayers of water; ATP splitting and the removal of ADP results in a movement of the system to a lower energy state in which the ion selectivity is lost as is the polarization of water".

Protons can be transferred along lipid/water interface in the absorbed water molecule network by a Grotthus-type mechanism (ref. in Tocanne and Teissié, 1990, see also IV.B.2.a).

C. ORGANIC DIPOLES

2) Multiple molecules (polar chains)

Charge transfer molecular interactions are of high significance in biology (Sklifkin, 1980). Electrons are delocalized in molecular conjugated systems (alternate single and double bonds). The polarization of these molecules is enhanced when they carry hydroxy-substituent(s) which behave as electron-attracting groups. Consequently, Pont and Pezet (1990) could suggest that "the polar interaction of these molecules with membraneous proteins could lead to a destruction of the cellular membranes by depolarization" (see IV.B.2.d). This could account for the biocidal effects of highly conjugated phenol derivatives such as the natural hydroxystilbenes which are efficient protectors of grape berries against the grey mold *Botrytis* (Pont and Pezet, 1990).

D. CHIRAL MOLECULES

Pasteur (1884, see I) audaciously extrapolated from molecular asymmetry the famous aphorism "la vie est apparue dans une brisure de symétrie". If we equate asymmetry and polarity, this would therefore mean that polarity is basic to the arisal of living matter.

From atoms to human beings, nature is asymmetric with respect to chirality (Gardner, 1979) and "clues are beginning to emerge that connect chirality on different levels". Thus, and as resulting from the weak nuclear Z force between electrons and nuclei, all atoms are also chiral. Consequently, the interaction that causes the helical motion does not conserve parity, and the mirror-image atom with a right-handed helical electron flow does not exist in nature (Hegstrom and Kondepudi, 1990, see I.B).

Chirality has its fundaments in the asymmetry between electron and positron; this asymmetry follows up in the hydrogen atom and reaches its full expression in the carbon asymmetry (see II.D). The basic molecules of life all have a specific handedness. They are therefore asymmetric (see I.B): its amino acids are left-handed, whereas its sugars are right-handed (see I). Chiral compounds which dissociate into enantiomers display a sharp difference in biologic activity. Chemists have been able

to induce a selection between two enantiomers and to develop methodologies for asymmetric syntheses initiated from prochiral center (Mosher, 1971, in Morrison, 1983-1985; see Oppolzer, 1987 and Holmstedt *et al.*, 1989). Chiral auxiliaries have been produced around asymmetric centers using organo-copper reagents. New bondings have thus been obtained with the concourse of highly stereo reactions in compounds such as diverse drugs (R(+)-S(-) thalidomide, etc.), pheromones, and perfumes (Oppolzer, 1987).

III. MACROMOLECULAR POLARITIES

A. FREE MACROMOLECULES

1. a) Deoxyribonucleic acid (DNA)

a¹ Structure. As noticed in I (Fig. 5) the two polynucleotide strands of the DNA double helix have opposite polarities and transcription only occurs from the sense strand (+) in the $5' \rightarrow 3'$ direction.

The bipolar pattern of the DNA double helix has important consequences on DNA recombination processes which involve restriction enzymes (Arber, 1974; Nathans and Smith, 1975). The recognition sequence for representatives of these site-specific endonucleases such as *Eco*RI and *Hind*III is a palindrome, i.e. a sequence of six inverted repeat base pairs showing a twofold rotational symmetry. The inverted polarity of the two DNA strands imposes a positioning of the cleavage sites outside the axis of palindromic symmetry. The ensuing asymmetric cutting produces single-stranded ends containing four bases of complementary sequences.

Seemingly, small variations in molecular structure or electrostatic potential at specific sites can make a critical difference in how the nucleic acid is organized and how it is recognized by other molecules in the intracellular environment. This is becoming increasingly clear from scanning tunnelling microscopy studies of calf thymus DNA and poly(rA)·poly(rU) which have shown that the helical pitch and periodic alternation of major and minor grooves can be visualized and reliably measured (Arscott *et al.*, 1989).

a⁴ Mutations. Those causing variegation are due to the action of transposons, a group of genetic elements known to move from one location in the genome to another. Certain strains of Saccharomyces cerevisiae contain an intron endowed with the ability for transposition in the gene coding for mitochondrial RNA which is absent from the corresponding gene of other strains; most of the progeny between intron plus and intron minus are positive (Dujon et al., 1974). This phenomenon, termed "polarity of recombination" by Bolotin et al. (1971) resembles a duplicative transposition which is characteristic for prokaryotic transposons.

As recently outlined by crystallographic structural studies of contacts in repressor-operator complexes, "positioning contacts" appear to be important conserved features within families of helix-turn-helix proteins (Pabo *et al.*, 1990).

b) Ribonucleic acid (RNA)

Antisense RNA molecules can selectively turn off genes and be used as antisense expression vectors to produce pigment variegations in flowers (Weintraub, 1990).

2. PROTEINS

As one of the recently described DNA-binding motifs, the zinc finger protein coordinates with a Zn²⁺ ion through paired cysteine and histidine residues along the amino-to-carboxyl protein dipole (Johnson and McKnight, 1989).

The thermodynamics of membrane-located proteins containing large (hundred of Debye units) permanent dipoles has been outlined by Schwarz (1978). Ordered water molecules can contribute directly to the properties of proteins by influencing their interaction with ligands. In their studies of atomic structures of the complexes of the L-arabinose-binding protein with sugars, Quiocho *et al.* (1989) have found that "two hydrogen-bonded water molecules in the site contribute further to tight binding of L-arabinose but create an unfavourable interaction with a methyl group of D-fucose".

5. ENZYMES

The distribution of charges within the charge-relay system (or "catalytic triad") at the active site of the serine proteinases has been further investigated. An Asp--Asn mutant in rat trypsin has been engineered by Craik *et al.* (1987). As reported by (Blow, 1990), this mutant showed that "the polarization of the histidine by the buried aspartate enhanced the reactivity of the serine". Warshel *et al.* (1989) have used the technic of computational chemistry "to estimate the effect of the charged carboxylate group and the polarized histidine on the reactivity of the serine side-chain surrounded by water".

B. AGGREGATES

2. c) Polar viral morphopoiesis

Packaging of bacteriophage λ DNA involves polarity of chromosome entry into the prohead (I, p.70) from the *Nu1* end to the *R* end (Becker and Murialdo, 1990).

IV. SUBCELLULAR POLARIZATIONS

B. SURFACE MEMBRANES

2. a) Biochemical properties

The bilayer membrane can be modelled electrically as a thin slab of non conducting material separating two aqueous solutions and thereby acts as a simple parallel-plate capacitor (Gennis, 1989). "Its dielectric constant is a measure of the polarizability of the material and the degree to which any permanent electric dipoles which may be present in the material respond to an electric field (voltage difference)".

The amphiphilic phospholipids form spontaneously well-organized bilayer structures in water which are the basic architecture of biomembrane. Evidence has been obtained with membrane model systems, which support the view that lateral proton conduction occurs at water/lipid interfaces (Tocanne and Teissié, 1990). The polarity at these interfaces in terms of dielectric constant is different of that of bulk water. This means that, "in terms of micropolarity or water molecular dipole moment, the lipid/water interface region is more than likely anisotropic both in terms of structural organization and electrical properties" (see B.2b).

As for the very low permeability of the lipid bilayer to cations as compared to anions (see Tocanne and Tessié, 1990) it is ascribed to the positive polarization potential of the surface membrane (see B.2.d) which would constitute an energy barrier against the transport of positively charged compounds across membranes.

Permeability coefficients have been determined for several kinds of small molecules. Among them, water can relatively easily penetrate the membrane bilayer. As commented by Gennis (1989) "It may seem surprising at first to learn that water can so readily penetrate the phospholipid bilayer". However, "there is no substantial water to be found inside the membrane beneath the carbonyl groups".

2. c) Energy transduction

In 1961, two proposals were made as to the way in which electron-transfer reactions of the cytochrome chain — the chain used in the oxidation of NADH by molecular dioxygen — could be connected to ATP formation without the intervention of chemical intermediates (Williams, 1989). Both mechanisms invoked the transduction of the energy of the oxidation/reduction reaction to a proton gradient before the gradient generates ATP. The two mechanisms, sometimes termed the delocalized (Mitchell) hypothesis and the localized (Williams) hypothesis, are very different: in the first, protons generated by oxidation appear only in aqueous phases;

even ATP is generated by an electric field acting on the ATP synthetase and not by proton flow; in the second, protons move in proteins within matrices and aqueous phase equilibrations are ignored in the development of proton gradients, in proton diffusion and in the ATP-synthesis step. To distinguish between these mechanistic possibilities, long series of experiments (Wikström, 1989) have been carried out on separate parts of the cytochrome chain, especially on the last stages of the electron-transfer reactions, those of cytochrome oxidase.

Electric currents produced by oxido-reduction reactions, also called Faraday currents, can be assayed by electrochemical methods such as those of polarography. The polarograph apparatus works with three electrodes (see Monnier *et al.*, 1979): an indicator capillary electrode on which oxido-reduction reactions occur at the surface of mercury drops, a reference electrode allowing to impose to the first one a constant potential while varying the voltage, and an auxiliary electrode insuring passage of current. Registered curves of intensity-potential of chemicals such as metal ions allow their quantitative assay. Dissolved O₂ can also be measured by the polarographic technique (Fork, 1972).

In artificial fuel cells, gases are combined electrochemically such that the exothermicity is converted directly to electrical energy and the only reaction product is water. Dyer (1990) observed gas — electrical energy conversion processes occurring whithin very thin films of gas-permeable, ionically conducting membranes of hydrated aluminium oxide, as a prototypical membrane. Both polarity and the magnitude of the voltage were unexpected. The covered inner platinum electrode was positive and the polarity of the cell could be changed in $H_2 + O_2$ mixtures only when the outer platinum catalyst was changed to a nickel catalyst. This shows the strong dependence of cell polarity on the metals used and their sequence, suggesting that "different electrochemical kinetics might establish the polarity observed" (Dyer, 1990).

2. d) Electric potentials

Many possible factors can contribute to the amount of electrical work to move a charge through a membrane (Gennis, 1989): a) associated work with dielectric constant; b) internal dipole potential by orientation of the dipoles at the membrane surface resulting in a positive potential in the center of the phosphatidylcholine bilayer; c) surface potential which, in most biomembranes, is negatively charged, usually due to the presence of acidic, anionic phospholipids; the electric potential at the shear plane which is the plane defining what migrates in the electric field is called the zeta potential (McLaughlin, 1977); it someway controls the electrophoretic mobility of charged vesicles (electrokinetic effects); d) transmembrane potential which is defined as the difference in the electric potentials of the two bulk aqueous phases separated by the membrane. The asymmetric charge distribution generates

transmembrane potentials which are usually negative inside and can be measured with fluorescence polarity methods using probes such as merocyanine or anilinonaphthalene.

The membrane surface potential (ΔV) is the sum of an electrical term (Ψ o) and a dipolar or "polarization" term (ΔV p) which exhibits high positive values (about 300 to 500 mV). The variously oriented and rotating strong dipoles of lipid polar heads would contribute to the surface polarization potential and this view (Tocanne and Teissié, 1990) has been correlated with the concept of "molecular electrometer" as developed by Seelig *et al.* (1987) on the ground of 2 H-NMR experiments using parameters such as the deuterium quadrupole splitting.

In fungi, marked changes in the membrane potential detected by [³H]tetraphenylphosphonium (TPP⁺) uptake rate have been caused by illumination of dark-grown mycelium of *Trichoderma viride*. An initial hyperpolarization of the plasma membrane was found to be accompanied by a rise in the intracellular ATP concentration and by changes in the intracellular level of cyclic AMP (Gresik *et al.*, 1988).

In higher plants, blue light is known to activate the electrogenic proton pump to hyperpolarize the plasmalemma (Assmann et al., 1985 and Shimazaki et al., 1986). Plasma membrane hyperpolarization caused by auxin (IAA), accompanied by short time oscillations in the electric potential of corn coleoptile cells, is paralleled by cytosolic pH drops as well as changes in Ca²⁺ activity (Felle, 1989). Moreover, the activity of the plant plasma membrane enzyme NADH oxidase which transfers the electrons from NADH to oxygen in the absence of added electron acceptors has been linked to membrane polarization (Novak and Ivankina, 1983). In Conjugatophycean green algae photoreception, a tetrapolar gradient of phytochrome created by light perception is achieved by the dichroitic orientation of plasma membrane-bound phytochrome molecules; blue-light also appears to mediate a tetrapolar gradient of the sensor pigment proper mediating tetrapolar actin anchorage sites on the plasmalemma (Grolig and Wagner, 1988).

Gating and ion selectivity of calcium channels have been further studied by electrophysiological experiments. Subtypes of calcium channels have been classified according to their voltage threshold for activation and by their inactivation characteristics (Wray et al., 1989). Current dependence of channel gating has been tentatively ascribed to the formation of dipoles along the trajectories of ion movement that exist during dipole relaxation time (Kostyuk et al., 1989). This new approach would assume that "ion transition through the open channel produces local displacements of charged molecular groups lining the wall of its steric region". During the process, the frequency of ion transitions would increase drastically and become comparable with frequency of dipole relaxation (Kostyuk et al., 1989).

Release of Ca²⁺ from the sarcoplasmic reticulum (SR) following depolarization of transverse tubules (T-tubules) triggers contraction of the skeletal muscle. The foot

structure of the SR is part of a molecular bridge which spans a short gap between the T-tubules and the terminal cisternae of the SR. Large cytoplasmic extensions of the molecule evidently attach to the dihydropyridine receptor complex in the T-tubules (Agnew, 1989). There is also evidence that the dihydropyridine receptor in the T-tubule membrane of skeletal muscle functions not only as slow calcium channel but also as an essential component of coupling, probably as the voltage sensor (Takeshima *et al.*, 1989). A model of the structure of the dihydropyridine-sensitive calcium channel has been proposed (Catterall *et al.*, 1989) in analogy with current models of the structure of voltage sensitive sodium channels.

Chloride (Cl⁻) channels (normal and pathological) were activated by patch excision which caused large membrane depolarization. This allowed Welsch *et al.* (1989) "to use depolarization as a "tool" to determine if a Cl⁻ channel was present in a patch". Active chloride transport can be light-driven by retinal proteins. These bacterio- or halorhodopsins function as inward-directed electrogenic pumps for Cl⁻ ions (Zimányi and Lanyi, 1989). Parallely, these pumps transport protons out of the cell interior, thereby generating an inside-negative membrane potential.

Opening and closing of chloride channels studied in the electric ray *Torpedo californica* are unequally timed. This asymmetric electric conduction increases with transmembrane electrochemical gradient for the chloride ion thus demonstrating that the channel-gating process is not at thermodynamic equilibrium (Richard and Miller, 1990).

2. e) Action potentials

They are not only generated in animals (see I) but also in fungi, algae and higher plants in response to light, heat, cold, chemicals, electrical stimulus, and wounding as reviewed by Pickard (1973) and Simons (1981). Davies (1987) considered action potentials as multifunctional signals in plants and proposed a unifying hypothesis to explain apparently disparate wound responses. Action potentials could also be a unifying factor to explain the involvement of an interaction between Ca²⁺ flux and auxin transport in the role of gravity in geotropisms (De la Fuente, 1984, also VIII.A.2.c⁴).

Cell electrophysiology and membrane transport in plants have been recently reviewed by Bentrup (1989) who stated that "the evergreen question of the role of Ca²⁺ during the characean action potential will remain elusive as long as the characean plasmalemma is not routinely accessible to patch clamp technics". In the *Characeae*, depolarization occurs by diffusive Cl⁻-efflux and repolarization by diffusive K⁺-efflux (Köhler *et al.*, 1986; Gradmann, 1989).

The role of K^+ in the mechanisms of action potentials has been further analyzed in the green alga *Eremosphaera viridis* by Köhler *et al.* (1985, 1986) who showed that it is caused by a transient opening of a K^+ channel which is not gated by the membrane potential.

In animals, action potentials experimentally evoked by electrical activity can suppress neurite elongation and growth cone motility (Cohan and Katter, 1986) and thereby may influence structure and connectivity within the nervous system (see also VI.A.2.i).

Following electrical activity in excitable cells, there is an increase in intracellular Ca²⁺ concentration. Silver *et al.* (1990) also report that clustering of L-type Ca²⁺ channels causes intracellular Ca²⁺ hotspots at the neural growth cone. Enzymes with a micromolar requirement for Ca²⁺ at the hotspots are therefore activated by the ensuing depolarization. The role of voltage-dependent calcium influx in controlling nerve cell outgrowth remains puzzling because "also raised intracellular Ca²⁺ concentration triggers outgrowth of the growth cone margin, neurite elongation requires low intracellular Ca²⁺ concentration". According to Silver *et al.* (1990), the fact that "electrical activity can selectively raise intracellular Ca²⁺ concentration in the growth cone, leaving neurite calcium concentration low would resolve this paradox".

C. ENDOMEMBRANAR AND VESICULAR SYSTEMS

1. Endoplasmic reticulum

In the endomembranar sorting process, proteins destined for transfer are sequestered within membrane vesicles that bud off from a donor organelle and then fuse with the appropriate acceptor organelle. Vesicle fusion in several distinct branches of this complex distribution network as well as transfer of vesicles between the rough endoplasmic reticulum (ER) and the Golgi complex require the same cytosolic protein, a tetrameric, N-ethylmaleimide-sensitive protein (NEM) called NSF (Beckers et al., 1989). Such transfer requires ATP and is inhibited by NEM or the monoclonal antibody against NSF. NSF is required in a late, calcium-dependent transfer step; this step is most likely the fusion step. Surprisingly, the deduced protein of cloned and sequenced NSF product showed sequence similarity with the product of a yeast gene (SEC18) previously shown by Schekman and Novick (1982) to control the transfer of vesicles between the rough endoplasmic reticulum and the Golgi complex; more recent studies suggested it has a function in endocytosis (Riezman, 1985). These results raise the possibility that "fusions between different organelles derived from the rough endoplasmic reticulum may all be catalyzed by the same set of proteins" (Schatz, 1989).

2. Golgi apparatus

This compact structure colocalizes with the microtubule organizing center (MTOC) in a perinuclear region of fibroblasts. Intact interphase microtubules but

not microfilaments appear to be required for this specific location of the Golgi apparatus. This has been demonstrated by the scattering of Golgi elements after treatment with the microtubule depolymerizing drug nocodazole, and by the subsequent reclustering of the Golgi elements when nocodazole is removed (Ho et al., 1989). A protein may be involved in linking the Golgi apparatus to the microtubule network and the MTOC in vivo (Allan and Kreis, 1986). A fungal antibiotic, brefeldin A, produces a reversal of traffic polarity i.e. a rearrangement of Golgi elements into the ER, thereby inducing a secretion block (Bosshart et al., 1990). Such "violation of the one-way system" has been further discussed by Armstrong and Warren (1990).

D. ORGANELLES

3. Chloroplasts and phototransducing membranes

Most of the chloroplast proteins are imported from the cytosol and polarly directed into six different compartments (Smeekens et al., 1990). Two sorting systems are involved in this import and intraorganellar transport of nuclear-encoded protoplast proteins. Additional sorting informations located at N- termini are contained in thylakoid lumen proteins. The information present in transit peptides, decoded by the chloroplast import machinery, is not yet known.

The electron transfer reactions in photosystem II take place within the so-called reaction center grouping numerous antenna pigment molecules (chlorophyll, etc.) as well as organic ions and charged atoms (manganese, calcium, etc.). The stepwise transfer of electrons through this reaction center succeeds in pulling far apart the mutually attractive positive and negative charges. The task of the photosystem II is thus to act as a tiny capacitor, storing energy by separating and stabilizing positive and negative charges on either side of the thylakoid membrane (Rutherford, 1989). The water-splitting reaction produces four protons and four electrons released simultaneously with O_2 in that water-oxidizing clock which is a cyclic mechanism of four states (Gowindjee and Coleman, 1990).

E. CYTOSKELETAL COMPONENTS

That the cytoskeleton is someway involved in plants intracellular movements, perception mechanism and transmission effects has again been emphasized by Hensel (1989b) who concluded that "the function of the cytoskeleton is to generate and maintain cell polarity".

As for fungal cells, they have been comprehensively surveyed in 1987 and 1989 by Hohl.

1-2. Microfilaments (actin-myosin)

Both actin and myosin filaments have definite polarities and well-ordered structures (see I). Actin filaments can move in opposite directions on tracks of myosin heads. They always move foreward but never backward reversing the polarity of the movement. According to Toyoshima *et al.* (1989) "The direction of movement is therefore determined by the polarity of the actin filament".

Myosin heads can form reverse chevrons and, when tethered in a single thick filament of a mutated *Drosophila* flight-muscle sarcomere, can bind with opposite rigor crossbridge angles to flanking thin filaments, which are apparently of opposite polarities (Reedy *et al.*, 1989).

The driving force for the rearrangements of the actin cytoskeleton in cell motility, division and differentiation is provided by actin-binding proteins. The addition of actin subunits to the barbed end of actin filaments and the nucleation of polymerizing actin *in vitro* are controlled by capping protein. Recent experiments suggest that capping protein regulates polar distribution *in vivo* of actin filaments. The actin cytoskeleton is disrupted in yeast capping protein mutants, indicating that "the asymmetric distribution of actin in budding yeast (see VI.A.1.a² in I) depends on the proper functioning of several actin-binding proteins with apparently different functions" (Amatruda *et al.*, 1990).

The uniform angle and conformation of myosin subfragment 1 (S1) bound to actin filaments (F-actin) "attest to the precise alignment and stereospecificity of the binding of these two contractile proteins. Because actin filaments are polar, myosin heads must swing or rotate about the head-tail junction in order to bind" (Reedy et al., 1989). Adams and Pollard (1989) have shown for the first time that the single-headed myosins called myosin-I can bind directly to NaOH-extracted membranes isolated from Acanthamoeba and to vesicles of pure lipids with an affinity sufficient for extensive binding in the cell. Membrane-bound myosin-I may provide a mechanism for many cellular movements previously thought to involve filamentous myosin-II (see V, in I) and for the specification of sites of cell surface growth (Drubin et al., 1990).

For a general review about cytoskeleton microfilaments, see Kristen (1987).

1-3. Microfilaments-microtubules (actin-tubulin)

In the cortex of the giant coenocytic green alga *Caulerpa*, amyloplasts are transported along microtubular strands as shown by the fact that both microtubule-and dynein-specific inhibitors block movements of these organelles. In contrast, chloroplast movement is blocked by cytochalasin but not by colchicine thereby showing that immobilization and movement of chloroplasts are dependent on intact microfilaments of actin but not on microtubules (Menzel and Elsner-Menzel, 1989).

F. NUCLEI AND MITOTIC FIGURES

2. Polewards chromosome movement

The bipolar attachment of chromosomes to the spindle occurs well before all the chromosomes congregate metaphasically. In the normal functioning of the mitotic spindle most of its growth and disassembly take place at the end of the microtubule away from the pole. All microtubules have the same polarity and the fibers behave differently depending on the structure in the spindle to which they bind. Most important as microtubule-organizing center is the centrosome which serves as a seed to start microtubule polymerization; thereby it defines their polarity. That polarity, or asymmetry, is crucial to the functioning of microtubules (see I) by at least two of its functional consequences: at the ends it causes the (+) end to add and lose subunits faster than the (-) end; along the surface it influences the orientation with which proteins will bind to the microtubule surface (McIntosh and McDonald, 1989).

The molecules involved in the mechanical forces moving polewards chromosomes begin to be unraveled (Vale and Goldstein, 1990). Among such mitotic motors there are kinesin motors and perhaps the newly discovered dynamin motor (Shpetner and Vallee, 1989) which forms cross-bridges and induces ATP-dependent sliding between antiparallel microtubules *in vitro* (McIntosh and Koonce, 1989). Kinesin is a microtubule-interactive, force-generating ATPase acting as a plus-end motor in intracellular transport of vesicles along microtubules (Vale, 1987, and others, see in \mathbf{I}). The inherent asymmetry of the polymer (actin or tubulin) and the motor is necessary for the unidirectional movement of the motor along the polymer. It is toward the barbed (or +) end of the actin filament that myosin motors such as myosin I (single ellipsoidal head) move.

A superfamily of kinesin motors acting in fungal nuclear fusion and division has now been described in *Saccharomyces cerevisiae* (Meluh and Rose, 1990) and in *Aspergillus nidulans* (Enos and Morris, 1990). Such kinesin motors bear either round or rectangle heads at the end of the α -helical coiled coils. Short single-headed kinesins analogous to myosin I, kinetochore-specific kinesins, and perhaps kinesins may also be expected to be involved in morphogen or RNA transport as force-producing proteins (Vale and Goldstein, 1990).

V. POLAR CELL MOVEMENTS

B.1. Cilia-flagella

In the green unicellular alga *Chlamydomonas*, a component of contractile flagella roots is the centrosome-associated phosphoprotein centrin. This type of structural organization contributes to define its cell polarity through cell axiation (Fig. 1, in Salisbury, 1989).

2. Gliding movements

Bacterial gliding motility appears to be dependent on the establishment of transmembrane potential and any depolarization (*not* depolymerization as wrongly written in I p. 132) by protonophores such as 2,4-DNP or CCCP results in a cessation of motility.

3. Amoeboid motion (transient polarity)

Both the single headed myosin I and the double headed myosin II are mechanochemical enzymes which generate force through the hydrolysis of ATP when complexed with F-actin.

Fukui et al. (1989) show by immunofluorescence microscopy that nonfilamentous myosin-I occurs at the leading edges of the lamellipodial projections of migrating Dictyostelium amoebae, which are devoid of myosin II, whereas filamentous myosin II is concentrated in the posterior zone of the cells. The authors suggested on the basis of these locations of the two forms of myosin and their known biochemical and biophysical properties that "actomyosin I may contribute to the forces that cause extension at the leading edge of a motile cell, while the contraction of actomyosin II at the rear squeezes the cell mass forward. Myosin I isoenzymes might have similar roles in metazoan cells, for example at the leading edges of neuronal growth cones, and in the extension of lamellipodia and pseudopodia of leukocytes, macrophages and fibroblasts." These observations suggest that "actomyosin I-dependent force-generating activity occurs at the leading edge (as in pseudopodia extension) and that actomyosin II-dependent force-generating activity occurs at the trailing end of a migrating *Dictyostelium* amoeba (causing the cell mass to move forwards)". This could explain "how myosin II-minus mutants can form smaller-than-normal pseudopodia at a relatively normal rate. Membrane-bound Acanthamoeba myosin I can generate force against actin cables however, and both Acanthamoeba and Dictyostelium myosin I will crosslink actin filaments and generate force between crosslinked filaments".

None of Fukui et al. (1989) observations is compatible with the participation either of other processes in amoeboid movement, such as membrane flow or the remodelling of the actin matrix, or of myosin I and myosin II in other motile activities. The significance of Fukui's team results is that they show the presence in the leading edge of a migrating cell of myosin I, which in conjunction with F-actin is known to be capable of producing force and movement.

To explain the rearward movements of membrane proteins in locomoting polymorphonuclear leukocytes, the experimentally best supported model implies the cytoskeleton (see I, pp. 133-137). The retrograde lipid flow hypothesis has been proposed by Bretscher (1984) as an alternative explanation for the rearward movements of membrane proteins. However, recently used techniques of low-light-level fluorescence microscopy and digital image-processing of photobleached images disprove that lipid flow model (Lee *et al.*, 1990). By further implicating cytoskeleton in proteins movements, they also validate the conclusion of Sheetz *et al.* (1989) that such a membrane flow in the leading edge of amoeboid cells does not drive rearward movements of membrane glycoproteins.

About the motor of amoeboid motion, there is much evidence linking actin-based system to the generation of motile structures in the cell (Bray and Vasiliev, 1989). Nevertheless, a mutant of *Dictyostelium discoideum* deficient in α-actinin and in which movements are unimpaired has been obtained by Gerisch's group (Wallraff *et al.*, 1986; Schleicher *et al.*, 1988). "Motile life without myosin" also exists as shown by mutants of *D. discoideum* that lack normal myosin-II (Knecht and Loomis, 1987; De Lozanne and Spudich, 1987, see I, p. 134). Since, André *et al.* (1989) have described a strain of this slime mold lacking severin (actin-filament fragmenting protein) even though still able to move. A relative interpretation of these findings is that "there is an extensive overlapping redundancy in the activity of actin-binding proteins *in vitro* and more than one way to crosslink, fragment or even to move actin filaments" (Bray and Vasiliev, 1989). There is analogy between the behavior of such parallely distributed processor of the locomotive cytoskeleton of *Dictyostelium* amoebae and of the cytoskeletal network intervening at yeast budding (see VI.A.1.a²).

VI. POLAR CELL GROWTH

A.1. MONOPOLAR OUTGROWTH (EMERGENCE)

In our present state of knowledge, cytoplasmic *microtubules* are dispensable for bud outgrowth (see I) but required for specific, single or double budding of yeast cells or fungal spores to direct their mono- or dipolar axiation toward the site(s) of bud formation. By contrast, polarly localized actin *microfilaments* appear to be an absolute requirement for the budding processes.

a² Yeast budding

The cortical actin cytoskeleton seems to specify sites of growth of the yeast cell surface (Adams and Pringle, 1984, see I; Novick and Botstein, 1985). An actin-binding protein (ABP1p) might be involved in the spatial organization of cell surface growth and the identification of C-terminal protein domains suggests that such domains might serve to bring together signal transduction proteins and their targets or regulators, or both, in the membrane cytoskeleton (Drubin et al., 1990).

The cytoskeletal network in the budding yeast cell (Saccharomyces cerevisiae) behaves as a parallely distributed processor, as suggested by the finding of a protein (SPA2) associated with actively growing regions of the cell surface (Snyder, 1989). Such polarization of the growth process is disturbed in mutant cells displaying an inability to stop growing under nutrient-limiting conditions which often results in multiple budding (multipolar growth, see I, p. 187).

b¹ Fungal spores

In the germinating spores of *Mucor rouxii* the change in growth pattern from spherical to polarized correlates with the degree of DNA methylation and this, in turn, may be controlled by polyamine levels. The establishment of the polarized phase of growth in *M. rouxii* probably occurs through the regulation of the genes involved in the synthesis of products necessary for apical growth of the hyphae (Cano *et al.*, 1988).

c) Dimorphism

Quite recently, Crombie et al. (1990) have shown that the sites of budding and germ tube formation on yeast cells of Candida albicans were polarized preferentially towards the cathode. Buds were found to be less polarized than germ tubes at any given applied voltage. Moreover, polarization of germ tubes was biphasic.

2. TIP GROWTH

b) Fungal hyphae

In the models of hyphal tip growth, electric current does not always enter the growing end (*Allomyces* hypha drives an outward protonic current, see Youatt *et al.*, 1988 in **I**). As recently commented by Gow (1989) "Most of the evidence suggesting that ionic currents are involved in establishing and maintaining polar growth is essentially correlative, and it is not yet clear whether the current is a cause or consequence of polarity". However, Gow leaves open the possibility that "Cytoplasmic proton and calcium-ion gradients and fixed-charged gradients resulting from asymmetric transport of calcium into a cell may be involved in localizing growth". The same conclusions have recently been reached about differentiation at egg germinations of brown and red algae (Quatrano and Kropf, 1989; Waaland, 1989; see VII.C.3.a).

In hyphal tips of the oomycete Saprolegnia ferax, Heath and Kaminskyj (1989) observed that "all the organelles and the microtubules are non uniformly distributed, each showing a characteristic longitudinal gradient starting at a different point behind the tip". A few microtubules can reach the extreme tip but they were more abundant sub-apically. The authors concluded that "the correlated patterns of organelle and cytoskeleton organization from this and previous work show that neither the microtubules nor the detected arrays of actin are sufficient to account for most organelle arrangements".

The role of microtubules at the onset and maintenance of polarized growth of hyphae is still unclear. Intact microtubular tracks are required to initiate dominant, monopolar outgrowth from macroconidia of *Neurospora crassa* (Caesar *et al.*, 1988, see in I). However, further elongation of hyphae deprived of microtubules can still occur contortionally, with a damped polarity (Howard and Aist, 1980, see I).

Germlings of the bean rust fungus *Uromyces appendiculatus* treated with the microtubule-binding drug griseofulvin continued polarized apical growth even though showing changes in the morphology of their apical and subapical regions (Hoch *et al.*, 1987).

i) Animal neurites

A major question in developmental neurobiology is how developing nerve cells accurately extend processes to establish connections with their target cells (see Lasek and Black, 1988). This unsolved problem of polarized growth involves "both the nature of cues for growth cone guidance and also the question of how growth cones survey their environment for cues and respond by altering their direction of migration" (Bentley and Toroian-Raymond, 1986, see I). According to Lamoureux *et al.* (1989) "there is also controversy over whether axonal elongation is the result of a pulling growth cone and the role of tension in axonal elongation".

Earlier in this decade, the consensus was that axons or neurites elongated from tension generated by forward motility of the growth cone (Landis, 1983; Letourneau, 1982). It was presumed that contractile filopodia were the source of the tension moving the growth cone (Bray, 1982; Trinkaus, 1985). But this view was challenged by experiments showing that neurites elongate, albeit abnormally, in the presence of cytochalasin, which inhibits growth cone and filopodial movements (Marsh and Letourneau, 1984).

Bentley and Toroian-Raymond (1986) also reported an examination of the migration of pioneer growth cones deprived of filopodia by culture in agents which disrupt actin microfilaments. Under these conditions, axons continue to extend but a large percentage of growth cones are highly disoriented. Their results indicate that filopodia are not necessary for axonal elongation *in vivo* but that they are important for correctly oriented growth cone steering.

Additionally, high resolution, video-enhanced observations of growth cone activity argue against filopodial shortening as a source of tension, suggesting instead that an extrusion of cytoplasm rather than a pulling process, is the key event in neurite elongation (Goldberg and Burmeister, 1986; Bray, 1986; Aletta and Greene, 1988, ref. in Lamoureux *et al.*, 1989). Studies of slow axonal transport (Lasek, 1986) indicate that much slower cytoskeletal pushing underlies axonal elongation and direct measurements of neurite force as a function of growth cone advance show that they are linearly related and accompanied by apparent neurite growth (Lamoureux *et al.* (1989). No increase in force occurs in neurites whose growth cone fails to advance.

According to Mitchison and Kirschner (1988) there are three phases of axonal development: an actin based-system in which the leading edge becomes orientated, a consolidation phase in which filopodial microtubules become stabilized in their direction of future growth and a conversion phase to stable microtubules bundled within the axonal tube. The protein factor tau stimulates the conversion phase. However, tau expression is insufficient to induce polarity but tau antisense oligonucleotides can inhibit neurite polarity (Kosik and Finch, 1987).

Pulse-labelling studies performed both in mature nerve and in cell culture provided most of our knowledge of the axonal transport of cytoskeletal proteins. In 1975, Ochs has put forward his unitary hypothesis of axonal transport according to which proteins achieve different transport rates by having different affinities for a single moving vector. Tubulin and actin molecules are the essential components of the axonal cytoskeleton and considered by some (Black and Lasek, 1980) as a static complex travelling down the axon, a view challenged by others (ref. in Okabe and Hirokawa, 1990) who observed a gradual recovery of photobleached zones rather than their movement or spreading along the axon, both in neurons injected with fluorescein-labelled tubulin and actin. Therefore, these cytoskeletal components can be considered as "dynamic structures that continue to assemble along the length of the axon" (Okabe and Hirokawa, 1990).

In most recent and interesting experiments, Schnell and Schwab (1990) have shown that axonal regeneration and elongation in the rat spinal cord can be produced by the neutralization by monoclonal antibodies of myelin-associated neurite growth inhibitors.

VII. POLARIZED CELL DIFFERENTIATION

B. APICAL DIFFERENTIATIONS

- 1. Monopolar patterns
- a) Fungal exosporulation: a² Sporangia

A unique capability of excised segments of sporangiophores of the terrestrial mold *Phycomyces* is to regenerate new sporangiophores with sporangia (Götze, 1918). The excised segments in the sporangiophore preferentially regenerate at the apical end. In addition to this segmental polarity, there is a polarity of the whole sporangiophore. Moreover, the fact that "polarity is not destroyed by acropetal or basipetal centrifugation seems to indicate that the plasma membrane or the cell wall (see also proposal for algal axiation in C.3.a) plays a crucial role in the polarity". Galland and Ootaki (1987) conclude from their comprehensive review that the molecular basis for this polarity is still obscure, and one of the challenging problems in *Phycomyces* differentiation remains to discover what molecules constitute the actual gradient and where are they located?

The tip of the growing zone of the sporangiophores of *Phycomyces* (Bergman *et al.*, 1969) is the site where the gravitropic bending occurs (Sachs, 1879, in Shropshire and Lafay, 1987, see VIII.A.2.c⁴).

a³ Basidiospores

Basidia of *Coprinus cinereus* continue differentiation when explanted to water agar and vegetative hyphal tips monopolarly elongate from the four apical sites of the basidium expected to produce sterigmata (Chiu and Moore, 1990).

C. APICO-BASAL DIFFERENTIATIONS

3. a) Algal eggs (rhizoid-thallic poles)

In model systems of early embryogenesis of the Fucales, the site of inward current precedes and accurately predicts the site of rhizoid outgrowth (see I) and the polar axis can be oriented by external vectors (light, etc.) and two unequal cells result from the first division. Experiments with inhibitors (i.e. the cytochalasins) clearly implicate microfilaments in the process of axis fixation. Moreover, such polarization of two-celled embryo cannot occur in absence of a cell wall, demonstrating that the

presence of this cellular component is an absolute requirement for axis fixation. From these results, Quatrano and Kropf (1989) derive their actual working hypothesis that "axis fixation involves transmembrane bridges at the presumptive rhizoid pole, from the cell wall to the microfilament cytoskeleton".

Using repair shoot cells and rhizoids of the red alga *Griffithsia*, Waaland (1989) tested Jaffe's hypothesis (1968, 1979, see I) that transcellular currents are responsible for establishing and maintaining sites of localized secretion and growth. However, in repair shoot cells, the inflowing current continued even when the cell repair hormone rhodomorphin was withdrawn and elongation stopped. Thus, in *Griffithsia* "transcellular currents *per se* do not appear to control localized organelle accumulation and localized growth".

6. Higher animal cells

b) Epithelia (apical-basolateral poles)

The apical and basolateral, macroscopic domains of polarized epithelial cells are mostly large, morphologically distinct regions of the cell surface which are separated by proteinous barriers.

The rapid diffusion and equilibration of lipophilic NH₃ across cell membranes and the accumulation of NH₄⁺ seem to be governed by pH differences between compartments. Kikeri *et al.* (1989) reported that renal tubule cells from the medullary thick ascending limb of Henle have an apical membrane which is not only virtually impermeable to NH₃, but is also highly permeable to NH₄⁺. They proposed a model which would explain how this renal epithelium can mediate vectorial movement of NH₄⁺ between compartments of equal pH.

A hierarchy of sorting information with multiple sorting signals — apical and basolateral — present in different domains of a given plasma membrane protein has been suggested from the evidence that covalently attached glycosyl-phosphatidylinositol (GPI) acts as a "dominant" apical targeting signal. Polarized epithelial protein sorting might therefore rely on glycolipids (Lisanti and Rodriguez-Boulan, 1990).

VIII. MORPHOGENETIC POLARIZATIONS

A. PLANTS

2. Organismic polarities

a) Mushrooms

These higher fungi grow upwards and should be responsive to the gravitational field. The problem will be to find the gravity sensor and the way its signals are interpreted (also for the model mold *Phycomyces*, see below).

c⁴) Polar auxin transport and tropic curvatures

Bioelectric gradients along axial organs demonstrate morphological and physiological polarity in higher plants (Fensom, 1959; Scott, 1967; Zatsepina and Tsaplev, 1980; Goldsworthy, 1986). This electric polarity probably controls the distribution of phytohormones (Clark, 1937). Changing the bioelectric gradients by an external electric field has various consequences on plant growth and development (Lund *et al.*, 1947; Cholodny, 1956; Jaffe and Nuccitelli, 1977; Ellis and Turner, 1978; Medvedev and Markova, 1990).

In studies of gravity-dependent plant responses provided by the special conditions of spaceflights, interfering accelerations are relatively small (below 10^{-3} g) and termed "microgravity" (see Hensel, 1989a).

Plant morphogenesis in general does not appear to be considerably disturbed by microgravity, as shown by the polar differentiation of anise callus cultures into somatic embryos (Theimer et al., 1986). Compared to ground controls the distribution of the amyloplasts is shifted towards the proximal pole in statocytes of space grown roots (ref. in Hensel, 1986). This polarity of statocytes does not require the continuous action of gravity but develops also at microgravity. In statocytes of lentil roots differentiated in microgravity, the nucleus was preferentially located toward the gravity center of the cell (Perbal and Driss-Ecole, 1989). Polar differentiation of statocytes was also disturbed but only at the level of endoplasmic reticulum (ER) in seedlings of Zea mays launched from earth after germination, while those germinated at microgravity had aggregated ER in root statocytes (Moore et al., 1987).

By comparison, the normally negatively gravitropic sporangiophores of the terrestrial mold *Phycomyces* (see VII.B.1a²) become disoriented when cultivated aboard an orbiting spacecraft (Parfyonov *et al.*, 1979). The nature of the gravity receptor is still unknown (Shropshire and Lafay, 1987).

As previously suggested, statocytes polarity depends on a genetically prepatterned program (Sievers *et al.*, 1976). Since, agravitropic mutants of roots have been discovered (see Scott, 1990). Such mutants exhibit morphological and physiological abnormalities which suggest that they are unable to respond to the plant growth hormone auxin, indole-3-acetic acid (Hicks *et al.*, 1989). The root cap plays a role in root geotropism (Pilet, 1978) and its removal can also lead to an agravitropism (Moore *et al.*, 1990). Gravity could thus induce a change in cellular structure which somehow generates a chemical and/or electrical signal in the cap.

The starch statolith hypothesis attempts to explain gravity perception in plants. Starchless (phosphoglucomutase deficient) mutants recently produced in Arabidopsis thaliania (Caspar and Pickard, 1989) showed a lower response to gravity. The authors concluded that a full complement of starch is necessary for full gravitropic sensitivity (Kiss et al., 1989). However, these mutants can still sense gravity also more slowly and less accurately. According to Bandurski (1990) "if an organism has a dense and heavy statolith then it will use the statolith to provide a very accurate and rapid gravity sensor. If however it does not have such a dense body then the organism uses some more subtle gravity sensing apparatus". Bandurski's guess is then "that the plant uses its own bioelectric fields as a sensor". With his collaborators he had developed a working theory postulating that "the perception of the gravitational stimulus involves a perturbation of the plant's bioelectric field" and that the transduction of the stimulus involves a hormone-transport voltage-gating mechanism (Bandurski et al., 1986).

In the provoking suggestions concluding his recent review on "Plant Movements and the Cytoskeleton", Hensel (1989b) suggests that the cytoskeleton has a general function to generate and maintain polarity of root cap statocytes but that the cytoskeleton is "indirectly involved in perception by generating and maintaining a structural polarity of statocytes". Interestingly "it maintains domains of ion pumps/channels and/or hormone receptors/channels in the plasma membrane". The cortical part of the cytoskeleton would be directly involved in mechanotransduction of statolith weight into shear forces, thus triggering a plasma membrane response.

B. ANIMALS

Polar axiation in the eggs and embryos as well as the mechanisms underlying these processes in annelids, arthropods, amphibia and mammals are further discussed in a symposium on "Cellular Basis of Morphogenesis" published by Wolpert in 1989.

2. BIAXIAL PATTERNS: i) Mammals

Homologous gene clusters have been recently compared in insects and vertebrates. Specific homologues of *Antennapedia* (*Antp*)-like homeobox genes in *Drosophila* (see VIII.B.2d, in I) have been characterized as *Hox* complexes in vertebrates (Duboule *et al.*, 1986). Corresponding murine genes and insect complexes show the same relative boundary of the expression along the antero-posterior (A/P) axis of the developing embryo (Akam, 1989). A model for the mouse forelimb budding has been proposed by Dollé *et al.* (1989) that accounts for the establishment of the expression of the *Hox-5* domain in relation to the existence of a morphogen released by the zone of polarizing activity.

3. TRIAXIAL PATTERNS (left-right polarities)

Handedness is a fundamental quality already appreciated by D'Arcy Thompson (1942, see I).

a) Helical bacteria

The twist model of the lytic-deficient mutations of *Bacillus subtilis* has recently reactivated the handedness principle (Mendelson and Thwaites, 1989). Growth of these lytic-deficient mutants does not result in increased numbers of individual bacteria but in long thread-like clones which may have an unusual double-helical morphology. These double-helical threads fold repeatedly to form helical, multicellular "macrofibres" ("macrobes") that, according to Galloway (1990) are structurally analogous to twisted textile yarns. A macrobe is therefore an amplifier of the cell wall structure-determining features of the individual cells and therefore has a helical structure.

On the basis of screw sense, some strains are left-handed, others right. Others again are "conditional" mutants — they may be either left or right, and the degree of twist can vary continuously between left-handed and right-handed extremes depending on environmental factors, such as temperature (Galloway, 1990). Right-hand clones are produced at lower temperatures, left-hand at higher ones (Mendelson et al., 1984). It seems that a protein is needed for left-handed structures but not right-handed.

e) Molluscs

Interestingly, a same asymmetric behaviour as in bacteria is seen in the early development of snails: right-handedness in *Lymnaea peregra* needs a protein, left-

handedness apparently does not (see I). In the interplay between molecular self-assembly into helicoidal structures and mechanical reorientation due to growth forces (Neville, 1985; Galloway, 1990), a central role has been suggested to microtubules in the formation of helical patterns (Lloyd, 1984).

EPILOGUE (complement to pp. 271-273 in I)

The predictor question — What is life? — asked by the brilliant physicist Erwin Schrödinger in 1945 has since been partially answered by the cracking of the enigma of the genetic code. However, it still leaves open the question of "How does this one-dimensional code specify a three-dimensional organism?", a question relevant of topobiology (Edelman, 1988). At this epigenetic level, organizational principles of inanimate objects appear to be still valid even though complexified for animate ones. Preeminent among such universal principles is polarity emerged from the primary asymmetries of particulate matter (see I.B) and multi-expanded into the numerous biopolarities.

To bridge genetics and epigenetics still remains the great question of how genes control the transduction of the intrinsic molecular polarities into those cellular and organismic biopolarities? The bridge starts to be completed at the cellular level with the recent unravelling of genes controlling polarity of cytoskeletal macromolecules such as actin, myosin and tubulins (see IV.E), themselves someway related to known cell positioning as examplified by our *Allomyces* "sexual dipoles" (Plate I). However, the link remains elusive at the organismic level where some types of interaction should intervene between macromolecular polarities and DNA-controlled directional (head or foot in the Hydra model) morphogenetic gradients.

I renew my gratefulness to ARIANE FEHR for her trustful technical help.

REFERENCES

- ADAMS, R. J. and T. D. POLLARD (1989). Binding of myosin I to membrane lipids. Nature 340: 565-568.
- AGNEW, W.S. (1989). Excitation-contraction coupling: cloning of the SR foot. Nature 339: 422-423.
- AKAM, M. (1989). Hox and HOM: homologous gene clusters in insects and vertebrates. Cell 57: 347-349.
- ALLAN, V. J. and T. E. Kreis (1986). A microtubule-binding protein associated with membranes of the Golgi apparatus. J. Cell Biol. 103: 2229-2239.
- AMATRUDA, J. F., J. F. CANNON, K. TATCHELL, C. Hug and J. A. Cooper (1990). Disruption of the actin cytoskeleton in yeast capping protein mutants. *Nature* 344: 352-354.
- André, E., M. Brink, G. Gerisch, G. Isenberg, A. Noegel, M. Schleicher, J. E. Segall and E. Wallraff (1989). A *Dictyostelium* mutant deficient in severin, an F-actin fragmenting protein, shows normal motility and chemotaxis. *J. Cell Biol.* 108: 985-995.
- ARBER, W. (1974). DNA modification and restriction. Prog. Nucl. Acid Res. Mol. Biol. 14: 1-37.
- ARMSTRONG, J. and G. WARREN (1990). Membranes: violating the one-way system. Nature 344: 383-385.
- ARSCOTT, P. A., G. LEE, V.A. BLOOMFIELD and D. F. EVANS (1989). Scanning tunnelling microscopy of Z-DNA. *Nature* 339: 484-486.
- Assmann, S., L. Simoncini and J. I. Schroeder (1985). Blue light activates electrogenic ion pumping in guard cell protoplasts of *Vicia faba*. *Nature* 318: 285-287.
- BANDURSKI, R. S. (1990). Oral communication, Feb. 1990.
- BANDURSKI, R. S., A. SCHULZE and W. DOMAGALSKI (1986). Possible effects of organelle charge and density on cell metabolism. *Adv. Space Res.* 6(12): 47-54.
- BARNES, P., J. L. FINNEY, J. D. NICHOLAS and J. E. QUINN (1979). Cooperative effects in simulated water. *Nature* 282: 459-464.
- BECKER, A. and H. MURIALDO (1990). Bacteriophage λ DNA: the beginning of the end. *J. Bacteriol.* 172: 2819-2824.
- BECKERS, C. J. M., M. R. BLOCK, B. S. GLICK, J. E. ROTHMAN and W. E. BALCH (1989). Vesicular transport between the endoplasmic reticulum and the Golgi stack requires the NEM-sensitive fusion protein. *Nature* 339: 397-398.
- Bentrup, F.-W. (1989). Cell electrophysiology and membrane transport. Progr. Bot. 51: 70-79.
- BERGMAN, K. et al. (1969). Phycomyces. Bacteriol. Rev. 33: 99-157.
- BLACK, M. M. and R. J. LASEK (1980). Slow components of axonal transport: two cytoskeletal networks. J. Cell Biol. 86: 616-623.
- BLOW, D. (1990). More of the catalytic triad. Nature 343: 694-695.
- BOLOTIN, M., D. COEN, J. DEUTSCH, B. DUJON, P. NETTER, E. PETROCHILO and P. P. SLONIMSKI (1971). La recombinaison des mitochondries chez *Saccharomyces cerevisiae*. *Bull. Inst. Pasteur* 69: 215-239.
- BOSSHART, H., P. STRAEHL, B. BERGER and E. G. BERGER (1990). Brefeldin A induces a microtubule-directed backflow of Golgi to ER membranes. *Experientia* 46: S2 3.
- BOUCHIAT, A.-M. and L. POTTIER (1984). An atomic preference between left and right. *Sci. Amer.* 250(6): 76-85.
- BRACK, A. et al., interrogés par E. Noël (1989). La Symétrie Aujourd'hui. Sciences, éd. du Seuil, Paris. 238 pp.
- Bray, D. (1982). Filopodial contraction and growth cone guidance. In *Cell Behaviour*. Pp. 299-317. R. Bellair, A. Curtis and G. Dunn (eds). Cambridge University Press.

- BRAY, D. and J. VASILIEV (1989). Networks from mutants. Nature 338: 203-204.
- Bretscher, M.S. (1984). Endocytosis: relation to capping and cell locomotion. Science 224: 681-686.
- BRODSKY, M. H. (1990). Progress in gallium arsenide semiconductors. Sci. Amer. 262(2): 56-63.
- CACERES, A. and K.S. KOSIK (1990). Inhibition of neurite polarity by tau antisense oligonucleotides in primary cerebellar neurons. *Nature* 343: 461-463.
- CAILLOIS, R. (1973). La Dissymétrie. Gallimard, Paris.
- CANO, C., L. HERRERA-ESTRELLA and J. RUIZ-HERRERA (1988). DNA methylation and polyamines in regulation of development of the fungus *Mucor rouxii*. J. Bacteriol. 170: 5946-5948.
- CASPAR, T. and B. G. PICKARD (1989). Gravitropism in a starchless mutant of *Arabidopsis*. *Planta* 177: 185-197.
- CATTERALL, W.A., M. J. SEAGAR, M. TAKAHASHI and K. NUNOKI (1989). Molecular properties of dihydropyridine-sensitive calcium channels. In *Calcium Channels: Structure and Function*. D. W.-Wray, R. I. Norman and P. Hess (eds). *Ann. N.Y. Acad. Sci.* 560: 1-14.
- CHIU, S. W. and D. MOORE (1990). Sporulation in *Coprinus cinereus*: use of an *in vitro* assay to establish the major landmarks in differentiation. *Mycol. Res.* 94: 249-253.
- Cholodny, N. G. (1956). Selected works in 3 volumes. Vol. 1. Materials for Plant Electrophysiology. Pp. 358-374. Kiev Acad. Sci. URSS.
- CLARK, W. G. (1937). Electrical polarity and auxin transport. Plant Physiol. 12: 409-440.
- CLEGG, J.S. (1982). Alternative views on the role of water in cell function. In *Biophysics of Water*. Pp. 365-383. F.F. Franks and S. Mathias (eds), John Wiley and Sons, Chichester, New York. 400 pp.
- CLINE, D. B. (1988). Beyond truth and beauty: a fourth family of particles. Sci. Amer. 259(2): 42-49.
- COHAN, C. S. and S. B. KATTER (1986). Suppression of neurite elongation and growth cone motility by electrical activity. *Science* 232: 1638-1640.
- CRAIK, C. S., S. ROCZNIAK, C. LARGMAN and W. J. RUTTER (1987). The catalytic role of the active site of aspartic acid serine proteases. *Science* 237: 909-913.
- CROMBIE, T., N. A. R. Gow and G. W. GOODAY (1990). Influence of applied electrical fields on yeast and hyphal growth of *Candida albicans*. J. Gen. Microbiol. 136: 311-317.
- DAVIES, E. (1987). Action potentials as multifunctional signals in plants: a unifying hypothesis to explain apparently disparate wound responses. *Plant Cell Environ*. 10: 623-631.
- Dollé, P., J.-C. Izpisua-Belmonte, H. Falkenstein, A. Renucci and D. Duboule (1989). Coordinate expression of the murine *Hox-5* complex homeobox-containing genes during limb pattern formation. *Nature* 342: 767-772.
- DROST-HANSEN, W. and J. L. SINGLETON (1989). Liquid asset. How the exotic properties of cell water enhance life. *The Sciences* Sept./Oct.: 38-42.
- DRUBIN, D. G., J. MULHOLLAND, Z. ZHU and D. BOTSTEIN (1990). Homology of a yeast actin-binding protein to signal transduction proteins and myosin-I. *Nature* 343: 288-290.
- DUBOULE, D., A. BARON, P. MÄHL and B. GALLIOT (1986). A new homeo-box is present in overlapping cosmid clones which define the mouse *HOX*-1 locus. *EMBO J.* 5: 1973-1980.
- DUJON, B., P. P. SLONIMSKI and L. WEILL (1974). Mitochondrial genetics IX: A model for recombination and segregation of mitochondrial genomes in *Saccharomyces cerevisiae*. *Genetics* 78: 415-435.
- DYER, C. K. (1990). A novel thin-film electrochemical device for energy conversion. *Nature* 343: 547-548.
- EDELMAN, G. M. (1988). Topobiology. An Introduction to Molecular Embryology. Basic Books, Inc., Publishers, New York. 240 pp.
- ELLIS, H.W. and E. R. TURNER (1978). The effect of electricity on plant growth. Sci. Prog. 65: 395-407.

REFERENCES 361

- ELLIS, J. (1990). Deep resonance of CP violation. Nature 344: 197-198.
- ENOS, A.P. and N.R. MORRIS (1990). Mutation of a gene that encodes a kinesin-like protein blocks nuclear division in A. nidulans. Cell 60: 1019-1027.
- ERNST, R. R., G. BODENHAUSEN and A. WOKAUN (1987). Principles of Nuclear Magnetic Resonance in One and Two Dimensions. Oxford Scientific Publ. Clarendon Press.
- Errera, J. (1928). Polarisation Diélectrique. Presses Universitaires de France.
- (1935). Le Moment Electrique en Chimie (2 vol.). Hermann, Paris.
- FARLEY, F. W. and G. M. McClelland (1990). Orientational dielectric relaxation of collisionless molecules. *Science* 247: 1572-1575.
- Felle, H. (1989). pH as a second messenger in plants. In Second Messengers in Plant Growth and Development. Plant Biology, vol.6, pp. 145-166. W. F. Boss and D. J. Morré (eds), Alan R. Liss, New York. 348 pp.
- FENSOM, D. S. (1959). The bio-electrical potentials of plants and their functional significance. 3. The production of continuous potentials across membranes in plant tissue by the circulation of the hydrogen ion. *Can. J. Bot.* 37: 1003-1026.
- FIELD, M. J. and R. W. RICHARDSON (1989). Arch. rational Mech. Anal. 105: 91-94; cited by I. Stewart (1989). Symmetry breakthrough. Nature 341: 389-390.
- FINNEY, J.L. (1982). Towards a molecular picture of liquid water. In *Biophysics of Water*. Pp. 73-95. F.F. Franks and S. Mathias (eds), John Wiley and Sons, Chichester, New York. 400 pp.
- FORK, D. C. (1972). Oxygen electrode. Methods Enzymol. 24B: 113-122.
- FOUKAL, P.V. (1990). The variable sun. Sci. Amer. 262(2): 26-33.
- FUENTE, de la, R.K. (1984). Role of calcium in the polar secretion of indoleacetic acid. *Plant Physiol.* 76: 342-346.
- FUKUI, Y., T.J. LYNCH, H. BRZESKA and E.D. KORN (1989). Myosin I is located at the leading edges of locomoting *Dictyostelium* amoebae. *Nature* 341: 328-331.
- GALLAND, P. and T. OOTAKI (1987). Differentiation and cytology. In *Phycomyces*. Pp. 281-316. E. Cerdá-Olmedo and E. D. Lipson (eds). Cold Spring Harbor Laboratory. 430 pp.
- GALLOWAY, J. (1990). Putting a twist in the tale. Nature 343: 513-514.
- GARDNER, M. (1979). The Ambidextrous Universe: Mirror Asymmetry and Time-Reversed Worlds. Second and Uptdated Edition. Charles Scribner's Sons.
- GENNIS, R. B. (1989). Biomembranes. Molecular Structure and Function. Springer-Verlag, New York, Berlin. 533 pp.
- GOLDSWORTHY, A. (1986). The electrical compass of plants. New Sci. 109: 22-23.
- GÖTZE, H. (1918). In Galland and Ootaki, 1987.
- GOVINDJEE and W. J. COLEMAN (1990). How plants make oxygen. Sci. Amer. 262(2): 42-51.
- Gow, N. A. R. (1989). Circulating ionic currents in micro-organisms. Adv. Microb. Physiol. 30: 89-123.
- GRADMANN, D. (1989). Electrogenic chloride transport in Algae. In Bicarbonate, Chloride, and Proton Transport Systems. J. H. Durham and M. A. Hardy (eds). Ann. N.Y. Acad. Sci. 574: 20-29.
- GRESIK, M., N. KOLAROVA and V. FARKAS (1988). Membrane potential, ATP, and cyclic AMP changes induced by light in *Trichoderma viride*. Exper. Mycol. 12: 295-301.
- GROLIG, F. and G. WAGNER. (1988). Light-dependent chloroplast reorientation in *Mougeotia* and *Mesotaenium*: Biased by pigment-regulated plasmalemma anchorage sites to actin filaments? *Bot. Acta* 101: 2-6.
- HARGITTAI, I. and M. HARGITTAI (1986). Symmetry Through the Eyes of a Chemist. VCH. 458 pp.

- HEATH, I.B. and S.G.W. KAMINSKYJ (1989). The organization of tip-growth-related organelles and microtubules revealed by quantitative analysis of freeze-substituted Oomycete hyphae. *J. Cell Sci.* 93: 41-52.
- HEGSTROM, R. A. and D. K. KONDEPUDI (1990). The handedness of the Universe. Sci. Amer. 262 (1): 98-105.
- HENSEL, W. (1986). Cytodifferentiation of polar plant cells. Use of anti-microtubular agents during the differentiation of statocytes from cress roots (*Lepidium sativum L.*). *Planta* 169: 293-303.
- (1989a). Physiology of movements in space experiments. *Progr. Bot.* 50: 158-162.
- (1989b). Plant movements and the cytoskeleton. Progr. Bot. 51: 165-172.
- HICKS, G. R., D. L. RAYLE and T. L. LOMAX (1989). The *Diageotropica* mutant of tomato lacks high specific activity auxin binding sites. *Science* 245: 52-54.
- Ho, W. C., V. J. Allan, G. van Meer, E. C. Berger and T. E. Kreis (1989). Reclustering of scattered Golgi elements occurs along microtubules. *Eur. J. Cell Biol.* 48: 250-263.
- HOCH, H.C., B. E. TUCKER and R. C. STAPLES (1987). An intact microtubule cytoskeleton is necessary for mediation of the signal for cell differentiation in *Uromyces. Eur. J. Cell Biol.* 45: 209-218.
- HOHL, H.R. (1987). Cytology and morphogenesis of fungal cells. Progr. Bot. 49: 13-28.
- (1989). Cytology and morphogenesis of fungal cells. *Progr. Bot.* 51: 1-20.
- HOLLENBECK, P. J. (1990). Cytoskeleton on the move. Nature 343: 408-409.
- HOLMSTEDT, B., H. Frank and B. Testa (eds) (1989). Chirality and Biological Activity. John Wiley and Sons, New York. 300 pp.
- JAEGER, F. M. (1925). Principe de Symétrie. Gauthier-Villars.
- JAFFE, L.F. and R. NUCCITELLI (1977). Electrical control of development. Ann. Rev. Biophys. Bioeng. 6: 445-476.
- JOHNSON, P. F. and S. L. McKnight (1989). Eukaryotic transcriptional regulatory proteins. *Ann. Rev. Biochem.* 58: 799-839.
- KIKERI, D., A. SUN, M.L. ZEIDEL and S.C. HEBERT (1989). Cell membranes impermeable to NH₃. *Nature* 339: 478-480.
- KISS, J. Z., R. HERTEL and F. D. SACK (1989). Amyloplasts are necessary for full gravitropic sensitivity in roots of *Arabidopsis thaliana*. *Planta* 177: 198-206.
- KNECHT, D. A. and W. F. LOOMIS (1987). Antisense RNA inactivation of myosin heavy chain gene expression in *Dictyostelium discoideum*. Science 236: 1081-1086.
- KÖHLER, K., W. STEIGNER, W. SIMONIS and W. URBACH. (1985). Potassium channels in *Eremosphaera* viridis. I. Influence of cations and pH on resting membrane potential and on an action potential-like response. *Planta* 166: 490-499.
- KÖHLER, K., W. STEIGNER, J. KOLBOWSKI, U. P. HANSEN, W. SIMONIS and W. URBACH (1986). Potassium channels in *Eremosphaera viridis*. II. Current- and voltage-clamp experiments. *Planta* 167: 66-75.
- KOSIK, K. S. and E. A. FINCH (1987). MAP2 and tau segregate into dendritic and axonal domains after the elaboration of morphologically distinct neurites: an immunocytochemichal study of cultured rat cerebrum. *J. Neurosci.* 7: 3142-3153.
- KOSTYUK, P., N. AKAIKE, Y. OSIPCHUK, A. SAVCHENKO and Y. SHUBA (1989). Gating and permeation of different types of Ca channels. In *Calcium Channels: Structure and Function*. D.W. Wray, R. I. Norman and P. Hess (eds). *Ann. N.Y. Acad. Sci.* 560: 63-79.
- Kristen, U. (1987). General and molecular cytology: the cytoskeleton: microfilaments. *Progr. Bot.* 49: 1-12.
- LAMOUREUX, P., R. E. BUXBAUM and S. R. HEIDEMANN (1989). Direct evidence that growh cone pull. *Nature* 340: 159-162.

REFERENCES 363

- L'ANGMUIR, I. (1932). Cited by Peratt, 1990.
- LANDIS, S. C. (1983). Neuronal growth cones. Ann. Rev. Physiol. 45: 567-580.
- LASEK, R. J. (1986). Polymer sliding in axons. J. Cell. Sci. (suppl.) 5: 161-179.
- LASEK, R. J. and M. M. BLACK (eds) (1988). Intrinsic Determinants of Neuronal Form and Function. Neurology and Neurobiology, vol. 37. Alan R. Liss, New York. 591 pp.
- LEE, J., M. GUSTAFSSON, K.-E. MAGNUSSON and K. JACOBSON (1990). The direction of membrane lipid flow in locomoting polymorphonuclear leukocytes. *Science* 247: 1229-1233.
- LENK, R., M. Bonzon and H. Greppin (1980). Dynamically oriented biological water as studied by NMR. *Chem. Phys. Lett.* 76: 175-177.
- Letourneau, P. C. (1982). Nerve fiber growth and its regulation by extrinsic factors. In *Neuronal Development*. Pp. 213-254. N. C. Spitzer (ed.). Plenum, New York.
- LINDEN, E. (1990). Can we really understand matter? Time, April 16, p. 41.
- LISANTI, M. P. and E. RODRIGUEZ-BOULAN (1990). Glycophospholipid membrane anchoring provides clues to the mechanism of protein sorting in polarized epithelial cells. *Trends Biochem. Sci.* 15: 113-118.
- LLOYD, C.W. (1984). Toward a dynamic helical model for the influence of microtubules on wall patterns in plants. *Inter. Rev. Cytol.* 86: 1-54.
- LUCKEN, E. A. C. (1969). Nuclear Quadrupole Constants. Academic Press, London, New York. 360 pp.
- Lund, E. J., R. I. Mahan and A. H. Hanszen (1947). Reversible total inhibition of polar growth in the root by an applied electric current and the nature of this control. In *Bioelectric Fields and Growth*. Pp. 186-197. E.J. Lund. (ed.). Univ. Press Austin, Tx., U.S.A.
- MARSH, L. and P. C. LETOURNEAU (1984). Growth of neurites without filipodial or lamellipodial activity in the presence of cytochalasin B. J. Cell Biol. 99: 2061-2067.
- McIntosh, J. R. and M. P. Koonce (1989). Mitosis. Science 246: 622-628.
- McIntosh, J. R. and K. L. McDonald (1989). The mitotic spindle. Sci. Amer. 261(4): 26-34.
- McLaughlin, S. (1977). Electrostatic potentials at membrane-solution interfaces. Current Topics in Membranes and Transport 9: 71-144.
- MEDVEDEV, S. S. and I. V. MARKOVA (1990). How can the electrical polarity of axial organs regulate plant growth and IAA transport? *Physiol. Plant.* 78: 38-42.
- MELUH, P.B. and M.D. Rose (1990). KAR3, a kinesin-related gene required for yeast nuclear fusion. Cell 60: 1029-1041.
- MENDELSON, N. H. and J. J. THWAITES (1989). Do forces and the physical nature of cellular materials govern biological processes? *Comments Theor. Biol.* 1: 217-236.
- MENDELSON, N. H., D. FAVRE and J. J. THWAITES (1984). Twisted states of *Bacillus subtilis* macrofibers reflect structural states of the cell wall. *Proc. Natn. Acad. Sci. U.S.A.* 81: 3562-3566.
- MENZEL, D. and C. ELSNER-MENZEL (1989). Actin-based chloroplast rearrangements in the cortex of the giant coenocytic green alga *Caulerpa*. *Protoplasma* 150: 1-8.
- MITCHISON, T. and M. KIRSCHNER (1988). Cytoskeletal dynamics and nerve growth. Neuron 1: 761-772.
- MONNIER, D., W. HAERDI, J. BUFFLE and Y. RUSCONI (1979). Chimie Analytique. Georg, Genève. 293 pp.
- Moore, R., C.E. McClelen, W.M. Fondren and C.L. Wang (1987). Influence of microgravity on root-cap regeneration and the structure of columella cells in *Zea mays. Am. J. Bot.* 74: 218-223.
- Moore, R., M. L. Evans and W. M. Fondren (1990). Inducing gravitropic curvature of primary roots of *Zea mays* cv ageotropic. *Plant Physiol.* 92: 310-315.
- NATHANS, D. and H.O. SMITH (1975). Restriction endonucleases in the analysis and restructuring of DNA molecules. *Ann. Rev. Biochem.* 44: 273-293.

- NEVILLE, A. C. (1985). Molecular and mechanical aspects of helicoidal development in plant cell walls. *BioEssays* 3: 4-8.
- NICOLLE, J. (1950). La Symétrie et ses Applications. Albin Michel, Paris.
- NOVAK, V.A. and N.G. IVANKINA (1983). Influence of nitro blue tetrazolium on the membrane potential and ion transport in water thyme. *Sov. Plant. Physiol.* 30: 845-853. (English translation).
- NOVICK, P. and D. BOTSTEIN (1985). Phenotypic analysis of temperature-sensitive yeast actin mutants. *Cell* 40: 405-416.
- OCHS, S. (1975). In *Recent Advances in Myology*. Pp. 189-194. W. G. Bradley *et al.* (eds). Excerpta Medica, Amsterdam. 578 pp.
- OKABE, S. and N. HIROKAWA (1990). Turnover of fluorescently labelled tubulin and actin in the axon. *Nature* 343: 479-482.
- OPPOLZER, W. (1987). Camphor derivatives as chiral auxiliaries in asymmetric synthesis. *Tetrahedron* 43: 1969-2004.
- OSIEWACZ, H. D. and U. HEINEN (1989). Recombination of mobile genetic elements from plants and cyanobacteria. *Progr. Bot.* 50: 174-197.
- PABO, C.O., A. K. AGGARWAL, S. R. JORDAN, L. J. BEAMER, U. R. OBEYSEKARE and S. C. HARRISON (1990). Conserved residues make similar contacts in two repressor-operator complexes. *Science* 247: 1210-1213.
- PARFYONOV, G. P. et al. (1979). Biological investigations aboard biosatellite cosmos-782. Acta Astronaut. 6: 1235.
- Pariselle, H. (1936). Polarimétrie et Chimie. Gauthier-Villars. Paris
- PERATT, A. L. (1990). Not with a bang. The Sciences Jan./Feb.: 24-32.
- PERBAL, G. and D. DRISS-ECOLE (1989). Polarity of statocytes in lentil seedling roots grown in space. (Spacelab D1 Mission). *Physiol. Plant.* 75: 518-524.
- PICKARD, B. G. (1973). Action potentials in higher plants. Bot. Rev. 39: 172-201.
- PILET, P.-E. (1978). The role of the cap in the geotropism of roots exposed to light. Z. Pflanzenphysiol. 89: 411-426.
- PONT, V. and R. Pezet (1990). Relation between the chemical structure and the biological activity of hydroxystilbenes against *Botrytis cinerea*. *J. Phytopathol*. (in press).
- POOL, R. (1989). Chaos in a hydrogen atom. Science 243: 894.
- PRICE, A. H. (1969). In *Dielectric Properties and Molecular Behavior*. Pp. 191-231. N.E. Hill (ed.). Van Nostrand Reinhold, New York.
- QUATRANO, R. F. and D. L. KROPF (1989). Polarization in *Fucus* (Phaeophyceae) zygotes: investigations of the role of calcium, microfilaments and cell wall. In *Algae as Experimental Systems*. Plant Biology, vol. 7, pp. 111-119. A. W. Coleman, L. J. Goff and J. R. Stein-Taylor (eds). Alan R. Liss, New York. 333 pp.
- QUIOCHO, F. A., D. K. WILSON and N. K. VYAS (1989). Substrate specificity and affinity of a protein modulated by bound water molecules. *Nature* 340: 404 and 732.
- REEDY, M. C., C. BEALL and E. FYRBERG (1989). Formation of reverse rigor chevrons by myosin heads. *Nature* 339: 481-483.
- REES, M. J. (1990). "Dead quasars" in nearby galaxies? Science 247: 817-823.
- RICHARD, E.A. and C. MILLER (1990). Steady-state coupling of ion-channel conformation to a transmembrane ion gradient. *Science* 247: 1208-1210.
- RIEZMAN, H. (1985). Endocytosis in yeast: Several of the yeast secretory mutants are defective in endocytosis. *Cell* 40: 1001-1009.

REFERENCES 365

- ROBERTS, R. G. (1990). Quark spins open to question. Nature 344: 708-709.
- RUTHERFORD, A.W. (1989). Photosystem II, the water-splitting enzyme. *Trends Biochem. Sci.* 14: 227-232.
- SAENGER, W. (1987). Structure and dynamics of water surrounding biomolecules. *Ann. Rev. Biophys. Biophys. Chem.* 16: 93-114.
- Salisbury, J. L. (1989). Algal centrin: calcium-sensitive contractile organelles. In *Algae as Experimental Systems*. Plant Biolology, vol. 7, pp. 19-37. A. W. Coleman, L. J. Goff and J. R. Stein-Taylor (eds). Alan R. Liss, New York. 333 pp.
- SCHATZ, G. (1989). Cell biology: biological cold fusion. Nature 339: 336.
- Schekman, R. and P. Novick (1982). The secretory process and yeast cell surface assembly. In *The Molecular Biology of the Yeast Saccharomyces cerevisiae:* metabolism and gene expression. Pp. 361-393. J. N. Strathern, E. W. Jones and J. R. Broach (eds). Cold Spring Harbor Laboratory, New York.
- Schleicher, M., A. Noegel, T. Schwarz, E. Wallraff, M. Brink, J. Faix, G. Gerisch and G. Isen-Berg (1988). A *Dictyostelium* mutant with severe defects in α-actinin: its characterization using cDNA probes and monoclonal antibodies. *J. Cell Sci.* 90: 59-71.
- Schnell, L. and M.E. Schwab (1990). Axonal regeneration in the rat spinal cord produced by an antibody against myelin-associated neurite growth inhibitors. *Nature* 343: 269-272.
- Schrödinger, E. (1945). What is life? The Physical Aspect of the Living Cell. Cambridge University Press; Macmillan, New York.
- SCHWARZ, G. (1978). On the physico-chemical basis of voltage-dependent molecular gating mechanisms in biological membranes. *J. Membrane Biol.* 43: 127-148.
- Scott, B. I. H. (1967). Electrical fields in plants. Ann. Rev. Plant Physiol. 18: 409-418.
- Scott, I.M. (1990). Plant hormone response mutants. Physiol. Plant. 78: 147-152.
- SCOTT, J. F. and C. A. PAZ DE ARAUJO (1989). Ferroelectric memories. Science 246: 1400-1405.
- SEELIG, J., P. M. MACDONALD and P. G. SCHERER (1987). Phospholipid head groups as sensors of electric charges in membranes. *Biochemistry* 26: 7535-7541.
- SHEETZ, M. P., S. TURNEY, H. QIAN and E. L. ELSON (1989). Nanometre-level analysis demonstrates that lipid flow does not drive membrane glycoprotein movements. *Nature* 340: 284-288.
- SHIMAZAKI, K., M. IINO and E. ZEIGER (1986). Blue light-dependent proton extrusion by guard-cell protoplasts of *Vicia faba*. *Nature* 319: 324-326.
- SHPETNER, H. S. and R. B. VALLEE (1989). Identification of dynamin, a novel mechanochemical enzyme that mediates interactions between microtubules. *Cell* 59: 421-432.
- Shropshire, W. Jr. and J.-F. Lafay (1987). Sporangiophore and mycelial responses to stimuli other than light. In *Phycomyces*. Pp. 127-154. E. Cerdá-Olmedo and E. D. Lipson (eds). Cold Spring Harbor Laboratory. 430 pp.
- SIEVERS, A., D. VOLKMANN, W. HENSEL, V. SOBICK and W. BRIEGLEB (1976). Cell polarity in root statocytes in spite of simulated weightlessness. *Naturwiss*. 43: 343.
- Simons, P. J. (1981). The role of electricity in plant movements. New Phytol. 87: 11-37.
- SILVER, R. A., A. G. LAMB and S. R. BOLSOVER (1990). Calcium hotspots caused by L-channel clustering promote morphological changes in neuronal growth cones. *Nature* 343: 751-754.
- SKLIFKIN, M. A. (1980). The significance of charge transfer interactions in biology. In *Molecular Interactions*, vol. 2, pp. 271-304. H. Ratajezak and W. J. Orville-Thomas (eds). John Wiley and Sons, New York.
- SMEEKENS, S., P. WEISBEEK and C. ROBINSON (1990). Protein transport into and within chloroplasts. Trends Biochem. Sci. 15: 73-76.

- SMITH, K.F. et al. (1990). A search for the electric dipole moment of the neutron. *Phys. Lett.* B 234: 191-196.
- SNYDER, M. (1989). The SPA2 protein of yeast localizes to sites of cell growth. J. Cell Biol. 108: 1419-1429.
- TAKESHIMA, H., S. NISHIMURA, T. MATSUMOTO, H. ISHIDA, K. KANGAWA, N. MINAMINO, H. MATSUO, M. UEDA, M. HANAOKA, T. HIROSE and S. NUMA (1989). Primary structure and expression from complementary DNA of skeletal muscle ryanodine receptor. *Nature* 339: 439-445.
- THEIMER, R., R. A. KUDIELKA and I. RÖSCH (1986). Induction of somatic embryogenesis in anise in microgravity. *Naturwiss*. 73: 442-443.
- THOMAS, J. O. and G. C. FARRINGTON (1982). Protonic beta-aluminas: model systems for proton transfer in biological processes. In *Biophysics of Water*. Pp. 215-217. F. F. Franks and S. Mathias (eds), John Wiley and Sons, Chichester, New York. 400 pp.
- TOCANNE, J.-F. and J. TEISSIÉ (1990). Ionization of phospholipids and phospholipid-supported interfacial lateral diffusion of protons in membrane model systems. *Biochim. Biophys. Acta* 1031: 111-142.
- TOYOSHIMA, Y. Y., C. TOYOSHIMA and J. A. SPUDICH (1989). Bidirectional movement of actin filaments along tracks of myosin heads. *Nature* 341: 154-156.
- TRINKAUS, J. P. (1985). Further thoughts on directional cell movement during morphogenesis. J. Neurosci Res. 13: 1-19.
- TURIAN, G. (1989). Polarity. From Dipoles to Biopolarizations. Arch. Sci. Genève 42: 1-323.
- (1990). Le couple divin (bipôles de vie). Campus (Uni-Genève) 1: 14-15.
- VALE, R. D. and L. S. B. GOLDSTEIN (1990). One motor, many tails: an expanding repertoire of force-generating enzymes. *Cell* 60: 883-885.
- Waaland, S. D. (1989). Cellular morphogenesis in the filamentous red alga *Griffithsia*. In *Algae as Experimental Systems*. Plant Biology, vol. 7, pp. 121-134. A.W. Coleman, L.J. Goff and J.R. Stein-Taylor (eds). Alan R. Liss, New York. 333 pp.
- WALLRAFF, E., M. SCHLEICHER, M. MODERSITZKI, D. RIEGER, G. ISENBERG and G. GERISCH (1986). Selection of *Dictyostelium* mutants defective in cytoskeletal proteins: use of an antibody that binds to the ends of α-actinin rods. *EMBO J.* 5: 61-67.
- WARSHEL, A., G. NARAY-SZABO, F. SUSSMAN and J.-K. HWANG (1989). How do serine proteases really work? *Biochemistry* 28: 3629-3637.
- WEINTRAUB, H.M. (1990). Antisense RNA and DNA. Sci. Amer. 262(2): 34-40.
- Welsh, M. J., M. Li, J. D. McCann, J. P. Clancy and M. P. Anderson (1989). Phosphorylation-dependent regulation of apical membrane chloride channels in normal and cystic fibrosis airway epithelium. In *Bicarbonate, Chloride, and Proton Transport Systems*. J. H. Durham and M. A. Hardy (eds). *Ann. N. Y. Acad. Sci.* 574: 44-51.
- WEYL, H. (1952). Symmetry. Princeton University Press, New Jersey, U.S.A. 168 pp.
- Wikström, M. (1989). Identification of the electron transfers in cytochrome oxidase that are coupled to proton-pumping. *Nature* 338: 776-778.
- WILLIAMS, E. R. (1988). The electrification of thunderstorms. Sci. Amer. 259(5): 48-65.
- WILLIAMS, R. J. P. (1989). Electron transfer: Energizing protons in membranes. *Nature* 338: 709-710.
- WOLPERT, L. (1989). Cellular Basis of Morphogenesis. Ciba Foundation Symposium 144. John Wiley and Sons. 280 pp.
- WRAY, D.W., R. I. NORMAN and P. HESS (eds) (1989). Calcium Channels: Structure and Function. Ann. N.Y. Acad. Sci., vol. 560. 475 pp.

REFERENCES 367

- ZATSEPINA, G. I. and J. B. TSAPLEV (1980). The nature of electric polarity in higher plants. *Biofizika* 25: 144-147 (In Russian).
- ZIMÁNYI, L. and J. K. LANYI (1989). Halorhodopsin: a light-driven active chloride transport system. In *Bicarbonate, Chloride, and Proton Transport Systems*. J. H. Durham and M. A. Hardy (eds). *Ann. N. Y. Acad. Sci.* 574: 11-19.

SUBJECT INDEX

Abiotic, 46 Acetone, 39 Achiral, 45 Acidic, 40, 62, 82, 101-103, 112, 167, 168, 218, 339 compartment, 167, 175, 185, 220 hyaloplasm, 154 pH, 235 polysaccharide, 207 protein, 154 Acidification, 101, 108, 151, 175, 186, 236 Acridine(s), 198, 199 Actin, 15, 38, 80, 81, 96, 112-118, 120, 125,	Agglutinin, 79, 80 Aggregates-aggregation, 68-71, 87, 96, 118, 137, 169, 209, 226, 228, 243, 265, 337 Agravitropism, 355 Algal axiation-differentiation, 195, 352 eggs-embryos, 157-159, 229, 352 elongation, 184 exosporulation, 195 rhizoids, 168 zygote, 171, 207 Algorithm, 61
129, 130, 132-136, 139, 144, 145, 152-154, 163, 167, 168, 177, 179, 180, 187, 200, 211, 218, 325, 344-347, 349, 350, 358 binding site, 115, 118, 344, 348 cable(s), 118, 130, 346 cortex, 180 dots, 151-154 F-, 111, 113, 114, 117, 145, 174, 346, 347	Amines, 40, 41, 95 Amino acids, 40, 41, 44-46, 57, 58, 60, 66, 67, 73, 81, 86, 95, 96, 167, 168, 249, 328, 334, 337 polar, 62 Ammonia (-ium), 68, 243, 353 Amoeboid (Amoeba), 15, 137-139, 141 cells, 147, 347 motion, 133-137, 346, 347 movement, 134, 135, 346
G-, 111, 114 G-F, 133	Amphipathic, 37, 40, 63, 65, 76-78, 80 Amphitropic, 77
microfilaments, 152, 167, 173, 174, 179, 207, 344, 348 molecule(s), 112, 114, 350 polymerization, 179 translocation, 180 Actin-binding protein(s), 115, 344, 348 Actin-like filaments, 154 Actin-mediated transport, 183	Ampholytes, 41 Amphoteric, 38, 41 Amyloplasts, 168, 236, 344, 354 Anaphase, 120, 125, 126 Anchorage, 63, 76, 81, 115, 153, 222, 339 Animal(s), 211, 212, 214-217, 223, 246, 250, 253, 255, 272, 325, 342, 353, 355
Action potential(s), 87, 88, 90-94, 341, 342 Activation, 52, 60, 64, 67, 80, 92, 102, 117, 127, 145, 148, 158, 162, 189, 198, 217, 222, 232, 243, 247, 248, 250, 341 Actomyosin, 89, 133, 134, 154, 346 Adhesion, 81, 115, 134, 136, 181 Affinity, 39, 40, 42, 60, 76, 106, 114, 194, 347	cell(s), 189, 211, 223, 262, 353 eggs-embryo 13, 211, 212, 225, 238, 325 gradient, 217, 255 pole, 190, 212, 217, 254, 255, 263 regeneration, 13 Animal-vegetal (A/V) axis, 212, 216, 253, 254, 262 gradient(s), 264, 265 halves, 255

polarity, 211, 214, 216, 217, 253, 254,	dominance, 164, 165, 231, 242
262, 263, 269	growth, 157, 162, 344, 345
Anion(s), 30, 91, 220, 339	meristem(s) 224, 229, 231, 238, 239
Anisometry, 39, 70, 150, 181	pattern(s), 189, 236
Anisotropy, 24, 185, 111, 112, 333, 338	pole(s), 162, 170, 218-220, 223, 225,
Ankyrin, 112, 221	248, 256, 266, 267, 272
Annihilation, 17, 21, 22	targeting signal, 348
Anode, 180	zone, 100, 163, 164, 168, 169, 174, 175,
Antenna, 108-111, 343	231
Anterio-posterior (A/P)	Apical and basolateral
axis, 245, 249, 250, 252, 256, 259, 260,	compartments (domains), 219, 220,
265, 267-270, 356	226, 353
differentiation, 228	gradients, 104, 171, 172
gradients, 264	poles, 218, 219, 248, 352
pattern(s), 240, 257	surfaces, 220, 221
polarities, 213, 241-249, 250-266,	Apolar (nonpolar), 37-40, 43, 50, 55, 61,
267-270	67, 75, 97, 147, 201, 264, 266, 271,
regeneration, 262	325
Anterior	cells, 14, 79, 87, 158, 160, 209, 264, 266,
pole, 213, 228, 229, 257-260, 263	271
segment, 256	egg, 158
structures, 259	growth, 149, 160
tip, 243, 257	Archegonia, 174, 197
Anterograde (inward) transport, 122,	Ascospores, 155
177-179	Assembly, 57, 69, 70, 99, 111, 114, 117,
Antheridia, 174, 197	119-123, 125, 126, 132, 179, 202,
Antibiotic, 86, 151, 183, 343	203, 219
Antibodies, 66, 115, 204, 220, 249, 353	Asymmetric
Antigenes, 66, 67, 150, 243	budding, 219
Antimatter, 18, 19, 21, 47, 327, 328	carbon, 32
Antiparallel, 25, 29, 49, 50, 64, 68, 86, 87,	distribution-transport, 80, 218, 339,
117, 120, 345	341, 345, 349, 350
Antiport, 82	division(s), 149, 150, 161, 189, 192,
Antisense, 58, 336, 350	200-202, 206, 210, 211, 222, 225,
Antisera, 267	229, 230, 252
Apex, 64, 129, 164, 165, 166, 169, 170, 172,	growth, 236
175, 180, 193-195, 197, 224, 225,	septation, 190, 191, 202, 205
227, 232, 238, 239, 241	structure(s), 74, 82, 202
Apical, 13, 100, 104, 137, 169-173, 208,	synthesis, 46, 335
218, 221, 225, 226, 231, 242, 271,	Asymmetry, 13, 18, 35, 44, 46-48, 50-52,
348, 352, 353	64, 66, 74, 77, 79, 82, 100, 102, 108,
axis, 184	109, 112, 114, 119, 123, 132, 145,
bud, 232	154, 159, 173, 181, 186, 194, 200,
cell(s), 160, 169, 172-174, 197, 205, 209,	205, 206, 208, 210, 213, 216, 225,
211, 268	230, 236, 237, 247, 266-269, 326,
differentiation(s), 192-221, 352, 353	327, 328, 334-336, 339, 341, 345,
division, 173	356, 358
dome, 194	Atomic, 17-32, 325-331

ATP, 81, 83-85, 100, 104, 106-110, 114,	shape, 150, 181
116, 117, 120, 122, 123, 128, 132,	sporulation (endospores), 190, 191
135, 140, 142, 143, 160, 167, 179,	Bacteriochlorophyll, 43
186, 192, 218, 327, 333, 338, 340,	Bacteriorhodopsin, 109, 142
345, 346	Band(s), 104, 110, 117, 130, 136, 148, 169,
pump, 106	172, 185, 195, 215
synthesis, 83-85, 339	Barbed end(s), 114, 130, 145
ATPase, 82-85, 90, 91, 102, 106, 110, 115,	Basal, 68, 122, 131, 137-139, 145, 147, 155,
118, 121-123, 126, 128, 129, 167,	160, 169-171, 193, 207, 208, 218,
168, 220, 345	225, 233, 235, 242, 244, 247, 249,
Auxin(s), 159, 160, 186, 222, 229-231,	255, 256
233-238, 340, 354, 355	bodies, 104, 244-246, 256
transport, 229, 230, 233, 235, 341, 354	cell, 204, 211, 225
Auxospores, 147, 184	compartment, 205
Axis-axes (axiation), 13, 21, 22, 29, 31, 32,	permeability, 235
47, 50, 60, 64, 71, 79, 86, 104, 105,	pole(s), 158, 170, 181, 221, 225
107, 123, 130, 132, 137, 144-148,	rhizoidal, 208
154, 157, 165, 170, 182, 188-190,	zone, 163, 164, 170
192, 198, 200, 201, 206, 207, 213,	Basidia, 227, 228, 352
216, 221-224, 227, 228, 231, 233,	Basidiocarp, 227
238, 241, 244, 246-250, 262-265,	Basidiospores, 155, 194, 195, 226, 352
271, 352-354, 356	Basolateral, 218, 220
division, 209	cell surface, 219-221
growth, 147, 158, 161, 165	domain, 220
polarity, 148, 154, 161, 185, 192, 200,	membrane, 98, 218, 220, 221, 267
205, 212, 213, 222, 227, 228, 233,	pumps, 220
239, 242, 248, 262, 266, 272, 348,	Bending, 50, 131, 168, 169, 174
354, 355	Benzene, 39, 43, 67
proximal-distal, 261	Biaxial patterns, 250-267, 356
Axial, 66, 133, 174, 224, 247, 249, 356	Bidirectional, 102, 123, 179
asymmetry, 13	Bifurcation, 143
development, 224	Big Bang, 17, 19, 327
head activator gradient, 248	Bilayer(s), 63, 75-78, 107, 333, 338, 340
regeneration, 13	Binary, 23, 26, 272, 330
symmetry, 64, 221, 249, 265	fission, 149, 150, 182, 201
Axon(s), 90-92, 94, 105, 122, 123, 177-181,	Binding, 17, 24, 33, 34, 39, 50, 52, 57, 60,
349-351	64, 66-70, 77, 80, 86, 90, 93, 96, 112,
cytoskeleton, 350	114, 123, 140, 337, 344
transport, 122, 177, 178, 350	protein(s), 60, 114, 337
Axoneme, 122	site, 66, 111
Axoplasm, 123, 179	Bioelectric field(s), 237, 354, 355
Pacterial 120 122 140 167 181 356	potentials, 166
Bacterial, 129-132, 140, 167, 181, 356	Bioelectrochemistry, 110
budding, 149, 150	Biosynthesis, 62, 73, 129, 155, 166
cell(s), 53, 182, 190	Biopolarity, 14, 273, 326, 358
chromosome, 183 division, 190	Bipolar, 13, 22, 64, 72, 107, 113, 115-118,
elongation, 167, 182	124, 130, 167, 170, 173, 181-186,
helical, 267, 356	195, 196, 202, 208, 209, 229, 247,
ilelical, 201, 330	

259, 271, 272, 325, 330, 348	channel(s), 89, 90, 175, 207, 216, 341,
axiation, 147, 181, 186, 192, 197, 204	342
	currents, 216
budding, 184, 187	gradients, 175, 262, 349
couple, 17, 21	ionophore, 209, 254, 261
differentiation(s), 181, 202, 204, 205,	pump, 106
229	transport, 106, 342, 349
field(s), 41, 222, 232, 240, 247	uptake, 106
filaments, 116, 117	Calcium, 23, 80, 88, 89, 93, 95, 102, 106,
germination, 184	116, 135, 139, 160, 237, 253, 254,
gradient, 200	340-342, 349
growth, 167, 181-186, 271	Calcofluor, 153
mating systems, 195	Callus, 147, 239, 354
mitochondrion, 107	Calmodulin, 95, 116, 139, 142, 237
pattern(s), 41, 195-201, 332, 336	Cambium, 189
regeneration, 251, 262	Cancer, 148
segregation, 184, 199	Capping, 80, 81, 113, 127
sexualization, 195, 196, 200	protein(s), 114, 344
Bipolarity, 22, 41, 70, 108, 113, 116, 117,	Capsid(s), 69
119, 183, 185, 200, 204	Carbohydrate(s), 13, 17, 72, 75, 78, 100,
Bipolarization, 70, 257, 258, 271, 272	109, 355
electric, 19-22, 330	Carbon, 38-40, 44, 60, 67, 75, 192, 248, 334
Bipolaron, 24	Carotene, 42, 158, 198, 199, 228
Blastocyst, 266, 267	Carotenoids, 109, 161
Blastomere(s), 252-254, 262, 264, 266, 267	Catalysis (-yst), 45, 56, 339
Blue light, 31, 169, 174, 209, 340	Cathode, 74, 180, 188, 348, 349
Bond(s), 24, 35, 39-41, 55, 59, 60, 62,	Cation(s), 23, 24, 30, 37, 38, 78, 81, 86, 90,
64-66, 68, 92, 332-335	92, 99, 135, 143, 338
Boson(s), 19, 24, 27, 29, 44, 327	Caulonema, 172
Branching, 93, 164, 165, 169, 195, 228, 231	Causal, 214, 217, 231, 255
Brevin, 113	CCl ₄ , 38, 39
Bridges, 45, 70, 115, 122, 345, 353	Cell(s), 13-15, 23, 30, 34, 38, 46, 52, 62, 64,
Bristle(s), 176, 261	68, 72-74, 77-81, 83, 85, 87-105,
Bud(s), 95, 97, 101, 103, 149-154, 163, 172,	107-109, 111-115, 117-119, 121-127,
184, 187, 231-233, 239, 348	129-154, 157, 160-166, 170-193, 196,
growth, 150-154	197, 200-226, 228-230, 232, 233,
meristem, 238	235-241, 243-247, 249-252, 254, 257,
polarity, 152	259, 261-268, 271, 272, 325, 326,
Budding, 102, 149-151, 153, 154, 162, 184,	333, 338, 342-348, 352-354, 358
187, 196, 246, 247, 342, 344, 348,	adhesion, 82
356	asymmetry, 205
bacterial, 149, 150	axes, 175, 190, 271, 346
forelimb, 265, 356	compartment(s), 191
polar, 151	cycle(s), 151, 153, 154, 165, 183, 197,
yeast, 150, 154, 184, 187, 326, 347, 348	201, 202, 203, 229, 243, 264, 267,
yeasi, 150, 154, 164, 167, 320, 347, 346	201, 202, 203, 229, 243, 264, 267, 270
Ca ²⁺ , 73, 81, 88-90, 95, 106, 114, 118,	cytoplasm, 191
141-143, 166, 168, 169, 175, 180,	differentiation, 81, 189-221, 228, 244,
194, 209, 232, 237, 261, 340-342	245, 272, 352-354

division(s), 12, 15, 147, 148, 152, 160, 165, 169, 173, 185, 187, 190, 191, 196, 200, 201, 203, 205, 206, 210, 222, 232, 244-246, 252, 253, 268, 272 elongation, 148, 169, 182, 185, 186, 222, 238 growth, 147-188, 348-351 membrane(s), 333, 353 movement(s), 123, 129-146, 346, 347 polarity, 15, 112, 148, 154, 159, 160, 186, 190, 191, 205, 235, 248, 261, 272, 344, 346 polarization, 205, 207 pole(s), 203, 204 surface(s), 63, 80, 89, 112, 149, 150, 157, 166, 181, 184, 204, 218, 219, 220, 246, 249, 255, 333, 348-350, 353 target(s), 345 Cellulose, 64, 129, 147, 176, 185-187, 206, 211, 241, 244 Centriole(s), 124, 127, 138, 195 Centrosome(s), 120, 124, 126-128, 141,	Chemo-structural gradient, 197 Chemotaxis, 60, 134, 139-141, 145, 241 Chirality, 15, 19, 43-48, 50, 53, 86, 124, 268, 325-328, 334, 335, 356 Chitin, 150, 153, 154 Chitosomes, 155, 157 Chloral, 160 Chloride, 85, 341 Chloroform, 39 Chloronema(ta), 172, 209 Chlorophyll, 60, 109, 110, 211, 343 Chloroplast(s), 50, 51, 83, 103, 108-111, 129, 130, 169-172, 195, 205, 209, 229, 343, 344 Choline, 65, 92 Chromatophore, 110 Chromosome(s), 52, 55, 120, 124-126, 128, 182, 192, 201, 203, 214, 252, 272, 337, 345 Cilia, 104, 112, 114, 118, 121, 123, 130-132, 141, 145, 346
144, 145, 151, 345, 346	Circular, 50, 51, 71, 86, 124, 327
Cephalon-abdomen, 215, 257, 258	polarization, 45, 47, 48
CH ₃ Cl, 39	Cis-, 53, 81, 89, 97-100, 102, 103, 257, 260
CH ₄ , 38, 39	Clathrin 101 102
Channel(s), 22, 66, 83, 86, 87, 90, 91, 93, 94, 110, 141, 142, 154, 167, 194, 221,	Clathrin, 101, 102
340-342, 355	Clay(s), 73, 74 Cleavage(s) 45, 57, 66, 101, 158, 194,
gated (-ing), 83, 340, 341	211-213, 215, 217, 218, 229, 249,
ionic (H ⁺ , K ⁺ , Na ⁺ , Ca ²⁺ , Cl ⁻), 86-95,	252, 253, 255, 258, 259, 262-264,
154, 175, 180, 207, 216, 221, 341	266, 268, 269, 336
ligand-gated, 87	CO ₂ , 38, 65, 109, 162, 192
polar, 90	Coalescence, 242
voltage gated, 87, 88	Coat, 71, 117, 202
Chaos, 17, 328	Code, 53, 54, 57, 260, 336, 358
Charge(s), 18-20, 22-24, 28, 33, 35, 42-44,	Coenzymes (see also NAD-NADP), CoA,
51, 52, 59-62, 64, 67, 68, 71, 74, 83,	65, 103
84, 86, 87, 92, 108-110, 131,	Colchicine, 141, 160, 173, 179, 186, 206,
135-137, 168, 271, 272, 327-334,	211, 251, 344
337, 343	Coleoptile, 186, 340
asymmetry, 271, 339	Commitment, 201, 210, 249
separation, 43, 343	Communication, 72, 89, 249
transfer, 328, 334	Compartment(s), 72, 84, 99, 102, 103, 106,
Chelation, 59	168, 205, 220, 346, 353, 355
Chemical communication, 149	Competence, 198
Chemical forces, 34	Complexity, 69, 145, 197, 226, 231
Chemiosmotic theory, 82-85, 186, 102, 109,	Computer(s), 23, 26, 61, 66, 146, 330, 337
235	Conductivity-conductance, 22-25, 86, 107,

220 222 220 241	210 212 214 217 210 227 227
329, 333, 338, 341	210, 212-214, 217, 218, 225, 235,
Conformation, 45, 49, 50, 70, 85-87, 89,	236, 255, 257, 259, 264, 269, 271,
117, 121, 344	272, 350
Conidia, 107, 155, 156, 184, 187, 188, 192,	Cytoplasmic, 15, 79-81, 88, 89, 94, 95, 100,
193, 196, 349	106, 111, 115, 123, 124, 126,
Copper $(Cu^{+/2+})$, 23, 24, 61, 65, 66, 335	129-131, 133, 139, 141, 146, 149,
Cortex, 100, 134, 136, 138, 144, 145, 202,	151, 152, 158, 161, 168, 169, 173,
210, 212, 216, 244, 263, 344, 346	179, 185, 189, 194, 197, 198, 200,
Cosmic, 17, 25, 325, 327, 328, 330	206, 209-214, 217, 221, 228, 229,
Coupling, 20, 24-26, 54, 82, 86, 89, 94, 97,	235, 252, 255, 261, 262, 264, 340,
102, 111, 148, 185, 223, 250, 326,	349, 350
341	basophilic gradient, 193
Crystal(s), 18, 21-24, 31, 32, 35, 37, 40, 55,	DNA, 198
63, 64, 68, 76, 86, 110, 271, 328-333	granules, 151
Current(s), 22, 23, 26, 34, 45, 62, 92, 95,	movements, 73, 129, 261
120, 129, 137, 139, 142, 143, 154,	streaming, 15, 129, 130, 137, 162, 164,
159, 161, 166-169, 194, 207, 216,	175
230, 235, 236, 262, 339, 341, 349,	zonation, 169
353	Cytoskeleton, 81, 111, 112, 124, 132, 138,
loop(s), 157, 168, 231, 272	139, 144, 145, 158, 162, 166, 173,
Cyanide, 132	174, 176, 177, 179, 180, 200, 204,
Cycle(s), 90, 102, 113, 133, 135, 136, 147,	206, 213, 214, 218, 219, 254, 264,
154, 165, 201, 203, 330	271, 325, 343-350, 353, 355, 358
Cyclic, 139, 179, 343	outgrowths, 261
adenosine monophosphate, 253	protein(s), 73, 77, 95, 99, 177, 350, 356
cAMP, 134, 140, 141, 157, 162, 163,	Cytosol, 65, 72, 73, 77, 102, 111, 118, 130,
240, 243, 340	172, 177, 179, 343
guanosine 5'-phosphate, 89	
photophosphorylation, 192	Dehydrogenases, 82, 84
Cyclosis, 129, 162, 164	Deoxyribonucleic acid (DNA), 45, 49-57,
Cylindrical, 69, 71, 78, 130, 137, 147,	61, 64, 70, 71, 81, 155, 189-191,
154-163, 165, 182, 183, 186, 221, 328	197-200, 203, 259, 260, 272, 336,
germ tubes, 154-161	337, 348, 358
Cysteine-Cystine, 40, 41, 337	packaging, 71, 337
Cytochalasin(s) B, E, H, 80, 113, 145, 179,	polarity, 49, 336
200, 206, 207, 229, 344, 350, 352	polymerase, 61
Cytochemical gradients, 166	positioning, 197-200
Cytochrome(s), 61, 66, 84, 198, 338, 339	replication, 52, 190, 203
oxidase, 198, 339	segregation, 191
Cytogel, 72	topology, 53
Cytokeratin, 111	
	transcription, 52-54
Cytokinesis, 136, 151, 153, 154, 161, 184,	Depolarization, 87-92, 94, 95, 108, 141,
200, 208 Cutalutia 00	142, 157, 216, 221, 251, 334, 340,
Cytolytic, 99	341, 346
Cytoplasm, 15, 54, 62, 72, 85, 89, 103, 104,	Design, 42, 69, 74
112, 122, 124, 129, 130, 136, 142,	Desmin, 111
158, 161, 164, 168, 172, 173, 176,	Detergent(s), 63, 76, 78, 110, 115
186, 189, 190, 194, 198, 199, 208,	Determinants, 122, 215, 253

Development, 15, 27, 72, 74, 91, 147, 152,	332, 344, 350, 352, 358
155, 157, 160, 164, 165, 167,	division, 148, 185, 186, 200
169-172, 176, 189, 193, 194, 203,	light, 209
204, 206, 208, 210, 212, 213, 217,	nucleation, 244
221, 222, 224-230, 232, 238, 240,	Dismutation, 66
243, 244, 247, 251-255, 260, 261,	Dissipation, 195, 197
263, 264, 270, 272, 338, 356	Dissymmetry, 33, 38
Dielectric, 19, 34, 61, 67, 332, 338, 339	Division(s) 79, 104, 120, 148, 149, 151, 160,
constant, 78, 338	161, 172-174, 176, 181-183, 186,
polarization, 332	189, 191, 192, 197, 200, 202, 203,
Dielectrophoresis, 78	205, 206, 209-213, 217, 222, 224,
Differentiation, 13, 15, 139, 181, 189-221,	225, 229, 232, 245, 252, 253, 267,
223, 225, 228, 230, 231, 233, 238,	268, 272, 345, 352
240, 243, 244, 247, 255, 256, 260,	nuclear, 125, 152, 205
264, 269, 352-354	unequal, 182, 197, 200, 205, 209, 212,
apical, 192-200	272
apico-basal, 200-221	Domain(s), 47, 58, 66, 67, 76, 79, 81, 103,
intercalary, 190-192	213, 218, 221, 254, 260, 329, 349,
Diffusion, 14, 25, 61, 76, 77, 101, 104, 154,	353-356
166, 235, 240, 272, 273, 333, 338,	Dominance, 13, 47, 144, 242
354	Dorsal, 263
gradient(s), 14, 224, 228	meristem, 231
Dimorphism, 117, 157, 161-163, 201, 203,	tissue(s), 250, 257
238, 348	products, 260
Diode, 23, 330	Dorso-ventral (D/V)
Diploid, 184, 196, 197	axis, 256, 262
Dipole(s), 13, 23, 26, 30, 33-39, 41-43, 58,	genes, 259
59, 61-63, 68, 75, 76, 79, 86, 87, 108,	gradients, 264
111, 249, 250, 271, 273, 325-334,	pattern(s), 256, 260
336, 340, 343, 358	polarities, 214, 250-266, 267-270
electric, 21, 22, 33, 34, 328, 329	structure, 173, 209
field, 25, 332	Dots, 152-154, 205
interactions, 76	Double
mineral, 34-38, 332, 333	gradient(s), 190, 255
moment(s), 25, 33, 37, 39, 58, 60, 63,	helix, 50, 51, 56, 57, 112, 336
76, 85-87, 89, 111, 329, 330, 332, 338	strands, 62, 86
organic, 38-43, 334	Driving force(s), 42, 43, 111, 134, 135, 147,
protein, 61, 337	166, 167, 186, 230, 345, 346
water, 34-38, 332, 333	Drug(s) 70, 80, 101, 121, 144, 145, 152, 188,
Diprotic, 41	267, 335, 343, 349
Direction (-ality), 17, 19, 23, 25, 26, 28-33,	Duality, 13, 54, 67, 108, 140, 142, 161, 166
37, 43, 45, 49-53, 57, 58, 61, 62, 82,	Duplex, 52, 53
86, 93, 99, 104, 106, 115-119,	Dyad, 51
121-123, 126, 129, 130, 133, 136,	Dynamics, 46, 111, 121, 126, 127
140-147, 159, 160, 164, 165, 173,	Dynamin, 345
174, 176, 179, 182, 185, 186, 189,	Dynein, 121, 123, 126, 129, 179, 344
192, 197, 199, 205, 207-210, 220,	
221, 224, 228, 231-234, 236, 241,	Ecto-meso-endoderms, 213, 248, 262, 264,
245, 249, 252, 254, 255, 261-263,	266, 269, 270

Ectoplasm, 129, 133, 134, 173, 215, 261 Electrogenic, 110, 235 Egg(s), 13, 14, 157, 158, 190, 201, 204, 206, pump(s), 154, 168, 339-341 208, 210-218, 225, 241, 252-265, Electromechanical, 168 268-270, 325, 349, 352, 355 Electromagnetic, 17-20, 26, 30, 31, 42, 45, axial polarity, 213, 217, 253, 254, 355 325, 327, 329 crescents field, 21, 31 grey, 255, 262, 263 Electron(s), 13, 17-24-31, 33, 35, 38-44, 47, yellow-orange-red, 216, 254, 261, 50, 51, 60, 62, 65, 66, 68, 78, 82, 84, 90, 106, 108-111, 143, 144, 186, 271, 262 272, 325, 332, 334, 339-341, 343 fertilized, 158, 159, 204, 210, 213, 215, 252, 253, 263 microscopy, 75, 98, 111, 114, 118, 119, Electric(al), 13, 14, 21-23, 26, 33-38, 60, 70, 124, 151, 155, 156, 183, 184 74, 84, 86, 87, 89, 90-93, 107, 123, polarization, 29, 47 transfer reactions, 338, 339, 343 131, 132, 143, 166, 168, 174, 209, transport, 82, 84, 110, 142, 186, 334 221, 248, 271, 272, 325, 338-342 biopolarity, 26, 107, 158, 272 tunnelling, 42 bipolarity, 13, 15, 23, 108, 136, 230, Electronic, 23, 30, 38, 43, 61, 68, 110, 330, 239, 271, 354 331 Electrophoresis, 14, 50, 106, 118, 172, 200, bipolarization, 15, 19, 22, 328 charge(s), 13, 14, 17-22, 27, 29, 30, 34, 35, 40, 41, 61, 62, 70, 85, 87, 93, 109, Electrostatic, 14, 35, 58-61, 66, 67, 75, 336 221, 271, 325, 328, 329, 343 Elongation, 152, 165, 180-186 current(s), 26, 34, 62, 73, 74, 78, 79, Embryo(s), 72, 126, 180, 189, 204, 210-215, 158, 159, 168, 204, 207, 209, 262, 217, 218, 223-225, 229, 230, 240, 325, 327, 339, 349 249-259, 261, 263-268, 270, 352-355 depolarization, 89 induction, 217 patterns, 256-260 dipole(s), 19, 21, 23, 27, 33-36, 63, 76, 87, 137, 272, 325, 328, 330, 339 polarity, 224, 257, 264 field(s), 19, 22-24, 31-34, 36, 37, 41, 61, Embryogenesis, 13, 210, 213, 216, 218, 224, 62, 78, 85, 87, 110, 137, 145, 160, 238, 241, 252, 255, 259, 262, 266, 167, 168, 176, 180, 188, 201, 209, 325, 352 210, 239, 247, 332, 338-340, 355 Enantiomer(s), 44-47, 325, 326, 328, 334, gradient, 102, 204 335 moment(s), 33, 38, 39, 272, 329, 331, Enantiomorphs, 32 332 Endocrine, 93, 220 poles, 27, 175, 325, 333 Endocytosis, 99, 101, 102, 180, 219, 342 potential(s), 83-86, 89, 90, 106-108, Endonuclease, 51 Endoplasm, 129, 133, 134, 215, 261 133, 142, 143, 159, 166, 170, 200, 235, 338-340 Endoplasmic reticulum (ER), 59, 65, 74, 88, 95-100, 103, 148, 155, 164, 166, signal(s), 87, 93, 94, 221 180, 236, 342, 343, 354 stimulation, 89, 154, 341 Energy, 17, 19, 21-24, 27, 30, 31, 35, 37, Electride, 30 Electrochemical, 68, 162, 176, 339 43-45, 51, 52, 58, 64, 67, 68, 74-76, 80-82, 84, 85, 87, 91, 92, 95, 100, force, 175 102, 104-111, 116, 119, 132, 140, gradient(s), 82, 84, 91, 102, 107, 109, 142, 167, 178, 198, 209, 235, 271, 140-142, 167, 168, 235, 339-341 potential, 82, 84, 102, 107, 110 328, 333, 334, 338-341, 343 transduction, 74, 82, 89, 102, 109, 338 Electrode(s), 92, 137, 167, 339 Entropy, 18, 19 Electrofusion, 78

Environment, 14, 30, 40, 43, 46, 47, 50, 51, 61, 66, 95, 101, 134, 142, 159, 181, 224, 233, 239, 352 Environmental, 138, 155, 163, 187, 197, 209, 226, 239, 252 factors, 129, 172, 272, 356 polarity, 79, 216 Enzyme(s), 46, 50-54, 56, 60, 64-66, 69, 73, 79-82, 85, 89, 93, 98-100, 103, 118, 123, 164, 166, 177, 186, 187, 194, 208, 220, 259, 337, 339, 340 Epidermis, 145, 175, 176, 185, 186, 210, 211, 228, 230, 261 Epigenetic(s), 201, 223, 224, 251, 358 Epithelium (epithelia), 104, 115, 146, 190, 218-221, 247, 260, 353 Equilibrium, 37, 102, 114, 119, 120, 159, 162, 222 Equipolar, 147, 181 Erythrocyte(s), 52, 62, 79, 137 Ethanol, 39, 43 Evolution, 46, 94, 192, 269, 271 Excitation, 42, 43, 51, 61, 74, 86, 88, 93, 108-111 Excited electrons, 23 ionization, 271, 327, 329 Excretion, 130, 230, 235 Exocytosis, 90, 94, 98-102, 135, 219, 220 Fascin, 114 Fat, 104 Fatty acids, 39, 40, 62, 73, 96 Feedback, 92, 168, 173, 250 Female, 174, 196-200, 327 basophilic gradient, 198, 199 gametangia, 192, 198, 199 Fermentation, 215	Fibroblast(s), 62, 135, 137, 146, 342, 346 Fibronectin, 78, 134, 135 Field(s), 19, 22-27, 30-33, 142, 165, 175, 180, 189, 210, 212, 222, 223, 240, 245-247, 252, 272, 330, 331, 355 polar, 205, 331 Filamin, 114 Filopodia, 122, 179, 180, 350 Fimbrin, 114 Flagella, 118, 121, 130-132, 138-140, 143, 195, 346 polar, 201 Flip-flop(s), 56, 332 Flippase, 77 Flowering, 161, 197, 238, 239 Fluid(s), 37, 72, 77, 93, 101, 136 mosaic model, 75, 77, 80 Fluorescence, 42, 61, 155, 175, 183, 188, 347 polarity, 340 Fluorophores, 61, 108, 350 Flux(es), 25, 143, 207 Fodrin, 95, 221 Folic acid, 141 Foot, 135, 225, 247, 250, 340, 358 Formaldehyde, 46, 67 Formic acid, 39 Freeze, 102, 269 Fungal, 63, 105, 155-157, 162, 164-168, 181, 184, 187, 192-195, 228, 341, 343, 345 conidia, 155, 181, 349 cell(s), 161, 164, 204 exosporulation, 192, 352 germ tubes, 155-157, 184, 187, 188 hypha(e), 107, 147, 165, 166, 169, 176, 186, 349
Fern(s) gametophyte, 147	186, 349 mycelia, 164 spore(s), 155, 160, 163, 325, 348, 352
leaf, 238 mother cell, 210 prothallia, 173, 174 spores, 160, 173 Ferritin, 79, 135 Ferroelectricity, 26, 330, 331 Ferromagnetism, 26 Fertilization, 89, 157-159, 201, 204-207, 210, 212, 213, 215, 216, 224, 225, 252-254, 259, 261-263, 268, 269	G-protein(s), 81, 89, 113, 149 Galvanotropism, 180 Gametangia, 197-199 Gamete(s), 170, 195, 198, 215, 252 Gametophyte(s), 160, 161, 173, 224 Gamma (γ)-rays, 19, 21, 31, 47 Gap junction(s), 219, 249 Gas, 17, 18, 25, 28, 29, 327, 339 Gel, 78, 114

Gene(s), 15, 52-58, 70, 74, 148, 162, 163,	ionic (H ⁺ , Ca ²⁺ ,Cl ⁻), 162, 163, 341,
170, 189, 192, 196-198, 200, 204,	349
213, 223, 229, 243, 244, 246,	light, 209
255-260, 268, 269, 272, 336, 356, 358	metabolic, 208, 217, 255
conversion, 56	morphogens, 15, 222, 223, 230, 240,
expression, 15, 52, 53, 55, 56, 163, 181,	247, 354, 358
189, 238	polarity, 205, 340
polarity, 52, 199, 259, 272, 356, 358	proton(s), 82, 83, 102, 108, 237, 338
targeting, 118	protoplasmic, 205
Genetic(s)	redox, 264
analysis, 55, 259	RNA, 198, 199
control, 208, 358	spatial, 180
information, 57, 74, 132	Grafting, 14, 208, 212, 225, 233, 242,
Genome, 50, 51, 57, 189, 191, 211, 223,	245-247, 249, 265
260, 336	Gramicidin, 86, 133
Germ tube(s), 108, 155-157, 160, 161, 163,	Gravitation, 26, 216
171, 181, 184, 187, 188, 204, 206,	field, 216, 234, 354
325, 348	forces, 19
Germination(s), 155-157, 160, 163, 187,	Gravitropism (Geotropism), 237, 341, 354,
196, 204, 205, 209, 348, 349, 354	355
Gliding, 132, 133, 139, 142, 143, 346	curvature, 168, 352
Globin, 52	response, 168, 186, 236
Globular, 76, 115, 117	Gravity, 19, 33, 39, 172, 173, 209, 216, 234,
molecules, 60, 76, 77, 106, 111, 113, 119	237, 238, 264, 265, 327, 341, 354,
Glucans, 166, 184	355
Clusoco 64 167 169 193	norgantian concer 254 255
Glucose, 64, 167, 168, 183	perception-sensor, 354, 355
Glutamic acid, 62	signal-stimulus, 236, 354
Glutamic acid, 62 Glycine, 39, 40, 41	signal-stimulus, 236, 354 Growth, 13, 52, 68, 81, 113, 114, 117, 119,
Glutamic acid, 62 Glycine, 39, 40, 41 Glycocalyx, 78, 81, 129	signal-stimulus, 236, 354 Growth, 13, 52, 68, 81, 113, 114, 117, 119, 122, 129, 131, 134, 146-150, 152,
Glutamic acid, 62 Glycine, 39, 40, 41 Glycocalyx, 78, 81, 129 Glycolipids, 99, 353	signal-stimulus, 236, 354 Growth, 13, 52, 68, 81, 113, 114, 117, 119, 122, 129, 131, 134, 146-150, 152, 154, 155, 158-160, 162-166, 169-174,
Glutamic acid, 62 Glycine, 39, 40, 41 Glycocalyx, 78, 81, 129 Glycolipids, 99, 353 Glycolysis, 73, 92, 215	signal-stimulus, 236, 354 Growth, 13, 52, 68, 81, 113, 114, 117, 119, 122, 129, 131, 134, 146-150, 152, 154, 155, 158-160, 162-166, 169-174, 176, 177, 179-188, 191-194, 201,
Glutamic acid, 62 Glycine, 39, 40, 41 Glycocalyx, 78, 81, 129 Glycolipids, 99, 353 Glycolysis, 73, 92, 215 Glycoproteins, 78, 80, 81, 98-100, 177, 347	signal-stimulus, 236, 354 Growth, 13, 52, 68, 81, 113, 114, 117, 119, 122, 129, 131, 134, 146-150, 152, 154, 155, 158-160, 162-166, 169-174, 176, 177, 179-188, 191-194, 201, 206, 208-210, 212, 224-226, 228,
Glutamic acid, 62 Glycine, 39, 40, 41 Glycocalyx, 78, 81, 129 Glycolipids, 99, 353 Glycolysis, 73, 92, 215 Glycoproteins, 78, 80, 81, 98-100, 177, 347 Glyoxysomes, 103	signal-stimulus, 236, 354 Growth, 13, 52, 68, 81, 113, 114, 117, 119, 122, 129, 131, 134, 146-150, 152, 154, 155, 158-160, 162-166, 169-174, 176, 177, 179-188, 191-194, 201, 206, 208-210, 212, 224-226, 228, 230-234, 236, 237, 241, 271, 344,
Glutamic acid, 62 Glycine, 39, 40, 41 Glycocalyx, 78, 81, 129 Glycolipids, 99, 353 Glycolysis, 73, 92, 215 Glycoproteins, 78, 80, 81, 98-100, 177, 347 Glyoxysomes, 103 Golgi (apparatus), 73, 74, 79, 95-104, 121,	signal-stimulus, 236, 354 Growth, 13, 52, 68, 81, 113, 114, 117, 119, 122, 129, 131, 134, 146-150, 152, 154, 155, 158-160, 162-166, 169-174, 176, 177, 179-188, 191-194, 201, 206, 208-210, 212, 224-226, 228, 230-234, 236, 237, 241, 271, 344, 345, 348-351, 353, 357
Glutamic acid, 62 Glycine, 39, 40, 41 Glycocalyx, 78, 81, 129 Glycolipids, 99, 353 Glycolysis, 73, 92, 215 Glycoproteins, 78, 80, 81, 98-100, 177, 347 Glyoxysomes, 103 Golgi (apparatus), 73, 74, 79, 95-104, 121, 148, 166, 175, 178, 206, 207, 216,	signal-stimulus, 236, 354 Growth, 13, 52, 68, 81, 113, 114, 117, 119, 122, 129, 131, 134, 146-150, 152, 154, 155, 158-160, 162-166, 169-174, 176, 177, 179-188, 191-194, 201, 206, 208-210, 212, 224-226, 228, 230-234, 236, 237, 241, 271, 344, 345, 348-351, 353, 357 axis, 147, 160, 173, 183
Glutamic acid, 62 Glycine, 39, 40, 41 Glycocalyx, 78, 81, 129 Glycolipids, 99, 353 Glycolysis, 73, 92, 215 Glycoproteins, 78, 80, 81, 98-100, 177, 347 Glyoxysomes, 103 Golgi (apparatus), 73, 74, 79, 95-104, 121, 148, 166, 175, 178, 206, 207, 216, 342, 343	signal-stimulus, 236, 354 Growth, 13, 52, 68, 81, 113, 114, 117, 119, 122, 129, 131, 134, 146-150, 152, 154, 155, 158-160, 162-166, 169-174, 176, 177, 179-188, 191-194, 201, 206, 208-210, 212, 224-226, 228, 230-234, 236, 237, 241, 271, 344, 345, 348-351, 353, 357 axis, 147, 160, 173, 183 cone(s), 176, 177, 179, 180, 342, 349,
Glutamic acid, 62 Glycine, 39, 40, 41 Glycocalyx, 78, 81, 129 Glycolipids, 99, 353 Glycolysis, 73, 92, 215 Glycoproteins, 78, 80, 81, 98-100, 177, 347 Glyoxysomes, 103 Golgi (apparatus), 73, 74, 79, 95-104, 121, 148, 166, 175, 178, 206, 207, 216, 342, 343 vesicles, 97, 99-101, 164, 168, 178	signal-stimulus, 236, 354 Growth, 13, 52, 68, 81, 113, 114, 117, 119, 122, 129, 131, 134, 146-150, 152, 154, 155, 158-160, 162-166, 169-174, 176, 177, 179-188, 191-194, 201, 206, 208-210, 212, 224-226, 228, 230-234, 236, 237, 241, 271, 344, 345, 348-351, 353, 357 axis, 147, 160, 173, 183 cone(s), 176, 177, 179, 180, 342, 349, 350
Glutamic acid, 62 Glycine, 39, 40, 41 Glycocalyx, 78, 81, 129 Glycolipids, 99, 353 Glycolysis, 73, 92, 215 Glycoproteins, 78, 80, 81, 98-100, 177, 347 Glyoxysomes, 103 Golgi (apparatus), 73, 74, 79, 95-104, 121, 148, 166, 175, 178, 206, 207, 216, 342, 343 vesicles, 97, 99-101, 164, 168, 178 Gradient(s), 14, 23, 54, 56, 72, 73, 82, 87,	signal-stimulus, 236, 354 Growth, 13, 52, 68, 81, 113, 114, 117, 119, 122, 129, 131, 134, 146-150, 152, 154, 155, 158-160, 162-166, 169-174, 176, 177, 179-188, 191-194, 201, 206, 208-210, 212, 224-226, 228, 230-234, 236, 237, 241, 271, 344, 345, 348-351, 353, 357 axis, 147, 160, 173, 183 cone(s), 176, 177, 179, 180, 342, 349, 350 differential, 186, 236, 237, 271
Glutamic acid, 62 Glycine, 39, 40, 41 Glycocalyx, 78, 81, 129 Glycolipids, 99, 353 Glycolysis, 73, 92, 215 Glycoproteins, 78, 80, 81, 98-100, 177, 347 Glyoxysomes, 103 Golgi (apparatus), 73, 74, 79, 95-104, 121, 148, 166, 175, 178, 206, 207, 216, 342, 343 vesicles, 97, 99-101, 164, 168, 178 Gradient(s), 14, 23, 54, 56, 72, 73, 82, 87, 93, 104, 133, 139-142, 146, 160, 162,	signal-stimulus, 236, 354 Growth, 13, 52, 68, 81, 113, 114, 117, 119, 122, 129, 131, 134, 146-150, 152, 154, 155, 158-160, 162-166, 169-174, 176, 177, 179-188, 191-194, 201, 206, 208-210, 212, 224-226, 228, 230-234, 236, 237, 241, 271, 344, 345, 348-351, 353, 357 axis, 147, 160, 173, 183 cone(s), 176, 177, 179, 180, 342, 349, 350 differential, 186, 236, 237, 271 direction, 147, 148, 186
Glutamic acid, 62 Glycine, 39, 40, 41 Glycocalyx, 78, 81, 129 Glycolipids, 99, 353 Glycolysis, 73, 92, 215 Glycoproteins, 78, 80, 81, 98-100, 177, 347 Glyoxysomes, 103 Golgi (apparatus), 73, 74, 79, 95-104, 121, 148, 166, 175, 178, 206, 207, 216, 342, 343 vesicles, 97, 99-101, 164, 168, 178 Gradient(s), 14, 23, 54, 56, 72, 73, 82, 87, 93, 104, 133, 139-142, 146, 160, 162, 163, 165, 170-172, 176, 180, 190,	signal-stimulus, 236, 354 Growth, 13, 52, 68, 81, 113, 114, 117, 119, 122, 129, 131, 134, 146-150, 152, 154, 155, 158-160, 162-166, 169-174, 176, 177, 179-188, 191-194, 201, 206, 208-210, 212, 224-226, 228, 230-234, 236, 237, 241, 271, 344, 345, 348-351, 353, 357 axis, 147, 160, 173, 183 cone(s), 176, 177, 179, 180, 342, 349, 350 differential, 186, 236, 237, 271 direction, 147, 148, 186 elongation, 182-186, 188, 349
Glutamic acid, 62 Glycine, 39, 40, 41 Glycocalyx, 78, 81, 129 Glycolipids, 99, 353 Glycolysis, 73, 92, 215 Glycoproteins, 78, 80, 81, 98-100, 177, 347 Glyoxysomes, 103 Golgi (apparatus), 73, 74, 79, 95-104, 121, 148, 166, 175, 178, 206, 207, 216, 342, 343 vesicles, 97, 99-101, 164, 168, 178 Gradient(s), 14, 23, 54, 56, 72, 73, 82, 87, 93, 104, 133, 139-142, 146, 160, 162, 163, 165, 170-172, 176, 180, 190, 195, 197-201, 205-207, 211, 215,	signal-stimulus, 236, 354 Growth, 13, 52, 68, 81, 113, 114, 117, 119, 122, 129, 131, 134, 146-150, 152, 154, 155, 158-160, 162-166, 169-174, 176, 177, 179-188, 191-194, 201, 206, 208-210, 212, 224-226, 228, 230-234, 236, 237, 241, 271, 344, 345, 348-351, 353, 357 axis, 147, 160, 173, 183 cone(s), 176, 177, 179, 180, 342, 349, 350 differential, 186, 236, 237, 271 direction, 147, 148, 186 elongation, 182-186, 188, 349 factor(s), 148, 149, 181, 226, 232, 354,
Glutamic acid, 62 Glycine, 39, 40, 41 Glycocalyx, 78, 81, 129 Glycolipids, 99, 353 Glycolysis, 73, 92, 215 Glycoproteins, 78, 80, 81, 98-100, 177, 347 Glyoxysomes, 103 Golgi (apparatus), 73, 74, 79, 95-104, 121, 148, 166, 175, 178, 206, 207, 216, 342, 343 vesicles, 97, 99-101, 164, 168, 178 Gradient(s), 14, 23, 54, 56, 72, 73, 82, 87, 93, 104, 133, 139-142, 146, 160, 162, 163, 165, 170-172, 176, 180, 190, 195, 197-201, 205-207, 211, 215, 217, 222, 223, 230-233, 237, 239,	signal-stimulus, 236, 354 Growth, 13, 52, 68, 81, 113, 114, 117, 119, 122, 129, 131, 134, 146-150, 152, 154, 155, 158-160, 162-166, 169-174, 176, 177, 179-188, 191-194, 201, 206, 208-210, 212, 224-226, 228, 230-234, 236, 237, 241, 271, 344, 345, 348-351, 353, 357 axis, 147, 160, 173, 183 cone(s), 176, 177, 179, 180, 342, 349, 350 differential, 186, 236, 237, 271 direction, 147, 148, 186 elongation, 182-186, 188, 349 factor(s), 148, 149, 181, 226, 232, 354, 355
Glutamic acid, 62 Glycine, 39, 40, 41 Glycocalyx, 78, 81, 129 Glycolipids, 99, 353 Glycolysis, 73, 92, 215 Glycoproteins, 78, 80, 81, 98-100, 177, 347 Glyoxysomes, 103 Golgi (apparatus), 73, 74, 79, 95-104, 121, 148, 166, 175, 178, 206, 207, 216, 342, 343 vesicles, 97, 99-101, 164, 168, 178 Gradient(s), 14, 23, 54, 56, 72, 73, 82, 87, 93, 104, 133, 139-142, 146, 160, 162, 163, 165, 170-172, 176, 180, 190, 195, 197-201, 205-207, 211, 215, 217, 222, 223, 230-233, 237, 239, 240, 247-250, 253, 255, 256, 261,	signal-stimulus, 236, 354 Growth, 13, 52, 68, 81, 113, 114, 117, 119, 122, 129, 131, 134, 146-150, 152, 154, 155, 158-160, 162-166, 169-174, 176, 177, 179-188, 191-194, 201, 206, 208-210, 212, 224-226, 228, 230-234, 236, 237, 241, 271, 344, 345, 348-351, 353, 357 axis, 147, 160, 173, 183 cone(s), 176, 177, 179, 180, 342, 349, 350 differential, 186, 236, 237, 271 direction, 147, 148, 186 elongation, 182-186, 188, 349 factor(s), 148, 149, 181, 226, 232, 354, 355 inhibitors, 236, 351
Glutamic acid, 62 Glycine, 39, 40, 41 Glycocalyx, 78, 81, 129 Glycolipids, 99, 353 Glycolysis, 73, 92, 215 Glycoproteins, 78, 80, 81, 98-100, 177, 347 Glyoxysomes, 103 Golgi (apparatus), 73, 74, 79, 95-104, 121, 148, 166, 175, 178, 206, 207, 216, 342, 343 vesicles, 97, 99-101, 164, 168, 178 Gradient(s), 14, 23, 54, 56, 72, 73, 82, 87, 93, 104, 133, 139-142, 146, 160, 162, 163, 165, 170-172, 176, 180, 190, 195, 197-201, 205-207, 211, 215, 217, 222, 223, 230-233, 237, 239, 240, 247-250, 253, 255, 256, 261, 262, 264, 272, 338-341, 349, 354, 358	signal-stimulus, 236, 354 Growth, 13, 52, 68, 81, 113, 114, 117, 119, 122, 129, 131, 134, 146-150, 152, 154, 155, 158-160, 162-166, 169-174, 176, 177, 179-188, 191-194, 201, 206, 208-210, 212, 224-226, 228, 230-234, 236, 237, 241, 271, 344, 345, 348-351, 353, 357 axis, 147, 160, 173, 183 cone(s), 176, 177, 179, 180, 342, 349, 350 differential, 186, 236, 237, 271 direction, 147, 148, 186 elongation, 182-186, 188, 349 factor(s), 148, 149, 181, 226, 232, 354, 355 inhibitors, 236, 351 orientation, 148, 185
Glutamic acid, 62 Glycine, 39, 40, 41 Glycocalyx, 78, 81, 129 Glycolipids, 99, 353 Glycolysis, 73, 92, 215 Glycoproteins, 78, 80, 81, 98-100, 177, 347 Glyoxysomes, 103 Golgi (apparatus), 73, 74, 79, 95-104, 121, 148, 166, 175, 178, 206, 207, 216, 342, 343 vesicles, 97, 99-101, 164, 168, 178 Gradient(s), 14, 23, 54, 56, 72, 73, 82, 87, 93, 104, 133, 139-142, 146, 160, 162, 163, 165, 170-172, 176, 180, 190, 195, 197-201, 205-207, 211, 215, 217, 222, 223, 230-233, 237, 239, 240, 247-250, 253, 255, 256, 261, 262, 264, 272, 338-341, 349, 354, 358 bioelectric, 354	signal-stimulus, 236, 354 Growth, 13, 52, 68, 81, 113, 114, 117, 119, 122, 129, 131, 134, 146-150, 152, 154, 155, 158-160, 162-166, 169-174, 176, 177, 179-188, 191-194, 201, 206, 208-210, 212, 224-226, 228, 230-234, 236, 237, 241, 271, 344, 345, 348-351, 353, 357 axis, 147, 160, 173, 183 cone(s), 176, 177, 179, 180, 342, 349, 350 differential, 186, 236, 237, 271 direction, 147, 148, 186 elongation, 182-186, 188, 349 factor(s), 148, 149, 181, 226, 232, 354, 355 inhibitors, 236, 351 orientation, 148, 185 pattern, 160, 175, 348, 349, 357
Glutamic acid, 62 Glycine, 39, 40, 41 Glycocalyx, 78, 81, 129 Glycolipids, 99, 353 Glycolysis, 73, 92, 215 Glycoproteins, 78, 80, 81, 98-100, 177, 347 Glyoxysomes, 103 Golgi (apparatus), 73, 74, 79, 95-104, 121, 148, 166, 175, 178, 206, 207, 216, 342, 343 vesicles, 97, 99-101, 164, 168, 178 Gradient(s), 14, 23, 54, 56, 72, 73, 82, 87, 93, 104, 133, 139-142, 146, 160, 162, 163, 165, 170-172, 176, 180, 190, 195, 197-201, 205-207, 211, 215, 217, 222, 223, 230-233, 237, 239, 240, 247-250, 253, 255, 256, 261, 262, 264, 272, 338-341, 349, 354, 358 bioelectric, 354 differentiation, 197-200, 217	signal-stimulus, 236, 354 Growth, 13, 52, 68, 81, 113, 114, 117, 119, 122, 129, 131, 134, 146-150, 152, 154, 155, 158-160, 162-166, 169-174, 176, 177, 179-188, 191-194, 201, 206, 208-210, 212, 224-226, 228, 230-234, 236, 237, 241, 271, 344, 345, 348-351, 353, 357 axis, 147, 160, 173, 183 cone(s), 176, 177, 179, 180, 342, 349, 350 differential, 186, 236, 237, 271 direction, 147, 148, 186 elongation, 182-186, 188, 349 factor(s), 148, 149, 181, 226, 232, 354, 355 inhibitors, 236, 351 orientation, 148, 185 pattern, 160, 175, 348, 349, 357 polar, 133, 147, 181, 348-351
Glutamic acid, 62 Glycine, 39, 40, 41 Glycocalyx, 78, 81, 129 Glycolipids, 99, 353 Glycolysis, 73, 92, 215 Glycoproteins, 78, 80, 81, 98-100, 177, 347 Glyoxysomes, 103 Golgi (apparatus), 73, 74, 79, 95-104, 121, 148, 166, 175, 178, 206, 207, 216, 342, 343 vesicles, 97, 99-101, 164, 168, 178 Gradient(s), 14, 23, 54, 56, 72, 73, 82, 87, 93, 104, 133, 139-142, 146, 160, 162, 163, 165, 170-172, 176, 180, 190, 195, 197-201, 205-207, 211, 215, 217, 222, 223, 230-233, 237, 239, 240, 247-250, 253, 255, 256, 261, 262, 264, 272, 338-341, 349, 354, 358 bioelectric, 354 differentiation, 197-200, 217 dissipation, 85	signal-stimulus, 236, 354 Growth, 13, 52, 68, 81, 113, 114, 117, 119, 122, 129, 131, 134, 146-150, 152, 154, 155, 158-160, 162-166, 169-174, 176, 177, 179-188, 191-194, 201, 206, 208-210, 212, 224-226, 228, 230-234, 236, 237, 241, 271, 344, 345, 348-351, 353, 357 axis, 147, 160, 173, 183 cone(s), 176, 177, 179, 180, 342, 349, 350 differential, 186, 236, 237, 271 direction, 147, 148, 186 elongation, 182-186, 188, 349 factor(s), 148, 149, 181, 226, 232, 354, 355 inhibitors, 236, 351 orientation, 148, 185 pattern, 160, 175, 348, 349, 357 polar, 133, 147, 181, 348-351 zone, 165, 169, 181
Glutamic acid, 62 Glycine, 39, 40, 41 Glycocalyx, 78, 81, 129 Glycolipids, 99, 353 Glycolysis, 73, 92, 215 Glycoproteins, 78, 80, 81, 98-100, 177, 347 Glyoxysomes, 103 Golgi (apparatus), 73, 74, 79, 95-104, 121, 148, 166, 175, 178, 206, 207, 216, 342, 343 vesicles, 97, 99-101, 164, 168, 178 Gradient(s), 14, 23, 54, 56, 72, 73, 82, 87, 93, 104, 133, 139-142, 146, 160, 162, 163, 165, 170-172, 176, 180, 190, 195, 197-201, 205-207, 211, 215, 217, 222, 223, 230-233, 237, 239, 240, 247-250, 253, 255, 256, 261, 262, 264, 272, 338-341, 349, 354, 358 bioelectric, 354 differentiation, 197-200, 217	signal-stimulus, 236, 354 Growth, 13, 52, 68, 81, 113, 114, 117, 119, 122, 129, 131, 134, 146-150, 152, 154, 155, 158-160, 162-166, 169-174, 176, 177, 179-188, 191-194, 201, 206, 208-210, 212, 224-226, 228, 230-234, 236, 237, 241, 271, 344, 345, 348-351, 353, 357 axis, 147, 160, 173, 183 cone(s), 176, 177, 179, 180, 342, 349, 350 differential, 186, 236, 237, 271 direction, 147, 148, 186 elongation, 182-186, 188, 349 factor(s), 148, 149, 181, 226, 232, 354, 355 inhibitors, 236, 351 orientation, 148, 185 pattern, 160, 175, 348, 349, 357 polar, 133, 147, 181, 348-351

H, see Hydrogen	Hydrogen
H_2O (see also Water), 15, 17, 35-38, 78, 82,	atomic (H), 13, 17, 18, 20-22, 27-29, 35,
271, 332-334	37-40, 49, 50, 55, 58-61, 66-68, 75,
Hadrons, 17, 19, 20	76, 82, 85-87, 101, 102, 107, 109,
Haem, 60, 62	116, 120, 142, 176, 186, 194, 199,
Hair(s), 175, 176, 208, 210, 211, 221, 246,	236, 265, 270, 271, 328, 337
261	bond(s), 34-37, 49, 55, 58, 63, 66,
Hair pin model, 82	76, 86, 96, 332, 333, 337
Handedness, 43-45, 48, 268, 269, 334, 356,	ionic (H ⁺), 84, 109, 169, 175, 271
357	channels, 87
Haploid, 137, 195	efflux, 235, 237
Hapten(s), 67, 66	gradient(s), 84, 102, 235
Head (cephalon), 15, 26, 59, 63, 115-117,	pump (ATPase), 167, 220
130, 211, 213, 218, 247-251, 257,	molecular (H_2), 17, 18, 82, 339
258, 260, 273, 345, 358	Hydrolytic (-ases), 65, 103
gradient(s), 50, 51, 87, 113, 247-249	Hydronium (H_3O^+), 37, 38, 168, 333
group(s), 76, 78, 340	Hydrophilic, 37, 39, 40, 63, 65, 75-77, 103,
	106, 110, 249
regeneration, 249, 250	Hydrophobic, 34, 37, 39, 40, 63, 65, 66, 75,
Heat, 29, 91, 190, 341	
Helical (helicoidal), 50, 51, 55, 57, 59, 68,	76, 78, 80, 84, 86, 96, 97, 106, 107,
70, 71, 86, 104, 115, 130, 131, 139,	141, 244
267, 328, 334, 336, 345, 356, 357	domain, 77
DNA, 50	Hydroxy(l) (OH ⁻), 37, 40, 68, 78, 82, 334
bacteria, 267, 356	Hymenium, 226, 228
protein, 59	Hyperpolarization, 89, 90, 108, 141, 142,
Helix, 44, 45, 47, 49, 50, 53, 58-60, 70, 71,	154, 187, 221, 236, 340
86, 87, 139, 327, 336	Hypha(e), 63, 107, 129, 147, 157, 161-168,
alpha, 76, 86, 106, 117	174, 181, 192-194, 197, 199, 204,
dipole(s), 58, 59	205, 225, 226, 228, 271, 325, 349
double, 49, 50, 113, 336	apex, 165, 190, 197, 200, 349
Heme, 65, 66	pole, 204, 345
Heterobipolar, 39, 41, 147, 181, 200-221,	tip(s), 105, 162-168, 174, 193, 194, 228,
326	349, 352
Heterogeneity, 79, 171, 172, 195	
Heterocysts (Cyanobacteria), 191, 192	Imaging, 124
Heterotrichous, 169	Immunochemistry, 66, 67, 103
Histidine, 40, 65, 140, 337	Immunofluorescence, 112, 117, 124, 152,
Hole(s), 22-24, 51, 109, 330	346
Homeobox, 260, 350	Incompatibility, 195, 196
Homeodomain, 260	Indole-3-acetic acid (IAA), 159, 186,
Homeotic mutants, 256, 260	230-232, 234, 235, 340, 355
Homobipolar, 13, 39, 147, 148, 181, 185,	asymmetry, 237
245, 326	Induction, 169, 200, 208, 215, 218, 222,
Hormone(s), 93, 95, 101, 102, 172, 186,	238, 239, 244, 246
209, 220, 227, 230-234, 236, 239,	endosporulation, 190
354, 355	polarity, 266, 267
Hyaloplasm, 72, 104, 111	prespore, 244
Hydration, 34, 73, 76, 330, 333	Information, 18, 23, 43, 56, 70, 89, 103,
Hydrocarbon, 39, 40, 42, 43, 63, 76, 78, 109	140, 144, 145, 149, 155, 162, 181,
22, 42, 43, 43, 70, 70, 70, 107	,,,,,,,,,,,,,,,,

197, 198, 200, 211, 214, 224, 227, 236, 243, 259, 262, 343, 353 Inheritance, 218 Inhibition, 64, 93, 139, 148, 155, 195, 231, 248, 249, 344	Kinase(s), 95, 148, 149, 162 Kinesin, 101, 123, 163, 179, 345 Kinetics, 43, 64 Kinetochore(s), 120, 124-128, 345 Kinetosomes, 244
contact, 80	Label(l)ing, 118, 131, 150, 153, 166, 204,
Inhibitor(s) 85, 99, 126, 187, 189, 192, 206,	350
211, 226, 228, 247, 351, 352	Lactic acid, 45
Insulin, 100, 101, 220 Integrin, 81	Lamellar, 63, 76
Interface, 34, 61, 75, 334, 338	Lamellipodia, 179, 180, 346
Internace, 34, 61, 75, 334, 336 Intermediate filaments, 111	Lamina-Laminin, 219, 220, 255, 256
Interposon, 55	Laser light, 23, 31
Intestinal, 219	Lateral-posterior direction, 265
Ion(s), 17, 25, 34, 37, 38, 41, 43, 65, 66, 68,	Lattice, 22, 24, 35, 38, 50, 64, 68, 72, 73, 79,
75, 82, 84-91, 93, 94, 102, 104, 110,	101, 111, 329 Layer(s) 75, 78, 130, 134, 150, 184, 201
135, 141, 142, 207, 249, 327, 329,	Layer(s), 75, 78, 129, 134, 150, 184, 201, 221, 225, 227, 232, 239, 330
333, 337, 340, 343, 355	Leading edge, 112, 135, 137, 144-146, 180,
Ionic	346, 347
channel(s), 86-89, 91, 93-95, 154, 180,	Leaf (leaves), 197, 205, 211, 230-232, 235,
340-342	238, 239
concentration, 201	Lectins, 79, 80
current flux(es), 110, 154, 181, 188, 230,	Left-right polarities, 267-270, 328, 356
349	Leucine, 40, 47, 48, 60, 87
gradient(s), 85, 175, 230, 236, 237	Leukocytes, 133, 137, 139, 141, 142, 146,
permeability, 90	346, 347
transport, 104, 107, 341	Li ⁺ (Lithium), 158, 217, 255
Ionization, 17, 31, 42, 237, 329	Life, 20, 34, 45, 46, 48, 74, 127, 138, 225,
Ionophore(s), 101, 133, 139, 160, 163, 170,	229, 267, 271, 273, 327, 334, 358
253, 261, 346 Iron (Fo ² +/3+), 14, 25, 27, 60, 66, 74, 84	Life cycle(s), 170, 238
Iron (Fe ^{2+/3+}), 14, 25-27, 60, 66, 74, 84	Ligand(s), 66, 76, 79, 80, 81, 85, 101, 113,
Irradiation, 169, 209 Isoelectric pH, 41, 164	141, 235, 337
Isometric, 38, 70, 133, 149, 205, 271	Light, 17, 23, 30-33, 43, 44, 46-48, 51, 66,
growth, 155	74, 80, 81, 87, 89, 94, 108-111, 129,
growth, 133	132, 142-144, 154, 159, 160, 169,
Innationa 112 218 267	170, 172, 174, 189, 194, 201,
Junctions, 112, 218, 267	205-210, 214-215, 226, 227, 236,
	242, 246, 258, 328-331, 340, 343, 352
K ⁺ (see also Potassium), 24, 40, 85, 87, 88,	bioelectric response, 239
91-94, 101, 106, 107, 141, 142, 154,	energy, 84, 109
158, 175, 237, 333, 341	excitation, 111
channels, 87, 88, 90-92, 94, 216, 221,	perception, 89, 336
341	polarization, 30, 32, 47, 124, 158, 174,
efflux, 194, 341	209, 210, 331
ionophore, 107	transducers, 108
Keratocytes, 145	Limb, 223, 231, 265, 356
Kidney (renal), 104, 105, 137, 218, 220,	Lipid(s), 62, 63, 75-79, 97, 164, 177, 198,
221, 353	331, 334, 338, 342, 344, 347

bilayer(s) (see also membranes), 72,	Mechanochemical
75-78, 87, 123, 338	enzymes, 346
flow model, 347	forces, 121, 123, 179, 180, 346
mosaic model, 76-78	Mechanoelectric transducers, 221
transport, 97	Meiosis, 252
Lipophilic, 235, 348	Membrane(s), 14, 63, 72-93, 95-102, 104,
Lipoprotein(s), 63, 77, 82	106-111, 114, 115, 123, 130-133,
Liver, 105, 249, 270	135, 137, 139-144, 148-150, 153,
Locomotion, 126, 130, 132-136, 138, 144,	154, 163, 177, 178, 180, 182, 186,
145, 179	191, 201, 204, 218-220, 267, 333,
Lomasome, 155	338-343, 346-348
Lymphocyte(s), 99	apical, 98, 175, 220, 353
Lysosome(s), 62, 95, 98, 99, 103, 164, 178,	asymmetric, 168
220	cellular, 15, 34, 58, 75, 76, 81, 86, 88,
Lysozyme, 59, 183	91-93, 95, 101, 104, 109, 115, 135,
	149
Macroconidium (-ia), 155, 349	differentiation, 99
Macrodipoles, 60, 61	domains, 204, 221
Macromolecule(s), 14, 15, 34, 49-71, 73,	plasmic (see Plasma membrane)
75, 79, 97, 155, 168, 170, 177, 206,	polarity, 85, 110, 191, 338, 339
262, 271, 332, 336, 337, 358	potential, 85-93, 106, 107, 146, 155-157,
conformation, 111	167, 188, 236, 239, 338-342
polarities, 49, 332, 358	primitive, 73
signals, 89	protein(s), 62, 76, 77, 79-81, 85, 96, 100,
Magnetic, 13, 24-28, 325, 330	103, 110, 175, 219, 221, 333, 343,
dipole(s), 30, 330	347
field(s), 14, 21, 24-26, 28, 30, 48, 175,	pump(s), 158, 167
327, 330, 331	receptor(s), 80, 89, 181
moment(s), 27, 29, 331	surface, 73-95, 101, 177, 332, 338-342
monopole(s), 26, 27	traffic-flow, 95, 102, 338, 347
particles, 108	Memory, 19, 23, 35, 95, 246, 329, 330
polarization, 25-30, 330, 331	Meristem, 210, 223, 224, 230, 238, 239
resonance, 30	cells, 147, 225, 230
Magnets, 13, 25, 27	Meromyosin, 112, 113, 121
Male, 161, 174, 197-200, 261, 327	Messenger RNA (mRNA), 53-55, 57, 58,
gametangia, 192, 198, 199	96, 155, 177, 208, 213, 214, 216, 217,
gradient, 199	257, 259, 261, 264, 272
mitochondria, 198	polarity, 58
mutant, 198	transport, 124
Maternal, 213, 218, 259, 267, 272	Messenger(s),
inheritance, 213, 214, 257, 259, 268, 269	chemical 89, 93, 94
mutations, 214, 256-259	Metabolism, 14, 53, 72, 73, 82, 85, 89, 91,
Mating, 152-154, 195, 197, 269	96, 100, 103, 132, 169, 170, 190, 217,
type(s), 56, 195-197	228, 233, 247, 255
Matrix, 42, 77, 78, 81, 84, 103, 105-107,	gradient(s), 208, 217, 247, 255
111, 130, 134, 181, 223, 255, 256,	transport, 82, 83
347	Metal(s), 22, 25, 68, 249, 339
Matter, 17-22, 24, 26, 27, 29, 44, 47, 67, 82,	ion(s), 40, 60, 65, 74, 337
147, 161, 271, 327-329, 334, 358	Metaphase, 120, 126, 128

M. d. 20, 20	Missania 100
Methane, 38, 39	Mitogenic, 199
Methanol, 39	Mitotic
Methylation, 140, 348	figures, 104, 122-128, 139, 151, 172, 345
Mg ²⁺ (magnesium), 81, 135	spindle, 112, 120, 124, 125, 151, 201,
Microbial spores, 14, 154	345
Microbodies, 103	Model(s), 19, 25, 49-51, 56, 61, 67, 70,
Microconidium (-ia), 193	75-78, 83, 86-88, 95, 97, 99, 101,
Microfibril(s), 162, 176, 211	103, 108, 117, 122, 124-126, 128,
Microfilament(s) (see also Actin-Myosin),	130, 132, 135, 137-139, 141-143,
15, 81, 99, 111-113, 115, 117, 119,	152, 157, 162, 167, 179, 183, 190,
121, 129, 130, 134, 138, 144, 157,	191, 199, 206, 210, 219, 224, 228,
163, 164, 175, 177, 188, 199, 200,	235, 237, 241, 243, 245, 247, 250,
211, 253, 267, 325, 344-346, 348, 350	254, 256, 260, 269, 332, 333, 340,
associated protein, 116	341, 347, 352, 356, 358
Microgravity, 236, 354	cell growth, 182, 183
Micropolarity, 338	membrane systems, 108, 109, 338
Microinjection, 142, 259	polarity, 22, 34, 88, 167, 183, 217, 220,
Microspikes, 179	240, 246, 250, 356, 358
Microtrabecular lattice, 73, 111	Modelling, 61, 67, 338
Microtubule(s) (see also Tubulins), 15, 81,	Molecular, 13, 14, 25, 29, 30, 34, 41, 42, 44,
95, 101, 105, 111-113, 118-129, 132,	46, 47, 50-52, 62, 63, 64, 66, 67, 69,
138, 139, 141, 144-146, 148,	71, 72, 75, 77, 85, 89, 94, 100, 109,
151-154, 157, 163, 164, 166, 167,	110, 112, 117, 131, 140, 144, 145,
173, 176-180, 186-189, 193, 199,	158, 170, 176, 177, 180, 189, 192,
200, 206, 211, 222, 229, 251, 253,	215, 240, 243, 244, 256, 269, 271,
271, 325, 342-344, 350, 357	272, 332-335
apical, 166	chirality, 43-48, 334-335
assembly-polymerization, 119, 121,	dipoles, 15, 33-43, 332
126, 187, 189, 222, 345	movement, 14, 33, 37
inhibitor, 152, 187	orientation, 14
MAPs (-associated proteins), 118, 350	polarity, 119, 121, 358
organizing centre (MTOC), 99, 121,	self-assembly, 350
342, 343	structure, 13, 62, 85, 337
polarity, 121, 122, 126, 127	Monensin, 99, 101
Microvillus, 115	Monoaxial patterns, 241-250
Microwave(s), 17, 37, 327	Monopodial, 133, 138
Mirror-image, 18, 44, 45, 69, 246, 268, 270,	Monopolar, 130, 132, 147-181, 197, 202,
334	272, 325, 348-351
Mitochondria, 74, 82-84, 96, 97, 103-108,	apical growth, 163-181, 197, 348, 349
110, 122, 130, 146, 154-157, 164,	assembly, 70
176, 168, 170, 173-175, 178, 179,	axiation, 107, 348, 349
184, 188, 190, 198, 205, 206, 209,	budding, 153
215, 218, 221, 261, 264, 336	differentiation, 202
apical-subapical, 155, 167	dominance, 163, 188, 271
ATP (ATPase H ⁺ pump), 82-85	germination, 155, 158, 171
bipolarity, 108	growth, 162, 167, 227, 348, 350
Ca^{2+} , 106	molecules, 39
membrane(s), 83, 104, 107, 108	outgrowth, 149-163
segregation, 198, 199, 261	pattern(s), 192-195, 352

regeneration, 251	Muscle(s), 85, 88-90, 93, 104-106, 112,
Monopolarity, 133, 165, 187, 352	114-117, 179, 190, 247, 340, 344
Monopole(s), 13, 27, 39, 61, 325	Mutant(s), 53, 54, 56, 58, 59, 118, 132, 140,
Morphogen(s), 14, 223, 228, 239, 240, 244,	142, 151, 152, 154, 183, 187, 188,
249, 250, 253, 256, 259, 265, 272,	193, 204, 226, 245-247, 256-261,
345, 356	268, 337, 344, 347, 355, 356
gradient, 14, 256	embryos, 257
transport, 248	morphological, 245, 256-261
Morphogenetic, 149, 151, 154, 162, 193,	Mutation(s), 53-55, 58, 70, 88, 154, 213,
200, 208, 209, 212, 213, 222-270,	252, 256, 257, 259, 260, 268, 269,
273, 325, 354-358	336, 356
determinants, 213, 253	Mycelium, 164-166, 193, 226, 340
factor, 265	
field(s), 14, 228, 245, 251	Myelin, 351
	Myofibril(s), 104
gradient(s), 217, 250, 255, 256, 264, 358	Myoplasm, 215, 216, 261
movement(s), 241, 264	Myosin, 15, 80, 81, 111-118, 125, 129, 130,
mutants, 154, 245, 257	134, 163, 167, 168, 179, 180, 211,
polarizations, 222-270, 354-358	344-347, 358
signal, 127	gene, 118
substances, 14, 354, 355	kinase, 116, 118
Morphopoiesis, 70, 337	fungal (myosin-like protein), 118
Morula, 266	Myxamoeba, 124
Mosaic model, 76, 77	N, see Nitrogen
Moss(es),	Na ⁺ , 37, 85, 89, 91-93, 106, 142, 158, 333
leaves, 211	-ATPase (pump), 91, 92, 221
protonema, 172, 173, 208-210	channels, 87, 91-94, 216
spores, 160, 208	NAD, 82, 84
Motion, 37, 43, 47, 51, 91, 110, 118, 125,	NADH, 186, 338, 340
131, 132, 135, 137, 139, 143, 146,	NADP, 84
179, 329, 332	NADPH, 109
Motor, 93, 94, 114, 118, 123, 131, 132, 140,	Negative, 18, 19, 22, 33, 35, 39, 41, 51, 53,
142, 143, 167, 179, 341, 343	61, 68, 70, 85, 88, 90, 92, 94, 106,
Movement(s), 14, 38, 52, 62, 81, 87, 88, 90,	108, 109, 131, 136, 140, 142, 143,
100-102, 112-116, 118, 120-123,	148, 167, 229, 325, 326, 330, 339,
125-146, 151, 162, 175, 178, 180,	345, 354
195, 214, 222, 231, 234, 262, 263,	charge(s), 20, 33-35, 39, 40, 57, 64, 67,
332, 336, 346, 347	82, 90, 108, 167, 175, 328, 345
auxin(s), 230, 233, 237, 354	electric potential, 235, 237, 327, 341
cytoplasm, 263	electrode, 209
intracellular, 114, 130, 346	Nerve(s), 68, 85, 86, 93, 149, 180, 181, 249,
polarity, 126, 266, 346, 347	252, 262, 342, 349, 350
Mucopolysaccharides, 64, 78	conduction, 91
	growth factor (NGF), 181
Multipolar, 147, 187, 188, 226, 326, 332,	induction, 238
348	
budding, 187, 247	Neurite(s), 126, 176-181, 271, 325, 342,
germination (outgrowth), 147, 187,	349-351 Neuron(s) 87-88-90-95-146-176-181-346
188, 348	Neuron(s), 87, 88, 90-95, 146, 176-181, 346, 350
moment, 33, 332	330

cytoskeleton-neurofilaments, 177, 179, 180	131, 138, 145, 146, 342-344, 349, 354
Neurotransmitter(s), 93-95, 102, 177, 178	Organizer (organizing center), 124, 238,
Neutron, 17, 22, 329, 333	241
NH ₃ (NH ₄), 38, 271, 353	Orientation, 13, 14, 29, 33, 41, 43, 62, 63,
Nigericin, 107, 108	79, 104, 112, 114, 116, 121, 122, 126,
Nitrogen (N ₂), 40, 58, 60, 165, 192	132, 142, 143, 148, 151, 158, 159,
fixing cell, 191	183, 185, 186, 190, 191, 205, 206,
Nuclear, 20, 30, 43, 45, 47, 57, 135, 164,	208, 211, 212, 229, 238, 244, 261,
169, 171, 198, 205, 208, 209, 244,	262, 331, 332, 336, 341
271, 327-329, 331, 345	Oscillator, 89
cap(s), 199	Osmotic pressure, 182, 193, 237
differentiation, 193, 254	Outgrowth(s), 108, 138, 149, 155, 157, 159,
dipoles, 30	160, 161, 169, 170, 180, 181, 184,
division, 125, 152, 205	210, 232, 342, 348, 352
fusion, 345	Oxalic acid, 39
magnetic resonance (NMR), 30, 73, 78,	Oxidation, 110, 340
331, 333, 340	phosphorylation, 85, 99, 106
matter, 17, 18	reduction potential(s), 162, 166, 339
membrane, 74, 95, 96, 128, 148, 151,	Oxygen (O-O ₂), 24, 25, 35, 37, 40, 58, 60,
152	65, 66, 82, 91, 92, 162, 186, 192, 198,
pore(s), 123, 124	327, 332, 339, 340, 343
Nucleation, 81, 114, 127, 128, 134, 344	Pancreas, 220
Nucleus (nuclei), 17, 19-21, 28-30, 33, 35,	Parallel
38, 72-74, 89, 97, 101, 104, 112-114,	dipoles, 38
123-128, 130, 137, 145, 146,	polarity, 50
148-150, 152, 158, 161, 170-173,	Paramagnetic, 26, 30
188, 189, 193, 195, 198-200, 205,	Paramyosin, 117
206, 208-211, 214, 216, 218, 225,	Parenchyma, 147, 233
236, 244, 252, 255, 258, 259, 271, 327-329, 331, 345	Parity, 18, 44, 45, 47, 327-330, 334
	Particle(s), 17-22, 26-31, 34, 43, 44, 48, 67,
Nucleic acid(s) (see also DNA, RNA), 15,	69-71, 77, 111, 122, 131, 133, 201,
49- 58, 69, 70, 74, 75, 214, 332, 336, 337	207, 271, 327-329
Nucleosomes, 53	Patches, 80, 81
Nucleosynthesis, 17	Pattern(s), 13, 14, 26, 61, 72, 115, 119, 131,
Nucleotide(s), 51, 57, 73, 119, 139, 157,	137, 138, 143, 145, 152, 154, 155,
174, 256	159, 160, 165, 170, 172, 175,
Nutrient(s), 60, 93, 101, 169, 175, 203, 348	182-185, 188, 189, 192-200, 207,
14411611(5), 60, 73, 101, 107, 173, 203, 346	212, 215-217, 222, 223, 227, 228,
O, see Oxygen	236, 238, 240, 241, 244, 246, 249,
Oligosaccharides, 75, 79	250, 255, 256, 259, 260, 265, 268,
Ontogenesis (ontogeny), 160, 193, 224	271, 272, 348, 349, 356, 357
Ooplasm, 215, 216, 254, 261	formation, 154, 215, 222, 238, 247
Operon, 53-55	polarizing current, 231
Optical, 23, 30, 44, 45, 48, 61, 119, 155, 328	regulation, 217, 240, 241
Oral (structures), 244-246	D . 147 104
	Pennate, 147, 184
Organelle(s), 72, 89, 94, 96, 97, 100-102,	Pennate, 147, 184 Peptide(s), 45, 53, 58-60, 62, 66, 80, 86, 87, 95, 107, 181, 249, 343

dipole(s), 59, 60	brown, 91, 159, 160, 169, 172, 195, 224,
Peptidoglycan, 182	244, 349
Perception, 236, 340, 343, 354, 355	green, 119, 130, 143, 147, 169, 170, 185,
Periodic, 88, 330, 336	188, 190, 208, 228, 340, 341, 349
Periplasm, 60, 194	grey, 255, 262, 263
Permeability, 82, 85, 87, 89, 91, 338	orange, 198, 199, 216, 254, 261, 262
Peroxide(s)-Peroxidase(s), 65, 66, 164	red, 75, 111, 172, 185, 216, 220, 349,
Peroxisomes, 103	353
pH, 37, 40, 41, 50, 80, 82, 84, 101, 106, 129,	yellow, 147, 169, 195, 216, 261,
159, 163, 166, 185-187, 201, 206,	Pinocytosis, 95
220, 223, 230, 235, 237, 340, 353	Plant(s), 13, 15, 32, 73, 79, 80, 89, 91, 101,
gradient, 84, 101, 102, 109, 159, 235	102, 124, 129, 147, 161, 169, 170,
intracellular, 168	172, 175, 185, 186, 189, 197, 200,
Phage(s), 53, 70, 71, 182, 337	204, 208-210, 212, 222-225, 229-233,
Phalloidin, 116, 152, 153, 174	235, 236, 238, 239, 268, 325, 340,
Phialide, 193	343, 354, 355
Phloem, 233	axis, 229, 233
Phosphate(s), 40, 70, 55, 62, 63, 92, 100,	cell elongation, 185
103, 149, 166	eggs-embryo, 154, 225, 325
Phospholipase, 88	growth substance(s), 186, 235, 355
Phospholipid(s), 62, 63, 65, 75-78, 80, 96,	Plasma membrane, 63, 74, 75, 77, 78,
97, 107, 109, 338, 339	80-82, 85, 89-91, 95, 97, 99-104, 106,
Phosphoprotein, 177, 346	111, 112, 115, 132, 134-136, 141,
Phosphorylation, 64, 84, 94, 103, 117, 118,	143, 149, 150, 154, 157, 162, 166,
134, 140, 141, 181	173, 175, 185, 186, 195, 207, 216,
Photo-	218, 219, 235, 269, 338, 352-355
chemistry, 42, 47, 48, 108	domain(s), 218-220
excitation, 42	polarization, 220
	Plasma, 24, 137, 214, 327
periodic induction, 239	Plasmalemma, 98, 142, 144, 164, 166, 167,
polarization, 206	173-176, 180, 184, 194, 235-237,
receptor(s), 89, 111, 142	254, 338
dipoles, 174	Plasmid, 118
synthesis, 43, 47, 48, 108-111, 142	
apparatus, 74	Plasmodium, 129, 133, 143
magnetic particles, 108	Plate(s), 32, 60, 70, 131, 174, 221, 228
reaction center, 110	Polar, 13, 14, 33, 35, 37-41, 43, 53-55, 58,
system(s), 110, 343	60-62, 64, 66-68, 70, 73, 75, 76, 78,
taxis, 142, 143	80, 83, 86, 87, 97, 108, 111, 112, 115,
transducers, 74	117, 119, 120, 125, 128-131, 135,
voltaic, 108	136, 137, 140, 143-145, 154, 160,
Photon(s), 17, 21-23, 27, 30-32, 47, 89,	163, 166, 170, 172, 173, 181, 183,
108-111, 209, 329	189, 191, 198, 199, 201, 204, 211,
Phycobilisomes, 111	214, 215, 224-226, 228, 232-236,
Phyllotaxis, 197, 232	241, 242, 244, 246, 247, 249, 267,
Phytochrome, 89, 174, 239, 340	268, 271, 272, 331, 346-349, 352-354
Pigment(s), 81, 89, 100, 110, 142, 207, 213,	blastomeres, 264
217, 242, 261, 269, 336, 340, 343	assembly, 179, 202
blue-green, 132, 133, 143	auxin transport, 233-235, 354

axiation, 13, 81, 147, 154, 160, 165, 173,	265-267, 271-273, 325-330, 332-334,
174, 200, 201, 206, 207, 212, 224,	336-358
228, 241, 242, 257, 258, 263, 265,	cellular, 157, 158, 229, 240
266, 271, 346, 353, 355	control(s), 22, 56, 62, 143, 157, 171,
bodies, 212, 218, 266	222, 236, 238, 244, 245, 248, 256,
bonds, 76	257
	cytoplasmic, 210, 211, 262
budding, 151	developmental, 227, 257, 352
cell(s), 97, 176, 217-222, 266	egg(s), 157-159, 210-212, 253-259
development, 73	
differentiation, 189, 193, 195, 201,	embryonic, 224, 225, 249, 257
346	hyphal, 166, 348, 349
divisions, 157, 158, 205, 210, 211,	intrinsic, 14, 15, 51, 115, 119-121, 132,
225	182, 244, 245, 248, 250, 271, 272,
growth, 147, 151, 164, 173, 176,	358
186, 194, 348	inverse, 49, 58, 242
	macromolecular, 14, 332, 334, 358
movements, 15, 84, 112, 162, 232,	organismic, 225-239, 358
233, 266	pigments, 216, 261, 340, 343
conduction, 38	replication, 52
cytoplasm, 13, 200, 208, 214, 215	reversal, 14, 50, 51, 92, 133, 141, 143,
domain(s), 64, 107, 203	174, 207, 220, 232-234, 247, 265,
elongation, 161, 171, 173, 180, 181,	The second secon
195, 201	268, 272, 343
genes, 259	systems, 244
gradient(s), 173, 176, 205, 234, 264, 345	tissular, 14, 262
granules, 256	Polarizability, 332
	Polarizable particles, 19, 23, 34, 330, 332
group(s), 50, 55, 63, 75-78	Polarization, 14, 17, 18, 24, 29-34, 37, 40,
liquid(s), 37, 68	41, 43, 45, 47, 48, 56, 89, 99, 101,
lobe(s), 213, 252, 253, 255, 269	104, 105, 107, 109, 123, 127, 132,
molecules, 14, 33, 35, 37-41, 43, 62, 63,	137, 141, 145, 146, 149, 152, 155,
78, 80, 350	157-160, 177, 186, 205, 209, 210,
mutants, 54	
pattern(s), 14, 26, 222, 239, 241	212, 216, 218, 220, 230, 254, 264,
polymerization, 114	266, 267, 327-333, 338-345, 348,
regeneration, 166, 234, 246, 251	352-358
segregation, 199, 343, 345	analyzers, 32
signal, 157, 261	field, 18
site, 154	light, 30-32, 331
	magnetic, 25-30, 330, 331
Polarimetry, 32, 331	Polarized, 13, 15, 23, 25, 28, 29, 31-34, 40,
Polarity, 13-15, 18, 23, 25, 34, 37, 40, 43,	47, 48, 50, 52, 60, 61, 64, 65, 79, 80,
50-57, 60-62, 66, 67, 72, 79, 81, 82,	97-99, 104, 106, 111, 113, 126, 131,
86, 94, 98, 99, 101, 102, 107,	134, 138, 141, 142, 144, 145, 153,
112-122, 124, 126-134, 137, 138,	155, 161, 162, 165, 166, 169, 173,
143-145, 147, 148, 154, 157-160,	176, 198, 201, 211, 216, 218-221, 232, 240, 242, 247, 256, 250, 265
165, 166, 168-176, 179, 181, 183,	232, 240-243, 247, 256, 259, 265,
185-191, 200, 201, 204-210, 212,	271, 273, 325-331, 336-341, 344,
214, 216-218, 220-222, 224, 225,	345, 349, 350, 352, 353
227-229, 231-234, 236, 238, 239,	actin, 168, 344
241, 242, 244-252, 256-258, 260-263,	beams of particles, 30

bonds, 62, 85	Polymerization, 15, 47, 70, 112-114,
communication, 243	119-121, 128, 131, 134, 179, 344-345
conductivity, 22-25, 329, 330	Polypeptides, 45, 55, 68, 196
control, 56	Polypodial, 133
direction, 230	Polypolarity (also Multipolar), 133, 147,
enzymatic reactions, 64, 65	326, 332
flow, 180	Polyribosome(s), 96, 103, 264
form, 177	Polysaccharide(s), 64, 134, 153, 175, 187,
gas, 29	206, 237, 355
genetic expression, 198, 199, 358	Pore(s), 75, 79, 86, 87, 99, 123, 124, 155,
growth, 15, 118, 129, 149, 155, 183,	160, 211
226, 348-351	Porphyrin(s), 42, 60, 74
ionic regulation, 232	Positional, 117, 172, 176, 180, 197, 199,
light, 32, 47, 124, 158, 174, 209, 210	223, 227, 241, 243, 244, 247, 256,
movements, 126, 266	265, 358
organelles arrangement, 180	control(s), 195, 222, 231, 243
organization, 14, 148, 172, 195, 236,	DNA, 198, 199, 336, 358
241, 246	information, 14, 173, 197, 209, 211,
orientation, 104	223, 228, 240, 252, 260
photons, 45, 329	Positive, 18-20, 22, 33, 35, 39, 40, 61, 65,
secretion, 101, 220, 342, 343	67, 68, 70, 83, 87, 90, 91, 93, 94, 102,
spores, 155-157, 187	107-109, 140, 142-144, 148, 167,
synapses, 93	168, 170, 325-327, 336-340
translocation-transport, 62, 84, 101,	charge(s), 20, 33, 35, 59, 68, 71, 82, 91,
106, 107, 166, 178, 229, 233-236, 272	94, 168, 327-329, 338, 343
Polarizing field, 34	current, 161, 175, 194
region(s), 265, 356	electric potential, 235
Polarography, 339	electrode, 209
Polaron(s), 24, 51, 56, 68	feedback, 230
Polarotropism, 174	gravitropism, 236, 237
Pole(s), 13, 14, 27, 33, 41, 49, 72, 82, 100,	hole(s), 22, 108, 109
107, 120, 124-126, 128, 130, 150,	ions, 236
158-161, 167, 171, 181-184, 190,	Positron(s), 17-19, 21, 22, 31, 44, 271, 327,
201, 204, 212-215, 217, 219, 220,	334
224, 229, 249, 254-256, 259, 263,	Posterior, 256
272, 325, 345, 352-355	pole, 213, 229, 245, 246, 256, 258, 263
north/south, 13, 28, 175, 246, 249, 325	region-segment, 244, 245, 250, 251,
rhizoidal, 158, 204-210, 352	257, 258, 261, 262-265, 346
thallic, 181, 204-210, 352	structures, 252, 259, 346
Poleward(s), 125-128, 345	Postsynaptic, 93-95, 180
Pollen	Potassium (see also K ⁺), 24, 85, 91-94
grain(s), 161, 163, 175, 205	Potential(s) (see also Electric-), 21, 30, 35,
tube(s), 147, 161, 164, 169, 174, 175,	42, 66, 74, 84-86, 88, 90, 91, 94, 109,
186	114, 131-133, 136, 142, 146, 148,
Polyamine(s), 155, 348	170, 176, 180, 225, 237, 243, 253,
Polyglutamic acid, 135	272, 338-341, 346
Polymer(s), 15, 64, 67, 68, 78, 112-114,	depolarization, 236, 334, 346
119, 121, 135, 183, 345	difference(s), 89, 170, 230
,,,,,	

polarization, 340 Prebiotic, 46-48, 57, 73 Precursor(s), 14, 57, 70, 74, 82, 103, 107, 124, 144, 166, 183, 191, 233, 252 Prepolar organization, 218 Pressure (see also Osmotic), 23, 35, 133, 182, 185, 193, 194, 237 Presynaptic, 93-95, 108 Primitive, 39, 73, 74, 108, 132, 137, 190, 265, 266 environment, 46 Probe(s) (see also Fluorescence), 50, 61, 152, 174, 204, 207, 340 Protein(s), 13, 15, 34, 38, 45, 50-53, 55-63, 66, 68-83, 86, 87, 89, 94-104, 106, 107, 109-119, 122-124, 126, 129, 131, 132, 134-136, 139, 140, 148, 149, 151, 154, 155, 158, 161-163, 168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Protonlail, 173, 174, 209 Protonlail, 174, 175, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 plow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonombotive force, 84, 106 Protonophore(s), 108, 346 Protonalian, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 237, 223, 243, 244, 2245-248, 346, 350, 350	gradient(s), 93, 94, 109, 170	Pseudoplasmodium, 241-243
Precursor(s), 14, 57, 70, 74, 82, 103, 107, 124, 144, 146, 183, 191, 233, 252 Prepolar organization, 218 Pressure (see also Osmotic), 23, 35, 133, 182, 185, 193, 194, 237 Presynaptic, 93-95, 108 Primitive, 39, 73, 74, 108, 132, 137, 190, 265, 266 environment, 46 Probe(s) (see also Fluorescence), 50, 61, 152, 174, 204, 207, 340 Profilin, 114 Promotor, 52, 53, 55 Protease, 194, 259, 337 Protein(s), 13, 15, 34, 38, 45, 50-53, 55-63, 66, 68-83, 86, 87, 89, 94-104, 106, 107, 109-119, 122-124, 126, 129, 131, 132, 134-136, 139, 140, 148, 149, 151, 154, 155, 158, 161-163, 168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 218-221, 225, 235, 237, 243, 244, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Protono(s), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 293, 102, 106, 154, 158, 186, 220, 221, 331, 341, 355 Purines, 49 Pyrinc, 108 Quantum, 17, 24, 26, 27, 29-31, 67, 42, 109, 329, 333 Quantum mechanics, 272, 328, 329, 333 Quantum mechanics, 272, 328, 329, 333 Quantum theory, 29, 333 Quantum theory, 29, 333 Quantum theory, 29, 333 Posteric, 45-48 Radial, 112, 241, 148, 226, 262, 264, 271 Radiation(s), 17, 21, 25, 30, 33, 45, 47, 48, 67, 327, 331 Reaction center(s), 110, 111, 343 -diffusion, 247, 249, 250 transfer, 339 Reactivity, 50, 337 Receptor(s), 60, 80-82, 93-96, 101-103, 107, 134, 135, 139-141, 143, 217, 220, 221, 235, 340, 341, 354, 355 Recognition sequence, 336 system(s), 51, 57, 58, 79, 81, 96, 162 Red light, 89, 142, 174 Redox, 17, 16, 68, 74, 82, 143, 166, 186, 169, 172, 173, 208-210 Protomotive force, 84, 106 Protonophore(s), 108, 346 Protonomior, 69, 154 Rediction, 19, 20, 29 Reactivity, 50, 337 Receptor(s), 60, 80-82, 93-96, 101-103, 107, 134, 135, 139-141, 143, 217, 220, 221, 235, 230, 240, 242, 245-248, 232,	polarization, 340	Pseudopodium-pseudopodia, 126, 133,
124, 144, 166, 183, 191, 233, 252 Prepolar organization, 218 Pressure (see also Osmotic), 23, 35, 133, 182, 185, 193, 194, 237 Presynaptic, 93-95, 108 Primitive, 39, 73, 74, 108, 132, 137, 190, 265, 266 environment, 46 Probe(s) (see also Fluorescence), 50, 61, 152, 174, 204, 207, 340 Profilin, 114 Promotor, 52, 53, 55 Protease, 194, 259, 337 Protein(s), 13, 15, 34, 38, 45, 50-53, 55-63, 66, 68-83, 86, 87, 89, 94-104, 106, 107, 109-119, 122-124, 126, 129, 131, 132, 134-136, 139, 140, 148, 149, 151, 154, 155, 158, 161-163, 168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 218-221, 225, 235, 237, 243, 244, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 20, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonomotive force, 84, 106 Protonomotive, 83, 102, 108, 235, 237, 243, 244, 244, 226, 262, 264, 271 Reaction sequence, 336 Reaction sequence, 336 Reaction, 13, 132, 134, 135, 139-141, 143, 217, 220, 221, 235, 340, 341, 354, 355 Receptorics), 60, 80-82, 93-96, 101-103, 107, 134, 135, 139-141, 143, 217, 220, 221, 235, 340, 341, 354, 355 Recognition sequence, 336 Reduction, 54, 82, 84, 90, 91, 109, 110, 2	Prebiotic, 46-48, 57, 73	134, 136, 138, 326, 342
Prepolar organization, 218 Pressure (see also Osmotic), 23, 35, 133, 182, 185, 193, 194, 237 Presynaptic, 93-95, 108 Primitive, 39, 73, 74, 108, 132, 137, 190, 265, 266 environment, 46 Probe(s) (see also Fluorescence), 50, 61, 152, 174, 204, 207, 340 Profilin, 114 Promotor, 52, 53, 55 Proteans, 194, 259, 337 Protein(s), 13, 15, 34, 38, 45, 50-53, 55-63, 66, 68-83, 86, 87, 89, 94-104, 106, 107, 109-119, 122-124, 126, 129, 131, 132, 134-136, 139, 140, 148, 149, 151, 154, 155, 158, 161-163, 168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 218-221, 225, 235, 237, 243, 244, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonmobre(s), 108, 346 Protonation, 50, 154 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 41 Resistance, 50, 67, 216	Precursor(s), 14, 57, 70, 74, 82, 103, 107,	Pump(s), 82, 93, 102, 106, 154, 158, 186,
Prepolar organization, 218 Pressure (see also Osmotic), 23, 35, 133, 182, 185, 193, 194, 237 Presynaptic, 93-95, 108 Primitive, 39, 73, 74, 108, 132, 137, 190, 265, 266 environment, 46 Probe(s) (see also Fluorescence), 50, 61, 152, 174, 204, 207, 340 Profilin, 114 Promotor, 52, 53, 55 Proteans, 194, 259, 337 Protein(s), 13, 15, 34, 38, 45, 50-53, 55-63, 66, 68-83, 86, 87, 89, 94-104, 106, 107, 109-119, 122-124, 126, 129, 131, 132, 134-136, 139, 140, 148, 149, 151, 154, 155, 158, 161-163, 168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 218-221, 225, 235, 237, 243, 244, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonmobre(s), 108, 346 Protonation, 50, 154 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 41 Resistance, 50, 67, 216	124, 144, 166, 183, 191, 233, 252	220, 231, 341, 355
Pressure (see also Osmotic), 23, 35, 133, 182, 185, 193, 194, 237 Presynaptic, 93-95, 108 Primitive, 39, 73, 74, 108, 132, 137, 190, 265, 266 environment, 46 Probe(s) (see also Fluorescence), 50, 61, 152, 174, 204, 207, 340 Profilin, 114 Promotor, 52, 53, 55 Protease, 194, 259, 337 Protein(s), 13, 15, 34, 38, 45, 50-53, 55-63, 66, 68-83, 86, 87, 89, 94-104, 106, 107, 109-119, 122-124, 126, 129, 131, 132, 134-136, 139, 140, 148, 149, 151, 154, 155, 158, 161-163, 168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 218-221, 225, 235, 237, 243, 244, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Protononos, 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protomophore(s), 108, 346 Protonophore(s), 108, 346 Protonation, 50, 154 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplasm(s), 78, 79, 162, 167, 202, 230,		
Presynaptic, 93-95, 108 Primitive, 39, 73, 74, 108, 132, 137, 190, 265, 266 environment, 46 Probe(s) (see also Fluorescence), 50, 61, 152, 174, 204, 207, 340 Profilin, 114 Promotor, 52, 53, 55 Protease, 194, 259, 337 Protein(s), 13, 15, 34, 38, 45, 50-53, 55-63, 66, 68-83, 86, 87, 89, 94-104, 106, 107, 109-119, 122-124, 126, 129, 131, 132, 134-136, 139, 140, 148, 149, 151, 154, 155, 158, 161-163, 168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 218-221, 225, 235, 237, 243, 244, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonomotive force, 84, 106 Protonophore(s), 108, 346 Protonation, 50, 154 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplasm(s), 78, 79, 162, 167, 202, 230,		
Presynaptic, 93-95, 108 Primitive, 39, 73, 74, 108, 132, 137, 190, 265, 266 environment, 46 Probe(s) (see also Fluorescence), 50, 61, 152, 174, 204, 207, 340 Profilin, 114 Promotor, 52, 53, 55 Protease, 194, 259, 337 Protein(s), 13, 15, 34, 38, 45, 50-53, 55-63, 66, 68-83, 86, 87, 89, 94-104, 106, 107, 109-119, 122-124, 126, 129, 131, 132, 134-136, 139, 140, 148, 149, 151, 154, 155, 158, 161-163, 168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 218-221, 225, 235, 237, 243, 244, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonomotive force, 84, 106 Protonophore(s), 108, 346 Protonation, 50, 154 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230,		mana [®] com
Primitive, 39, 73, 74, 108, 132, 137, 190, 265, 266 environment, 46 Probe(s) (see also Fluorescence), 50, 61, 152, 174, 204, 207, 340 Profilin, 114 Promotor, 52, 53, 55 Protease, 194, 259, 337 Protein(s), 13, 15, 34, 38, 45, 50-53, 55-63, 66, 68-83, 86, 87, 89, 94-104, 106, 107, 109-119, 122-124, 126, 129, 131, 132, 134-136, 139, 140, 148, 149, 151, 154, 155, 158, 161-163, 168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 218-221, 225, 235, 237, 243, 244, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonomotive force, 84, 106 Protonophore(s), 108, 346 Protonophore(s), 108, 346 Protonomotive force, 84, 106 Protonophore(s), 108, 346 Protonophore(s), 108, 346 Protonophore(s), 108, 346 Protonation, 50, 154 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230,	The second secon	
265, 266 environment, 46 Probe(s) (see also Fluorescence), 50, 61, 152, 174, 204, 207, 340 Profilin, 114 Promotor, 52, 53, 55 Protease, 194, 259, 337 Protein(s), 13, 15, 34, 38, 45, 50-53, 55-63, 66, 68-83, 86, 87, 89, 94-104, 106, 107, 109-119, 122-124, 126, 129, 131, 132, 134-136, 139, 140, 148, 149, 151, 154, 155, 158, 161-163, 168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 218-221, 225, 235, 237, 243, 244, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonmotive force, 84, 106 Protonophore(s), 108, 346 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230,		Quadrupole(s), 38, 326, 331, 333, 340
environment, 46 Probe(s) (see also Fluorescence), 50, 61, 152, 174, 204, 207, 340 Profilin, 114 Promotor, 52, 53, 55 Protease, 194, 259, 337 Protein(s), 13, 15, 34, 38, 45, 50-53, 55-63, 66, 68-83, 86, 87, 89, 94-104, 106, 107, 109-119, 122-124, 126, 129, 131, 132, 134-136, 139, 140, 148, 149, 151, 154, 155, 158, 161-163, 168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 218-221, 225, 235, 237, 243, 244, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonomotive force, 84, 106 Protonophore(s), 108, 346 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 420, 242, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 420, 242, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 420, 242, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 420, 242, 245-248, 250, 251, 351, 352		
Probles (see also Fluorescence), 50, 61, 152, 174, 204, 207, 340 Profilin, 114 Promotor, 52, 53, 55 Protease, 194, 259, 337 Protein(s), 13, 15, 34, 38, 45, 50-53, 55-63, 66, 68-83, 86, 87, 89, 94-104, 106, 107, 109-119, 122-124, 126, 129, 131, 132, 134-136, 139, 140, 148, 149, 151, 154, 155, 158, 161-163, 168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Protononotive force, 84, 106 Protonomotive force, 84, 106 Protonoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplask(s), 78, 79, 162, 167, 202, 230,		
152, 174, 204, 207, 340 Profilin, 114 Promotor, 52, 53, 55 Protease, 194, 259, 337 Protein(s), 13, 15, 34, 38, 45, 50-53, 55-63, 66, 68-83, 86, 87, 89, 94-104, 106, 107, 109-119, 122-124, 126, 129, 131, 132, 134-136, 139, 140, 148, 149, 151, 154, 155, 158, 161-163, 168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonnotive force, 84, 106 Protonophore(s), 108, 346 Protonophore(s), 108, 346 Protonophore(s), 108, 346 Protonoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplasm(s), 78, 79, 162, 167, 202, 230,		
Profilin, 114 Promotor, 52, 53, 55 Protease, 194, 259, 337 Protein(s), 13, 15, 34, 38, 45, 50-53, 55-63, 66, 68-83, 86, 87, 89, 94-104, 106, 107, 109-119, 122-124, 126, 129, 131, 132, 134-136, 139, 140, 148, 149, 151, 154, 155, 158, 161-163, 168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonnotive force, 84, 106 Protonophore(s), 108, 346 Protonophore(s), 108, 346 Protonophore(s), 108, 346 Protonoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplasm(s), 78, 79, 162, 167, 202, 230,		Secretary Sea Secretary Secre
Promotor, 52, 53, 55 Protease, 194, 259, 337 Protein(s), 13, 15, 34, 38, 45, 50-53, 55-63, 66, 68-83, 86, 87, 89, 94-104, 106, 107, 109-119, 122-124, 126, 129, 131, 132, 134-136, 139, 140, 148, 149, 151, 154, 155, 158, 161-163, 168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 218-221, 225, 235, 237, 243, 244, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 19, 20, 329 Quartz crystal(s), 46 Quinone, 42 Racemic, 45-48 Radial, 112, 241, 148, 226, 262, 264, 271 Radiaton(s), 17, 21, 25, 30, 33, 45, 47, 48, 67, 327, 331 Reaction center(s), 110, 111, 343 -diffusion, 247, 249, 250 transfer, 339 Reactivity, 50, 337 Receptor(s), 60, 80-82, 93-96, 101-103, 107, 134, 135, 139-141, 143, 217, 220, 221, 235, 340, 341, 354, 355 Recognition sequence, 336 system(s), 51, 57, 58, 79, 81, 96, 162 Red light, 89, 142, 174 Redox, 17, 61, 68, 74, 82, 143, 166, 186, 223, 264, 327, 339 Reduction, 54, 82, 84, 90, 91, 109, 110, 264 Regeneration, 13, 92, 132, 170-173, 177, 207, 225, 230, 240, 242, 245-248, 250, 251, 351, 352 Regulation, 51, 54 Protonomotive force, 84, 106 Protonophore(s), 108, 346 Protonophore(s), 108, 346 Protonophore(s), 108, 346 Protonophore(s), 211 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 230, 230, 240, 242, 211, 133, 136, 340 Reproduction, 712, 179, 200, 238, 241 Reaction Rediation(s), 17, 21, 25, 30, 33, 45, 47, 48, 46d, 67, 327, 331 Reaction center(s), 10, 111, 343 -diffusion, 247, 249, 250 transfer, 339 Reactivity, 50, 337 Receptor(s), 60, 80-82, 93-96, 101-103, 107, 134, 135, 139-141, 143, 217, 220, 221, 235, 340, 341, 354, 355 Recognition sequence, 336 system(s), 51, 57, 58, 79		0.00
Protease, 194, 259, 337 Protein(s), 13, 15, 34, 38, 45, 50-53, 55-63, 66, 68-83, 86, 87, 89, 94-104, 106, 107, 109-119, 122-124, 126, 129, 131, 132, 134-136, 139, 140, 148, 149, 151, 154, 155, 158, 161-163, 168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonation, 50, 154 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 230, 66, 68-83, 86, 87, 89, 94-104, 106, 108-110, 124, 143, 167, 168, 185 Protonomotive force, 84, 106 Protonoplore(s), 108, 346 Protonoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 230, 230, 230, 231, 232, 232, 242, 245-248, 250, 251, 351, 352		
Protein(s), 13, 15, 34, 38, 45, 50-53, 55-63, 66, 68-83, 86, 87, 89, 94-104, 106, 107, 109-119, 122-124, 126, 129, 131, 132, 134-136, 139, 140, 148, 149, 151, 154, 155, 158, 161-163, 168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonophore(s), 108, 346 Protonophore(s), 108, 346 Protonoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 230, 230, 237, 247, 205, 2211 Protoplast(s), 78, 79, 162, 167, 202, 230, 230, 236, 237, 247, 249, 250 protection (s), 12, 144, 143, 167, 168, 185, 175, 176, 186, 185, 176, 186, 185, 176, 186, 185, 176, 186, 185 Protonomotive force, 84, 106 Protonoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 230, 230, 230, 230, 230, 231, 232, 243, 244, 242, 111, 133, 136, 340 Recemic, 45-48 Radial, 112, 241, 148, 226, 262, 264, 271 Readiation(s), 17, 21, 25, 30, 33, 45, 47, 48, 67, 327, 331 Reaction center(s), 110, 111, 343 -diffusion, 247, 249, 250 transfer, 339 Reactivity, 50, 337 Receptor(s), 60, 80-82, 93-96, 101-103, 107, 134, 135, 139-141, 143, 217, 220, 221, 235, 340, 341, 354, 355 Recognition sequence, 336 system(s), 51, 57, 58, 79, 81, 96, 162 Red light, 89, 142, 174 Redox, 17, 61, 68, 62, 20, 341 sink(s), 167, 168, 185 Protonomotive force, 84, 106 Protonophore(s), 108, 346 Protoplasm, 74, 209 Protoplasm,		
66, 68-83, 86, 87, 89, 94-104, 106, 107, 109-119, 122-124, 126, 129, 131, 132, 134-136, 139, 140, 148, 149, 151, 154, 155, 158, 161-163, 168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 218-221, 225, 235, 237, 243, 244, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonophore(s), 108, 346 Protonophore(s), 108, 346 Protonophore(s), 108, 346 Protonoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230,		
107, 109-119, 122-124, 126, 129, 131, 132, 134-136, 139, 140, 148, 149, 151, 154, 155, 158, 161-163, 168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 218-221, 225, 235, 237, 243, 244, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Prothallia, 173, 174, 209 Prothonosis, 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonophore(s), 108, 346 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 230, 238, 241 Protoplast(s), 78, 79, 162, 167, 202, 230, 230, 230, 240, 228, 241 Reacmic, 45-48 Radial, 112, 241, 148, 226, 262, 264, 271 Radiation(s), 17, 21, 25, 30, 33, 45, 47, 48, 67, 327, 331 Reaction center(s), 110, 111, 343 -diffusion, 247, 249, 250 transfer, 339 Reactivity, 50, 337 Reccpitor(s), 60, 80-82, 93-96, 101-103, 107, 134, 135, 139-141, 143, 217, 220, 221, 235, 340, 341, 354, 355 Recognition sequence, 336 system(s), 51, 57, 58, 79, 81, 96, 162 Red light, 89, 142, 174 Redox, 17, 61, 68, 74, 82, 143, 166, 186, 223, 264, 327, 339 Reduction, 54, 82, 84, 90, 91, 109, 110, 264 Regeneration, 13, 92, 132, 170-173, 177, 207, 225, 230, 240, 242, 245-248, 250, 251, 351, 352 Regulation, 51, 54, 86, 91, 114, 148, 157, 162, 184, 188, 189, 203, 350, 331		(
131, 132, 134-136, 139, 140, 148, 149, 151, 154, 155, 158, 161-163, 168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 218-221, 225, 235, 237, 243, 244, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonophore(s), 108, 346 Protonophore(s), 108, 346 Protonophore(s), 108, 346 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 236, 237, 241, 241, 148, 226, 262, 264, 271 Radial, 112, 241, 148, 226, 262, 264, 271 Radiation(s), 17, 21, 25, 30, 33, 45, 47, 48, 67, 327, 331 Reaction center(s), 110, 111, 343 -diffusion, 247, 249, 250 transfer, 339 Reactivity, 50, 337 Receptor(s), 60, 80-82, 93-96, 101-103, 107, 134, 135, 139-141, 143, 217, 220, 221, 235, 340, 341, 355 Recognition sequence, 336 system(s), 51, 57, 58, 79, 81, 96, 162 Red light, 89, 142, 174 Redox, 17, 61, 68, 74, 82, 143, 166, 186, 223, 264, 327, 339 Reduction, 54, 82, 84, 90, 91, 109, 110, 264 Regeneration, 13, 92, 132, 170-173, 177, 207, 225, 230, 240, 242, 245-248, 250, 251, 351, 352 Recution(s), 12, 25, 30, 33, 45, 47, 48, 67, 327, 331 Reaction center(s), 110, 111, 343 -diffusion, 247, 249, 250 transfer, 339 Reactivity, 50, 337 Receptor(s), 60, 80-82, 93-96, 101-103, 107, 134, 135, 139-141, 143, 217, 220, 220, 231, 235, 340, 341, 355, 139-141, 143, 217, 220, 220, 231, 235, 340, 341, 355, 359 Reduction, 54, 82, 84, 90, 91, 109, 110, 264 Reduction, 54, 82, 84, 90, 91, 109, 110, 264 Reduction, 54, 82, 84, 90, 91, 109, 110, 264 Regeneration, 13, 92, 132, 170-173, 177, 207, 225, 230, 240, 242, 245-248, 250, 251, 351, 352 Resultation(s), 12, 25, 331, 334,		Racemic, 45-48
149, 151, 154, 155, 158, 161-163, 168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 218-221, 225, 235, 237, 243, 244, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 230, 238, 241 Protoplast(s), 78, 79, 162, 167, 202, 230, 230, 236, 216		
168, 169, 172, 174, 175, 177, 179-181, 198, 203, 204, 214, 218-221, 225, 235, 237, 243, 244, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Protonos), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonomotive force, 84, 106 Protonophore(s), 108, 346 Protonplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 236, 67, 216		
179-181, 198, 203, 204, 214, 218-221, 225, 235, 237, 243, 244, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonphore(s), 108, 346 Protonophore(s), 108, 346 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 363, 377, 384, -diffusion, 247, 249, 250 transfer, 339 Reactivity, 50, 337 Receptor(s), 60, 80-82, 93-96, 101-103, 107, 134, 135, 139-141, 143, 217, 220, 221, 235, 340, 341, 354, 355 Recognition sequence, 336 system(s), 51, 57, 58, 79, 81, 96, 162 Red light, 89, 142, 174 Redox, 17, 61, 68, 74, 82, 143, 166, 186, 223, 264, 327, 339 Reduction, 54, 82, 84, 90, 91, 109, 110, 264 Regeneration, 13, 92, 132, 170-173, 177, 207, 225, 230, 240, 242, 245-248, 250, 251, 351, 352 Regulation, 51, 54, 86, 91, 114, 148, 157, 162, 184, 188, 189, 203, 350 Relativity, 17 Relaxation, 24, 42, 111, 133, 136, 340 Replication, 52, 53, 74 Reproduction, 172, 197-200, 238, 241 Resistance, 50, 67, 216		
218-221, 225, 235, 237, 243, 244, 256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonphore(s), 108, 346 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 230, 236, 237, 248, 250, 251, 351, 352 Recenter(s), 110, 111, 343 -diffusion, 247, 249, 250 transfer, 339 Reactivity, 50, 337 Receptor(s), 60, 80-82, 93-96, 101-103, 107, 221, 235, 340, 341, 354, 355 Recognition sequence, 336 system(s), 51, 57, 58, 79, 81, 96, 162 Red light, 89, 142, 174 Redox, 17, 61, 68, 74, 82, 143, 166, 186, 223, 264, 327, 339 Reduction, 54, 82, 84, 90, 91, 109, 110, 264 Regeneration, 13, 92, 132, 170-173, 177, 207, 225, 230, 240, 242, 245-248, 250, 251, 351, 352 Recuptor(s), 60, 80-82, 93-96, 101-103, 107, 221, 235, 340, 341, 354, 355 Recognition sequence, 336 system(s), 51, 57, 58, 79, 81, 96, 162 Red light, 89, 142, 174 Redox, 17, 61, 68, 74, 82, 143, 166, 186, 220, 230, 246, 327, 339 Reduction, 54, 82, 84, 90, 91, 109, 110, 264 Regeneration, 13, 92, 132, 170-173, 177, 207, 225, 230, 240, 242, 245-248, 250, 251, 351, 352 Regulation, 51, 54, 86, 91, 114, 148, 157, Relaxition, 24, 42, 111, 133, 136, 340 Replication, 52, 53, 74 Reproduction, 172, 197-200, 238, 241 Reproduction, 172, 197-200, 238, 241 Resistance, 50, 67, 216		
256-261, 264, 269, 333, 336, 337, 339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonphore(s), 108, 346 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 240, 242, 245, 241 Protoplast(s), 78, 79, 162, 167, 202, 230, 240, 242, 245, 241 Protoplast(s), 78, 79, 162, 167, 202, 230, 240, 242, 245, 241 Resistance, 50, 67, 216		
339, 342-346, 350, 353 Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38, 41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonphore(s), 108, 346 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, Protoplast(s), 78, 79, 162, 167, 202, 230, Reactivity, 50, 337 Receptor(s), 60, 80-82, 93-96, 101-103, 107, Recptor(s), 60, 80-82, 93-96, 101-102, 107, Recptor(s), 60, 80-82, 93-96, 101-103, 107, Recptor(s), 60, 80-82, 93		
Proteoglycans, 78, 99, 219 Prothallia, 173, 174, 209 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38,		
Prothallia, 173, 174, 209 Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38,		
Proton(s), 13, 17-22, 27, 29, 30, 35, 37, 38,		
41, 44, 60, 65, 82, 83, 87, 102, 106, 108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonophore(s), 108, 346 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 402, 242, 238, 241 Protoplast(s), 78, 79, 162, 167, 202, 230, 403, 341, 354, 355 Recognition sequence, 336 system(s), 51, 57, 58, 79, 81, 96, 162 Red light, 89, 142, 174 Redox, 17, 61, 68, 74, 82, 143, 166, 186, 223, 264, 327, 339 Reduction, 54, 82, 84, 90, 91, 109, 110, 264 Regeneration, 13, 92, 132, 170-173, 177, 207, 225, 230, 240, 242, 245-248, 250, 251, 351, 352 Regulation, 51, 54, 86, 91, 114, 148, 157, 162, 184, 188, 189, 203, 350 Relativity, 17 Relaxation, 24, 42, 111, 133, 136, 340 Replication, 52, 53, 74 Reproduction, 172, 197-200, 238, 241 Resistance, 50, 67, 216		
108-110, 132, 140, 142, 143, 167, 168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 protonema, 172, 173, 208-210 Protonophore(s), 108, 346 Protonophore(s), 108, 346 Protonation, 50, 154 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, Resistance, 50, 67, 216 Recognition sequence, 336 system(s), 51, 57, 58, 79, 81, 96, 162 Red light, 89, 142, 174 Redox, 17, 61, 68, 74, 82, 143, 166, 186, 223, 264, 327, 339 Reduction, 54, 82, 84, 90, 91, 109, 110, 264 Regeneration, 13, 92, 132, 170-173, 177, 207, 225, 230, 240, 242, 245-248, 250, 251, 351, 352 Regulation, 51, 54, 86, 91, 114, 148, 157, 162, 184, 188, 189, 203, 350 Relativity, 17 Relaxation, 24, 42, 111, 133, 136, 340 Replication, 52, 53, 74 Reproduction, 172, 197-200, 238, 241 Resistance, 50, 67, 216		the control of the co
168, 175, 176, 186, 220, 230, 235, 271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonophore(s), 108, 346 Protonophore(s), 108, 346 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 341 sink(s), 78, 79, 162, 167, 202, 230, 324, 242, 245-248, 250, 251, 351, 352 Requence, 336 system(s), 51, 57, 58, 79, 81, 96, 162 Red light, 89, 142, 174 Redox, 17, 61, 68, 74, 82, 143, 166, 186, 223, 264, 327, 339 Reduction, 54, 82, 84, 90, 91, 109, 110, 264 Regeneration, 13, 92, 132, 170-173, 177, 207, 225, 230, 240, 242, 245-248, 250, 251, 351, 352 Regulation, 51, 54, 86, 91, 114, 148, 157, 162, 184, 188, 189, 203, 350 Relativity, 17 Relaxation, 24, 42, 111, 133, 136, 340 Replication, 52, 53, 74 Reproduction, 172, 197-200, 238, 241 Resistance, 50, 67, 216		
271, 325, 333, 338, 339, 341, 343, 349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonophore(s), 108, 346 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 341, 349 system(s), 51, 57, 58, 79, 81, 96, 162 Red light, 89, 142, 174 Redox, 17, 61, 68, 74, 82, 143, 166, 186, 223, 264, 327, 339 Reduction, 54, 82, 84, 90, 91, 109, 110, 264 Regeneration, 13, 92, 132, 170-173, 177, 207, 225, 230, 240, 242, 245-248, 250, 251, 351, 352 Regulation, 51, 54, 86, 91, 114, 148, 157, 162, 184, 188, 189, 203, 350 Relativity, 17 Relaxation, 24, 42, 111, 133, 136, 340 Reproduction, 52, 53, 74 Reproduction, 172, 197-200, 238, 241 Resistance, 50, 67, 216		
349 flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonphore(s), 108, 346 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, Red light, 89, 142, 174 Redox, 17, 61, 68, 74, 82, 143, 166, 186, 223, 264, 327, 339 Reduction, 54, 82, 84, 90, 91, 109, 110, 264 Regeneration, 13, 92, 132, 170-173, 177, 207, 225, 230, 240, 242, 245-248, 250, 251, 351, 352 Regulation, 51, 54, 86, 91, 114, 148, 157, 162, 184, 188, 189, 203, 350 Relativity, 17 Relaxation, 24, 42, 111, 133, 136, 340 Replication, 52, 53, 74 Reproduction, 172, 197-200, 238, 241 Resistance, 50, 67, 216		TO BE CHARLES STREET PROPERTY WITHOUT NEEDED THE TOTAL TO PROPERTY
flow, 22, 176, 339 gradient(s), 82, 83, 102, 108, 235, 237, 335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonphore(s), 108, 346 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, Redox, 17, 61, 68, 74, 82, 143, 166, 186, 223, 264, 327, 339 Reduction, 54, 82, 84, 90, 91, 109, 110, 264 Regeneration, 13, 92, 132, 170-173, 177, 207, 225, 230, 240, 242, 245-248, 250, 251, 351, 352 Regulation, 51, 54, 86, 91, 114, 148, 157, 162, 184, 188, 189, 203, 350 Relativity, 17 Relaxation, 24, 42, 111, 133, 136, 340 Replication, 52, 53, 74 Reproduction, 172, 197-200, 238, 241 Resistance, 50, 67, 216		
gradient(s), 82, 83, 102, 108, 235, 237,		
335, 339 motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonophore(s), 108, 346 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, Reduction, 54, 82, 84, 90, 91, 109, 110, 264 Regeneration, 13, 92, 132, 170-173, 177, 207, 225, 230, 240, 242, 245-248, 250, 251, 351, 352 Regulation, 51, 54, 86, 91, 114, 148, 157, 162, 184, 188, 189, 203, 350 Relativity, 17 Relaxation, 24, 42, 111, 133, 136, 340 Reproduction, 52, 53, 74 Reproduction, 172, 197-200, 238, 241 Resistance, 50, 67, 216		
motor, 167 pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonophore(s), 108, 346 Protonation, 50, 154 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, Regeneration, 13, 92, 132, 170-173, 177, 207, 225, 230, 240, 242, 245-248, 250, 251, 351, 352 Regulation, 51, 54, 86, 91, 114, 148, 157, 162, 184, 188, 189, 203, 350 Relativity, 17 Relaxation, 24, 42, 111, 133, 136, 340 Reproduction, 52, 53, 74 Reproduction, 172, 197-200, 238, 241 Resistance, 50, 67, 216		
pump(s), 82, 102, 186, 220, 341 sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonmotive force, 84, 106 Protonophore(s), 108, 346 Protonation, 50, 154 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, 207, 225, 230, 240, 242, 245-248, 250, 251, 351, 352 Regulation, 51, 54, 86, 91, 114, 148, 157, 162, 184, 188, 189, 203, 350 Relativity, 17 Relaxation, 24, 42, 111, 133, 136, 340 Replication, 52, 53, 74 Reproduction, 172, 197-200, 238, 241 Resistance, 50, 67, 216	· · · · · · · · · · · · · · · · · · ·	
sink(s), 167, 168, 185 Protonema, 172, 173, 208-210 Protonmotive force, 84, 106 Protonophore(s), 108, 346 Protonation, 50, 154 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, Protonema, 172, 173, 208-210 Regulation, 51, 54, 86, 91, 114, 148, 157, 162, 184, 188, 189, 203, 350 Relativity, 17 Relaxation, 24, 42, 111, 133, 136, 340 Replication, 52, 53, 74 Reproduction, 172, 197-200, 238, 241 Resistance, 50, 67, 216		
Protonema, 172, 173, 208-210 Protonmotive force, 84, 106 Protonophore(s), 108, 346 Protonation, 50, 154 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, Regulation, 51, 54, 86, 91, 114, 148, 157, 162, 184, 188, 189, 203, 350 Relativity, 17 Relaxation, 24, 42, 111, 133, 136, 340 Replication, 52, 53, 74 Reproduction, 172, 197-200, 238, 241 Resistance, 50, 67, 216		20 At 10 At
Protonmotive force, 84, 106 Protonophore(s), 108, 346 Protonation, 50, 154 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, Resistance, 50, 67, 216		
Protonophore(s), 108, 346 Protonation, 50, 154 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, Relativity, 17 Relaxation, 24, 42, 111, 133, 136, 340 Replication, 52, 53, 74 Reproduction, 172, 197-200, 238, 241 Resistance, 50, 67, 216		
Protonation, 50, 154 Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, Relaxation, 24, 42, 111, 133, 136, 340 Replication, 52, 53, 74 Reproduction, 172, 197-200, 238, 241 Resistance, 50, 67, 216		
Protoplasm, 13, 34, 72, 129, 133, 159, 165, 194, 205, 211 Protoplast(s), 78, 79, 162, 167, 202, 230, Replication, 52, 53, 74 Reproduction, 172, 197-200, 238, 241 Resistance, 50, 67, 216		I select and the select and the selection of the selectio
194, 205, 211 Reproduction, 172, 197-200, 238, 241 Protoplast(s), 78, 79, 162, 167, 202, 230, Resistance, 50, 67, 216		
Protoplast(s), 78, 79, 162, 167, 202, 230, Resistance, 50, 67, 216		
		Maria Mail II Maria Mari
	237	Respiration, 82, 103, 106, 166, 187, 188,

190, 198, 200, 215	Septation, 158, 181-183, 190, 191, 194, 195,
Retina cells, 104, 181, 236	202, 205,
Retinal, 89, 341	Sequence(s), 15, 50-58, 81, 86, 87, 94, 96,
Retinoic acid(s), 240, 244	97, 107, 121, 137, 157, 165, 184, 196,
Rhizoid(s), 147, 158-160, 164, 168-173,	204, 220, 226, 231, 256, 258, 336
181, 201, 204-210, 352, 353	Serine, 40, 87, 141, 259, 337
elongation, 172	Severin, 114, 343
outgrowth, 205, 206, 352	Sex (differentiation-disjunction), 153, 174,
Rhodamine, 152, 153, 155, 156, 174, 188	197-200, 358
Rhodopsin, 81, 89, 141	Shoot, 172, 185, 209, 229, 231-235, 237,
Ribonucleic acid (RNA), 46, 52, 56-58, 70,	353
96, 124, 161, 166, 198, 199, 208, 214,	apex, 225, 229, 231, 232, 238, 239
257, 336, 345	axis, 224, 225, 230
polymerase, 52, 57	meristem, 231
viruses, 57	Signal(s), 14, 53, 70, 81, 82, 87, 93, 95-97,
gradient, 198, 199	100, 101, 106, 107, 118, 138, 140,
Ribosome(s), 58, 72, 83, 96, 97, 124, 172,	142, 143, 145, 148, 179, 182, 203,
198, 199, 205	209, 230, 231, 236, 237, 239, 240,
nuclear cap, 198	243, 252, 253, 256, 265, 266, 342,
Root(s), 147, 163, 181, 185, 209, 210,	353-355
229-234, 236, 237, 268, 354, 355	inhibition, 185
apex, 225, 229, 231	intracellular, 102
cap, 176, 236, 355	morphogens, 240
hair(s), 147, 164, 169, 175, 176, 186,	peptide(s), 82, 88, 97, 106
205, 210, 211	receptor-recognition, 96, 97
meristem, 231	transduction, 79, 350
-shoot axis, 224, 225, 230	target protein, 103, 353
Rotation, 37, 44, 45, 52, 51, 64, 69, 139,	Silicon (silica), 22, 108, 184, 330
140, 245, 265, 268, 331, 333	Simulation, 222
Ruthenium red, 106	Sink(s), 92, 157, 167, 168, 185, 222, 233,
	240
Sarcomere(s), 116-118, 344	Size(s), 20, 27, 34, 35, 40, 81, 93, 117, 137,
Sarcoplasmic reticulum, 89, 340	148, 149, 157, 161, 170, 173, 177,
Scaffolding, 70, 112	182, 185, 189, 198, 200, 225, 228,
Scale(s), 18, 46, 82, 99, 100, 244, 327	229, 240, 247, 250, 262, 331
Scattering, 23, 29-31, 124, 227, 343	Sliding, 76, 117, 120, 122, 125, 175, 345
Sclerotium, 226	Sodium (see also Na+), 23, 49, 63, 68, 76,
Secretion, 86, 94, 98, 99, 101, 103, 104, 132,	90-93, 333, 341
137, 149, 150, 152, 177, 205, 207,	Solar, 45, 48, 108, 109, 327, 330
219, 220, 342, 343, 353	Solitons, 51, 52, 68
Segmentation, 169, 224, 259, 260	Solvation, 333
genes, 256, 259, 260	Sorocarp(s), 241, 242
Segregation, 189, 200, 218	Sorting, 96-98, 100, 102, 103, 161, 177, 219,
Self-assembly, 70, 189, 222, 357	220, 342, 343, 353
Self-electrophoresis, 175	-out, 154, 198
Semiconductor(s), 22, 68, 108, 329	Source(s), 22, 30, 45, 47, 48, 61, 92, 107,
Senescence, 238	132, 140-142, 173, 222, 233, 240,
Sensor(s), 141, 340, 341, 354, 355	206, 209, 328, 350

Spatial, 14, 15, 85, 95, 110, 121, 140-142,	Stimulus (Stimuli), 22, 86, 95, 139-141,
163, 180, 190, 203, 209, 214, 221,	143, 145, 206, 210, 220, 221, 223,
223, 259, 265	239, 243, 253, 329, 341, 355
asymmetry-symmetry, 69, 271	transducers, 236
organization, 112, 239, 259, 348	Stomata, 205, 211
pattern, 207, 240, 252	Stress, 115, 126, 147, 149, 182. 186
polarity, 214, 257, 259	Structural proteins, 58, 73
segregation, 183	Subapical, 155, 165, 168, 169, 173, 175,
Spatio-temporal, 73, 325	235, 349
Spectra-spectrum, 30, 31, 61, 86, 89, 108,	zone(s), 163, 164
331-333	Substitution, 54
Spectrin, 112, 114, 221	Sucrose, 206, 207, 222
Sperm, 89, 254, 261-263	Sugar(s), 44-46, 60, 64, 73, 79, 165, 233,
cells, 161, 218	334
Spherical, 34, 40, 63, 68, 69, 108, 131, 147,	Sulfur (SH ₂ , SO ₂), 38, 84
149-154, 158, 159, 161-163, 187,	Superconductivity, 23-25
204, 229, 264, 325, 330, 348	Superoxide dismutase (SOD), 61, 66
bud(s), 149, 157	Surface
growth, 157	charges, 237
stage, 154, 325	membrane(s), 73, 81, 90, 91, 101, 177,
Spin(s) 17, 22, 25, 27-30, 43, 44, 140, 331,	333, 337-339
333	potential(s), 230, 340
Spindle, 120, 124-128, 345	proteins, 77, 332, 337, 347
pole bodies, 151-153	structures, 203
Spiral, 45, 245, 268	Symmetry, 13, 18, 19, 44, 45, 47, 51, 64,
Spitzenkörper, 105, 164, 166	68-70, 72, 79, 123, 147, 154, 158,
Sporangia, 191, 194, 195, 352	159, 182, 221, 226, 246, 262, 265,
Sporangiospore(s)-phores, 157, 163, 352,	327-330, 334, 336
354	bilateral, 328
Spore(s) 13, 155, 160, 169, 172, 173, 181,	mutant, 268
187, 190, 191, 194-196, 202-205,	pattern, 255, 269, 330
208-210, 224, 228, 241, 325, 348, 349	radial, 226, 262
differentiation, 202	spatial, 69
germination, 155, 173, 208, 348	Symport, 82, 167, 168
induction, 195	Synapse(s), 87, 93-95, 105, 122, 178, 180,
unpolarized, 205	181
Sporulation	polarity, 177, 178
endo-, 190-192, 202	transmission, 94, 95, 180
exo-, 192-195, 352	vesicle(s), 90, 94, 177, 178
Spreading, 145, 199, 350	Synapsin, 94
Stability, 13, 18, 22, 69, 127, 132, 183, 229	Synthetic, 23, 52, 74, 77, 93, 109, 110, 166,
Stars, 25	185, 198, 235
	enzymes, 183
Statocyte(s), 236, 354, 355	polymers, 67
Statolith(s), 168, 236, 355	
Stereospecificity, 15, 44-46, 86	Target(s), 30, 53, 81, 89, 96, 97, 99, 106,
Steroids, 63, 78	122, 157, 162, 177, 178, 180, 181,
Stimulation, 90, 92, 141, 148, 227, 235	219, 230, 349, 350

Targeting, 81, 97, 101-103, 107, 219	Transcellular, 104, 137, 154, 168
signal(s), 97, 103, 106, 353	calcium, 106
Tartrate, 44	electrical potential, 170
Taxis (Chemo-, Photo-), 139-144	ion current(s), 154, 161, 167, 349, 353
Taxol, 121	Transduction, 140, 142, 143, 236, 338, 348,
Temperature, 17, 24, 25, 63, 68, 76, 80,	355, 358
129, 154, 159, 161, 165, 175, 182,	Transformation, 80, 133, 137, 138, 139,
187, 193, 226, 267, 268, 356	157, 162
Template(s), 35, 52, 53, 55, 57, 58, 69, 122,	Translation(s), 44, 51, 53-55, 58, 60, 69, 70,
132, 243	96, 97, 132, 196, 214
Temporal, 15, 95, 133, 140, 141, 143, 163,	Translocation, 51, 52, 54, 82, 83, 85, 96, 97,
180, 190, 203, 251	106, 107, 123, 132, 169, 203, 204,
Tetrahedral, 39, 64, 65, 332	236, 239, 333
Tetrapolarity, 196, 340	Transmembrane, 77, 81, 84, 86, 87, 90, 94,
Thalli, 181, 204-210, 352	102, 110, 132, 136, 140, 340, 341,
Theory, 14, 17, 20, 24-27, 31, 92, 131, 134,	346, 353
223, 255, 327, 355	domains, 261
metabolic gradients, 247, 255	potential(s), 89, 107, 339
Thermodynamic(s), 14, 75, 128, 272, 337,	receptor protein(s), 42, 77, 86, 90, 112
341	Transmission, 32, 86, 93-95, 105, 107, 343
Thiol, 40, 162	Transport(s), 24, 60, 62, 78, 79, 82, 83, 85,
Thylakoid(s), 74, 79, 108-110, 143, 343	86, 89, 94, 97-106, 112, 122, 140,
Time, 17, 18, 30, 37, 51, 70, 72, 87, 92, 94,	143, 152, 163, 177-179, 191, 207,
96, 98, 102, 112, 124, 125, 139-141,	219, 220, 231, 233, 235, 338, 341,
157, 159, 164, 175, 179, 181, 182,	342, 343, 355
190, 195, 200, 206, 207, 209, 210,	Transposons, 55, 336
214, 215, 218, 226, 231, 234, 240,	Treadmilling, 113, 114, 118, 120, 125
244, 253, 257, 269, 270, 273, 330,	Tree, 147
341	Triaxial patterns, 267-270, 356
Tip(s), 114, 115, 131, 144, 164, 167-169,	Trichoblasts, 205, 211
173, 175, 176, 193, 195, 233, 349	Trichome, 107, 133, 143, 211
acidic, 167, 168	Trigger(s), 90, 93, 203, 215, 233, 239, 243
growth, 150, 162, 163, 165, 167, 168,	Triplet(s), 54, 57
169, 352	Tripolar, 326, 329
organelles, 167, 173, 178	Trophoblast, 266
Tissue(s), 62, 130, 137, 147, 169, 177, 186,	Tropisms (tropic curvatures), 181, 235-238,
189, 201, 208, 224-228, 230-233,	354-355
237-240, 247-250	Tropomyosin, 114
polarity, 230-234, 261, 262	Tubular, 70, 99, 104, 163, 252, 340, 341
regeneration, 241, 242, 248	Tubulin(s), 15, 81, 112, 114, 118-121, 125,
Titanate, 23, 330	128, 129, 139, 151, 153, 163, 174,
	177, 180, 187, 194, 200, 218, 325,
Topology, 26, 50, 83, 102, 162, 174, 191,	344, 349, 358
358 Torque(s) 26, 27, 140	20 Perc 1976 W Hardware wheel o windows
Torque(s), 36, 37, 140	α - and β -, 111, 119, 121, 152
Trace, 108 Traffic (introcallular), 07, 100, 122, 148	gene(s), 151, 187
Traffic (intracellular), 97-100, 122, 148,	membrane, 187
177, 256, 343 Tranducers, 100	mutations, 193
Trans. 52, 97, 101, 102, 260	Tumoral cells, 115, 149, 160
Trans, 52, 97-101, 103, 260	Tunnelling, 35, 38, 42, 67, 272, 336

Turgor pressure, 147, 162, 166, 186, 187, 230 Uncoupling agents, 108 Unidirectional, 15, 90, 97, 107, 111, 115, 119, 120, 132, 136, 182, 345 Unification, 20, 26, 27, 45 Unipolar, 147, 183, 240 Unity, 20 Universe, 17-19, 26, 27, 45, 271, 327, 328 Unpolarized, 28, 62, 205, 212, 329 Uptake, 94, 102, 106, 110, 340 Urea, 115	Vibrating probe (electrode), 154, 168, 194, 207, 230 Villin, 114 Vimentin, 111 Vinculin, 144 Virus(es), 57, 69-71, 117, 219 DNA, 58, 337 form (TMV), 69, 70 RNA, 57 Vision, 89 Vitamin B ₁ , 160 Vitellogenesis, 100 Voltage gradient, 167
Urea, 115 UV-irradiation, 214, 215, 257, 258 Vacuole(s), 129, 130, 137, 155, 164, 173, 184, 185, 194, 195, 225, 244 Valence, 22, 23, 24, 68, 328 Valinomycin, 107, 108 Vascular systems, 224, 225, 229, 231-233, 237 Vector(s), 26-29, 33, 44, 249, 336, 350, 352 Vectorial, 79, 82, 85, 89, 124, 142, 151, 166, 219, 220 movement, 102, 200, 353 mitochondrion, 106 Vegetal, 212, 214-216, 253, 255 cortex, 214, 261 pole, 190, 212, 253-255, 261, 263, 264, 269 region, 217, 252, 255, 262 Vegetative, 147, 164, 165, 169, 190, 194-197, 203, 208, 226, 229, 231, 238, 239, 241, 352 buds, 239 cell(s), 161, 191, 192, 202, 205, 228 differentiation, 193 growth, 238, 241 pole, 161	Wall(s), 15, 100, 129, 133, 134, 148, 150-153, 155, 157-159, 161, 162, 165-168, 175, 176, 182, 183, 185, 186, 193-195, 199, 206, 207, 209-212, 224, 225, 230, 236, 237, 256, 267, 341 apical, 164, 166 cell, 64, 79, 149-155, 157, 162, 166, 169, 175, 176, 183-187, 189, 201, 203, 206, 207, 211, 222, 223, 226, 230, 235, 352, 356 vesicles, 167 Water (see also H ₂ O), 17, 34-41, 51, 55, 58-68, 73, 75-77, 104, 110, 155, 159, 169, 175, 228, 233, 246, 268, 325, 327, 332-334, 338, 343, 352 cell, 331 dipolar, 34-38, 332, 333, 338 surface-interfaces, 63, 76, 333, 338 Wing, 240, 260, 261, 265 Wound, 176, 250, 251, 341 X-ray(s), 35, 37, 78, 86, 110 Xylem, 233
Ventral, 231, 256, 263 Vesicle(s), 74, 85, 94, 95, 97-102, 105, 110, 122-124, 137, 151, 153, 155, 157, 161-164, 166-168, 173, 175, 177-180, 183, 184, 194, 195, 204, 206, 207, 215, 235, 340, 342-345 membrane, 102, 109, 342 secretion (granules), 101, 207 transport, 151	Yeast(s) budding(s), 150, 154, 184, 187, 344, 348 cell cycle, 151-154 cell polarity, 152, 161 cytoskeletal network, 342, 344, 348 elongation (fission-), 183, 184 forms, 161-163 mating, 152-154, 195-197 Yellow egg crescent, 216, 261

Yin-Yang, 13, 138, 148, 271 Yolk, 190, 212, 216-218, 269, 270

Zinc (Zn²⁺), 58, 59, 61, 337 Zone(s) (cellular zonation), 104, 117, 129, 133, 134, 161, 165, 168, 171, 173-178, 185, 195, 199, 214, 230, 236, 237, 245, 248, 256, 262, 350, 352, 356 elongation, 230 exclusion (mitochondria), 103-105, 166-168, 193

Zoospore(s), 134, 169, 195, 204

Zwitterion, 41

Zygospore, 169

Zygote(s), 151, 158-160, 169-171, 184, 195, 196, 201, 204-207, 210, 213, 214, 224, 225, 252

differentiation, 14

germination, 205

TAXONOMIC INDEX

Acanthamoeba, 116, 344, 346	Bacillus, 167, 202
Acetabularia, 170, 171	B. brevis, 86
A. mediterranea, 208	B. subtilis, 182, 191, 267, 356
A. Wettsteinii, 208	Bacteria, 41, 53, 55, 60, 103, 110, 130-132,
Achlya, 194	139, 140, 149, 150, 164, 182, 190,
A. bisexualis, 165,	201, 267, 356
A. debaryana, 194	budding, 149, 150
Actinomycetes, 164	helical, 267, 356
Agaricus, 226, 227	photosynthetic, 110
A. bisporus, 225, 226	Bacteriophages (fd, l), 53, 69-71, 337
Agrobacterium tumefaciens, 164	Basidiomycetes, 225, 226, 228
Algae, 73, 130, 132, 147, 168, 181, 188, 228,	Bipolaris, 184
229, 340, 341, 344, 346, 352, 353	Birds (chick), 81, 121, 146, 181, 217, 265,
Allium, 211	266
Allomyces, 147, 166, 181, 192, 194, 195,	Blastocaulis, 150
197, 199, 200, 204, 205, 349, 358	Blastocladiella emersonii, 194
A. arbuscula, 194, 197-199	Blepharisma, 268
A. javanicus (hybrid), 199	Boltenia, 216, 254
A. macrogynus, 197, 199	B. villosa, 261
Amblystoma mexicanum, 262	Botrydium, 169
Amoebae, 133, 134, 136, 138-141, 243, 326	Botrytis cinerea, 187, 334
Amoeba proteus, 134, 136, 143	Bollytis Cinerea, 187, 334
Amphibians, 204, 212, 216, 217, 223, 238,	Canarhabditis alagans 213 251 252
262-264, 270, 272, 355	Caenorhabditis elegans, 213, 251, 252 Callitriche, 211
Anabaena, 192	Candida albicans, 162, 163, 348
Anise, 349	Capsella bursa-pastoris, 225
Anthoceros, 225	Caulerpa, 344
Aquaspirilla, 267 Arabidopsis thaliana, 355	Caulobacteria, 201-203
	Caulobacter crescentus, 202, 203
Arthropade 355	Cecidomyids, 214
Arthropods, 355	Chaetophorales, 169
Ascaris, 252	Charges 168 338
Ascidians, 215, 216, 254, 261, 262, 272	Characeae, 168, 338
Ascobolus, 56	Chara, 130, 163, 168, 169, 205
A. immersus, 56	Chick, 266
Ascomycetes, 155, 196	Chironomids, 215
Aspergilli, 193	Chlamydomonas, 119, 121, 132, 143, 228,
A. nidulans, 193, 345	346
A. aureolatus, 193	Chlorococcales, 229
Azolla, 268	Chlorophyceae (Green algae), 119, 130,

143, 147, 169, 170, 185, 188, 190, 208, 228, 340, 341, 342, 344, 346 Ciliates, 244 Cladophorales, 147 Cladophora glomerata, 207 Clavelina, 262 Coleoptera, 214 Chrysomelid, 214 Coelenterates, 246, 250 Conjugatophyceae, 340 Coprinus, 225, 226 C. cinereus, 226, 352 C. congregatus, 226 C. stercorarius, 225 Corn (see also Zea), 354 Corynebacterium, 164 Cryptogams (see also ferns and mosses), 181, 208 Cyanobacteria (Blue-green algae), 111, 132, 133, 142, 143, 190-192 Dentalium, 253, 269 Desmid(s), 147, 188 Diatom(s), 132, 147, 185 Cymbella, 184 Navicula, 184 Dictyostelium, 116, 118, 134, 242, 243, 346,

Cymbella, 184
Navicula, 184
Dictyostelium, 116, 118, 134, 242, 243, 346, 347
D. discoideum, 124, 133, 140, 141, 143, 241-244, 347
Dictyota, 205
Diptera, 214, 257
Drosophila, 13, 62, 124, 213, 214, 216, 217, 256-261, 344, 356
D. melanogaster, 259, 261
Dryopteris, 160, 169, 174
D. filix-mas, 174

Echinoderms (sea-urchin), 89, 120, 126, 212, 253, 254, 264, 272

Enteromorpha, 208

Eremosphaera viridis, 341

Equisetum, 160, 205, 209, 210, 224

Escherichia, 88

E. coli, 52, 55, 64, 82, 140, 182

Euglena gracilis, 143

Eukaryotes, 107, 151

Ferns, 147, 160, 173, 174, 189, 190, 210, 232, 238, 268

Fishes, 147, 217, 265 shark, 265 Flagellate(s), 99 Flammulina (Collybia) velutipes, 226 Flexibacter, 132 Fomes, 225 Foraminifera, 268 Formica, 214 Frog, 263 Fucales, 201, 352 Fucoid algae, 182 Fucus, 81, 157-159, 204, 206, 208, 210, 224 F. furcatus, 158 Funaria, 160, 163, 164, 173, 209 F. hygrometrica, 160, 172 Fungi, 56, 101, 129, 134, 147, 155, 163-165, 168, 184, 194-196, 225, 340, 341, 352 higher (mushrooms), 227, 354 imperfecti, 155

Globigerina, 268 Griffithsia, 172, 353 G. pacifica, 185

Halicystis, 91
Halobacterium halobium, 110, 142
Hansenula, 64
Human, 133, 266, 273, 334
Hydra, 246-250, 358
H. attenuata, 248
Hydractinia, 250
Hydroids, 170
Hydrozoa, 241, 246, 249, 272

Ilyanassa, 253 Insects, 217, 256-260, 270, 356 Iris, 326

Kloeckera, 187

Laminaria, 224
Leafhoppers (Cicadellids), 215
Lentil, 349
Lepidoptera, 214
Lilium grandiflorum, 161
Lilium longiflorum, 175
Liverworts, 173, 174, 209

Pasteuria, 149 Lymnaea, 269 Pea, 234 L. peregra, 268, 269, 356 Pelvetia, 207, 210 L. suturalis, 269 P. fastigiata, 207 Mammals, 212, 266, 356 Penicillia, 193 Marchantia polymorpha, 173, 209 P. claviforme, 193 Micrasterias, 147, 188 Phaeophyceae (Brown algae), 91, 159, 160, Molluscs, 212, 213, 252, 268, 269, 272, 356 169, 172, 195, 224, 352 Monilia, 185, 193 Phormidium, 133, 142, 143 M. fructigena, 105 P. uncinatum, 132 Monstera, 210 Phycomycetes, 204 Mosses, 147, 160, 172, 173, 188, 197, Phycomyces, 352, 354 208-211 P. blakesleeanus, 129 Mouse, 266, 356 Phyllanthes, 238 Mucorales, 162, 196 Physarum, 129, 138 Mucor racemosus, 157, 162 P. polycephalum, 133, 137, 143 M. rouxii, 162, 163, 348 Physcomitrella, 209 Mycetozoa, 241 Pinus sylvestris, 197 Mycobacteria, 164 Planaria maculata, 251 Mycobacterium, 164 Planctomyces, 149, 150 Mycoplasma, 132, 133 Pleurococcoid, 147 Myxobacteria, 132 Pleurochrysis, 99 Podospora anserina, 196 Nadsonioideae, 184 Polyporaceae, 225 Nadsonia, 187 Polyporus brumalis, 226, 227 Naegleria, 138, 139 Prokaryotes, 53, 54, 96, 129, 132, 133, 139, N. gruberi, 138 140, 142, 143, 149, 164, 182, 336 Nassarius, 253 Protists, 107 Neurospora, 118, 165, 196 Protosiphon, 169 N. crassa, 56, 102, 108, 129, 155, 156, Protozoa, 104, 132, 141, 244, 245, 268, 272 165, 167, 187, 193, 195, 349 N. sitophila, 195 Rat, 351 Nicotiana tabacum, 239 Reptiles, 217 Nitella, 116, 130, 185 Rhodnius, 176 N. axiliaris, 185 Rhodophyceae (Red algae), 111, 172, 185, Nitrobacter, 150 345, 353 Nocardia corallina, 164 Rhodopseudomonaceae, 110 Nostocaceae, 142 Rodents, 267 Oedogonium, 169 Sabella, 251 Saccharomyces cerevisiae, 150-153, 336, Oncopeltus, 176 345, 348 Onoclea sensibilis, 173 Oomycete(s), 166, 349 S. uvarum, 152 Saccharomycodes ludwigii, 184 Oscillatoriaceae, 142 Salmonella, 131 Osmonda, 160 Saprolegnia, 194 Paracentrotus lividus, 213 S. ferax, 349 Scenedesmaceae, 229 Paramecia, 244 Schizophyllum commune, 196, 226 Paramecium, 130, 141, 142, 244

Schizosaccharomyces pombe, 183 Selaginella, 231 Seliberia, 150 Serratia marcescens, 242 Sinapis alba, 239 Slime mold(s), 116, 118, 129, 133, 134, 137-140, 241, 242, 346, 347, 348 Smittia, 257 Snails, 356 Sphacelaria, 172 Sphagnum, 197, 205, 211 Spirillum, 131 Spirochaeta, 131 Spirogyra, 147, 169 Squid, 90, 91, 94, 122, 179 Stentor, 245, 246, 268 Streptomyces, 164 S. streptomycini, 164 Styela, 216, 261

Tetrahymena, 57, 244-246, 268 Torpedo californica, 341 Trichoderma viride, 155, 156, 340 Tubularia, 247 Tunicates, 215, 261

Uromyces appendiculatus, 349

Vaucheria, 147, 169, 195

Viruses, 57, 69, 219
Tobacco Mosaic Virus (TMV), 69
Volvocaceae, 228
Eudorina, 229
Gonium, 229
Pandorina, 229
Volvox, 229

Worms,

Annelids, 213, 249, 355 Nematodes, 213, 251, 252 Planarians, 250, 251 Polychaetes, 251 Sabellids, 268

Xanthophyceae (Yellow algae), 147, 169, 195 Xenopus, 217 X. laevis, 262

Yeast(s), 56, 64, 97, 101, 107, 149-153, 161, 162, 181, 184, 187, 196, 326, 345, 347, 348 fission, 183

Zea (see also corn), 231 Z. mays, 354 Zygnematales, 147 Zygomycetes, 196