Zeitschrift: Archives des sciences et compte rendu des séances de la Société

Herausgeber: Société de Physique et d'Histoire Naturelle de Genève

Band: 47 (1994)

Heft: 3: Archives des Sciences

Artikel: Proton transport driven by the plasma membrane ATPase from

Spinacea oleracea leaves : biochemical characterization

Autor: Bellamine, Jalil / Greppin, Hubert

DOI: https://doi.org/10.5169/seals-740192

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 25.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Archs Sci. Genève Vol. 47 Fasc. 3 pp. 287-296 Décembre 1994

Communication présentée à la séance du 16 juin 1994

PROTON TRANSPORT DRIVEN BY THE PLASMA MEMBRANE ATPASE FROM SPINACEA OLERACEA LEAVES. BIOCHEMICAL CHARACTERIZATION

BY

Jalil BELLAMINE* & Hubert GREPPIN*

ABSTRACT

Proton transport driven by the plasma membrane ATPase from Spinacia oleracea leaves. Biochemical characterization. – The plasma membrane of spinach leaves was purified by two phase partitioning and the H+pumping activity associated to the purified plasma membrane was characterized. The pH optimum was 6.7 similar to that of the ATPase activity and the substrat specificity was more obvious for ATP with a Km_{app} of 0.697. The H+pumping activity was more or less inhibited by different ATPase inhibitors, and especially by VO₄ with I_{50} of 160 μ M. The SDS-PAGE of the purified plasma membrane proteins revealed the presence of a polypeptide of about 100 kDa which cross-reacted with the polyclonal antibodies against the plasma membrane ATPase of Arabidopsis thaliana.

Key-words: Spinacia oleracea, Adenosine triphosphatase, H⁺pump.

Abbreviations: ADP, Adenosine diphosphate; ATP, Adenosine triphosphate; DCCD, Dicyclohexylcarbodiimide; DCMU, 3-(3,4-Dichlorophenyl)-1,1-dimethylurea; DES, Diethysilbestrol; GTP, Guanosine triphosphate; EB, Erythrosine B; PEP, Phosphoenolpyruvate; PP_i, Pyrophosphate. SDS, Sodium dodecylsulfate.

INTRODUCTION

Many important aspects of plant physiology, including growth, development, nutrient transport and stomata mouvements, seem to be controlled by the H+pumping ATPase of plasma membrane (PM) (Serrano,1989). This could be supported by the fact that the early event in the action of growth-modifying pathogens, hormones, and light is an alteration in the PM proton pumping activity (Assmann *et al.*, 1985; Cleland, 1987; Bidwai & Takemoto, 1987).

^{*} Laboratoire de Biochimie et Physiologie végétales, 3, pl. de l'Université, CH-1211 Genève 4

In vitro activation of the H+pump of the purified PM, by auxins has been demonstrated for tobacco (Barbier-Brygoo, 1989; Santoni et al., 1991), Petunia (François et al. 1992) and Spinach (Bellamine et al. 1993 a), suggesting that the pump is one of the earliest responses to the phytohormone (Cleland, 1987; Rayle & Cleland, 1992).

The PM H+ATPase of higher plants belongs to the P-type family of cation-translocating enzymes and generates an electric potentiel and pH gradiant (H+motive force) that drive solute uptake across the PM. That type of enzyme is sensitive to vanadate (Jacobs & Taiz, 1980) and forms a phosphorylated intermediate (Briskin & Hanson, 1992). On the basis of the polypeptide composition and sensitivity to inhibitors, the plant PM H+ATPase is readily distinguished from that found in membrane derived from chloroplast, mitochondria and vacuole (Pederson & Carafoli, 1987). The plant PM H+ATPase was well characterized by biochemical and electrophysiological techniques, and the purified enzyme contains a single polypeptide of about 100 Kda (Grouzis *et al.*, 1990; Becker *et al.*, 1993). Based on molecular approach, this enzyme presents many transmembrane segments (Harper *et al.* 1989).

The genes of different bacterial, fungal, animal and plant ATPases of this family have already been cloned and sequenced (Serrano, 1988; Harper *et al.* 1989). The predicted amino acid sequence of *Arabidopsis thaliana* H+ATPase is more closely related to fungal and protozoan H+ATPase than bacterial K+ATPases or to animal (Na+/K+)- (H+/K+)- and Ca²⁺ ATPases (Pardo & Serrano, 1989).

In this study we reported a biochemical Characterization of the PM H+ATPase from spinach leaves. The plasma membrane was purified by phase partitioning and characterized as in a previous work (Bellamine *et al.*, 1993 a).

MATERIALS AND METHODS

Plant Material

Spinach (*Spinacia oleracea*, cv.Nobel) plants at the vegetative stage were grown in a phytotron for 4 weeks under non inductive short days (SD) of 8 h light (8:00 am to 4:00 pm; 400 µmol m⁻² s⁻¹) and 16 h dark (4:00 pm to 8:00 am). The temperature was set at $20 \pm 1^{\circ}$ C and the relative humidity at 80 ± 5 % during the light period and 60 ± 5 % during the darkness.

Preparation of microsomal vesicles

Crude microsomal vesicles preparation and PM purification were as described by Bellamine *et al.* (1993 a).

ATPase activity

The plasmalemma ATPase activity was measured at 37°C as described by Bellamine *et al.* (1993 b).

Proton pumping activity

The initial rate of quinacrine fluorescence quenching was utilized to measure the plasmalemma proton translocation activity as described by Bellamine *et al.* (1993 a).

Protein determination

Membrane aliquots were diluted 20 fold with cold water and centrifuged at 96,100 x g for 30 min at 5 °C. The pellet was resuspended in cold water (160 µl) and proteins were determined using Bio-Rad solution. 50 µl of protein suspension were diluted in 750 µl of water and 200 µl of Bio-Rad solution were added. The obtained solution was mixed and incubated at room temperature for at least 5 min. The OD at 595 nm was measured and proteins were quantified using BSA as the standard.

SDS PAGE and Western blot

Purified plasma membrane vesicles were washed in a solution containing 10 mM Tris, 20% glycerol, 1 mM EDTA Na₂, 1 mM PMSF, 50 mg chymostatin and adjusted to pH 6.7 with HCl. Equivalent to 90 µg of proteins were pelleted by centrifugation at 96,500 g for 30 min at 5°C (40,000 rpm, Beckmann T50 rotor). The resulting pellet was resuspended in a sample buffer, as described by Laemmli (1970), supplemented by 1 mM PMSF and 50 mg/ml Chymostatin. After the SDS polyacrylamide gel electrophoresis (SDS PAGE), the peptides were transferred to a nitrocellulose sheet as described by Towbin *et al.* (1979), and revealed using antibodies to the central part of *Arabidopsis thaliana* plasmalemma ATPase (gift from R. Serrano). Immunodetection in western blot was made with TBS as a basic medium. Immune serum was diluted to 1/1000 and the second antibody was diluted 1/2000 (Anti-Rabbit IgG (whole molecule) alkaline phosphatase conjugate provided from sigma). The phosphatase alkaline substrates were NBT and BCIP.

RESULTS AND DISCUSSION

pH effect on ATP dependent H+pumping activity and on ATP hydrolase activity

The H+pumping and the phosphohydrolyzing activities were tested as described in Materials and Methods according to the pH of the reaction medium (Fig, 1). Both activities showed a maximum at pH 6.7 described for other plant species (De Michelis & Spanswick, 1986; Olivari *et al.*, 1992).

Vanadate effect on H+pumping activity

The sodium orthovanadate, a poison well known to inhibit plant PM H+ATPase, inhibits also the H+pumping activity of spinach leaves PM. Figure 2 shows the kinetic of inhibition by vanadate of the initial rate of H+pumping measured as described in




Fig. 1.

pH effect of the reaction medium on the ATP hydrolase and H⁺pumping activities of the purified plasma membrane as described in Materials and Methods. The pH of the medium was adjusted by varying the proportions of BTP and Mes (50 mM final concentration).

Materials and Methods. The inhibition increases with vanadate concentration over the whole concentration range tested. The concentration of vanadate required to inhibit the H+ATPase activity by 50% (I_{50}) was about 160 μ M, higher than that found for the same enzyme in other plants (Rasi-Caldogno *et al.*, 1985; De Michelis & Spanswick, 1986; Olivari *et al.*, 1992). However, the I_{50} value of vanadate depends on the plant species and on the experimental conditions for the same plant specie (Rasi-Caldogno *et al.*, 1985; De Michelis & Spanswick, 1986).

Substrat specificity of the H+pumping activity

Table 1 shows that the H+pumping requires rather ATP as source of energy than other phosphorylated substrats tested. The apparent Michaelis-Menten constant (Km_{app}) for ATP was 0.697, similar to that obtained for the same enzyme in other plant species (Rasi-Caldogno *et al.*, 1985; De Michelis & Spanswick, 1986).




Fig. 2.

Effect of increased vanadate concentration on the ATP dependent H+pumping activity of the purified plasma membrane. The reaction was started by addition of 1.5 mM Mg ATP, after a 10 min preincubation at room temperature in presence of indicated concentration of vanadate.

TABLE 1.

Effect of some phosphorylated substrats on H⁺pumping activity of the purified plasma membrane as described in Materials and Methods. The reaction was started by addition of 1.5 mM of the substrat, and the activities were expressed in AU min⁻¹ mg⁻¹ protein ± standard deviation. All the substrats were sodium salt except the PEP which was tricycloammonium salt. The reaction medium contained 5 mM MgSO₄. Values between brackets represent the activities in the presence of various substrats as a percent of the activity in presence of ATP.

Substrate	Initial rate of substrat-dependent quenching of quinacrine fluorescence (AU min ⁻¹ mg ⁻¹ protein)
ATP	$61.3 \pm 4.7 (100)$
GTP	4.0 (6.5)
PPi	13.7 (22.3)
ADP	16.3 (26.6)
PEP	10.3 (16.8)

Effect of some inhibitors on H+pumping activity

H+pumping was tested in presence of Ca²⁺ channel antagonist (verapamil), animal Na+/K+ ATPase inhibitor (ouabain), plant ATPase inhibitors (EB, DES, VO₄, DCCD), and photosynthetic electron transport inhibitor (DCMU).

The obtained results indicated that the plant ATPase inhibitors, inhibited the H+pump activity of spinach leaves with more effective inhibition obtained for VO₄ and DCCD than for EB or DES. Some experimental problems with these later inhibitors have limited the study of there effect on the H+ATPase according to there concentration.

The ouabain had almost no effect on the H+pump suggesting that the secondary structure of the animal Na+/K+ATPase is different from that of the spinach H+ATPase. Suprisingly DCMU inhibited also the enzyme by about 50% at the indicated concentration. This could support the mechanism of the H+ transport by the ATPase proposed by Briskin *et al.* (1992).

The verapamil known as a plant Ca²⁺ channel antagonist inhibited strongly the H+ATPase of spinach leaves. This result could explain, in the context of the acid growth theory, the inhibition of the elongation rate of spinach petiole as described by Malatialy (1990).

TABLE 2.

Effect of some inhibitors on the H⁺pumping activity of the spinach leaves plasma membrane. The plasma membrane was incubated in the reaction medium in the presence of the inhibitors for 10 min and the reaction was started by addition of 1.5 mM Mg ATP. The activities in the presence of various inhibitors were expressed as the percent of the activity of the control (without inhibitors). Stock solutions of DCCD and DES were prepared in ethanol and the specific activities were compared to the activity of the control assay containing 1% ethanol which is very similar to that of the control without ethanol.

Inhibitors	Percent of the specific activity to the control
Control	100
Verapamil 100µM	23
Ouabain 100µM	93
DCMU 100µM	47
Erythrosin B	
lμM	91
2.5μΜ	67
DES	
30µM	85
100μM	63
Na3VO4	
20µM	95
200μΜ	49
2000μM	23
DCCD 500µM	22

SDS PAGE and Western blot

SDS-PAGE of the purified PM of spinach leaves revealed after Coumassie brillant blue G250 staining, a large number of bands of different molecular weight, particularly a polypeptide of about 100 kDa (fig. 3 A) which corresponds in molecular mass to the PM ATPase. This polypeptide cross-reacted with the polyclonal antibodies directed against the PM ATPase of *Arabidopsis thaliana* (arrow), kindly gived by Prof. R. Serrano (fig. 3 B). On the same blot we can observe a contaminating band of lower molecular weight which probably resulted from proteolysis of the 100 kDa polypeptide.




Fig. 3.

A. SDS polyacrylamide gel electrophoresis of the purified plasma membrane (90 µg protein). The polypeptides were stained by Coumassie brillant blue G250. B. immunoblot analysis of the purified plasma membrane. The arrow indicates the polypeptide of about 100 kDa. The experimental procedure was described under Materials and Methods.

ACKNOWLEDGMENTS

We would like to thank Professor R. Serrano from Department of Biotechnology (Valencia, Spain) for his kind gift of polyclonal antibodies against PM ATPase of *Arabidopsis thaliana*.

RÉSUMÉ

La membrane plasmique des feuilles d'épinard a été purifiée par partition de phase et l'activité pompe à proton associée à la membrane plasmique caractérisée. Elle présente un pH optimum à 6.7 similaire à celui de l'activité ATPase, et une spécificité du substrat plus évidente pour l'ATP avec un Km_{app} de 0.697. L'activité de cette pompe est plus ou moins inhibée par différents inhibiteurs des ATPases, et spécialement par le vanadate avec un I₅₀ de 160 µM. L'électrophorèse sur gel de polyacrylamide, des protéines de la membrane plasmique purifiée, a révélé la présence d'une bande à 100 kDa qui a réagit avec les anticorps polyclonaux anti ATPase de la membrane plasmique de l'*Arabidopsis thaliana*.

REFERENCES

- ASSMANN S. M., SIMONCINI L. & SCHROEDER J. I. (1985). Blue light activates electrogenic ion p'umping in guard cell protoplasts in *Vicia faba*. *Nature*, 318: 285-287.
- BARBIER-BRYGOO H., EPHRITIKHINE G., KLÄMBT D., GHISLAIN M. & GUERN J. (1989). Functional evidence for an auxin receptor at the plasmalemma of tobacco mesophyll protoplasts. *Proc. Natl. Acad. Sci. USA.*, 86: 891-895.
- BECKER D., ZEILINGER C., LOHSE G., DEPTA H. & HEDRICH R. (1993). Identification and biochemical characterization of the plasma membrane H+ATPase in guard cells of *Vicia faba* L. *Planta*, 190: 44-50.
- Bellamine J., Penel C. & Greppin H. (1993 a). Proton pump and IAA sensitivity changes in spinach leaves during the flowering induction. *Plant Physiol. Biochem.*, 31: 197-203.
- Bellamine J., Arrizabalaga Ph. & Greppin H. (1993 b). Characterization and identification of plasma membrane before and after flowering induction by Fourier transform infrared spectroscopy using horizontal attenuated total reflectance. *Archs Sci. Genève*, 46: 361-372.
- BIDWAI A. P. & TAKEMOTO J. Y. (1987). Bacterial phytotoxin, syringomycin, induces a protein kinase-mediated phosphorylation of red beet plasma membrane polypeptides. *Proc. Natl. Acad. Sci. USA*, 84: 6755-6759.
- BRISKIN D. P. & HANSON J. B. (1992). How does the plant plasma membrane H+ATPase pump protons? J. Exp. Bot., 43: 269-289.
- CLELAND R.E. (1987). How hormones work-Auxin and cell elongation. *In: Plant hormones and their role in plant growth and development*. P.J. Davies (Ed), Martinus Nijhoff Publishers, pp.132-149.
- DE MICHELIS M. I. & SPANSWICK R. M., (1986). H+ pumping driven by the vanadate-sensitive ATPase in membrane vesicles from corn roots. *Plant Physiol.*, 81: 542-547.
- François J. M., Bervillé A. & Rossignol M. (1992). Development and line dependent variations of Petunia Plasma membrane H+ATPase sensitivity to auxin. *Plant Sci.*, 87: 19-27.
- GROUZIS J. P., GIBRAT R., RIGAUD J., AGEORGES A. & GRIGNON C. (1990). Potassium stimulation of corn root plasmalemma ATPase. I. Hydrolytic activity of native vesicles and purified enzyme. *Plant Physiol.*, 93: 1175-1182.
- HARPER J. F., SUROWY T. K. & SUSSMAN M. R. (1989). Molecular cloning and sequence of cDNA encoding the plasma membrane proton pump (H+ATPase) of Arabidopsis thaliana. *Proc. Natl. Acad. Sci.*, 86: 1234-1238.
- JACOBS M. & TAIZ L. (1980). Vanadate inhibition of auxin-enhanced H+ secretion and elongation in pea epicotyls and oat coleoptiles. *Proc. Natl. Acad. Sci.* USA, 77: 7242-7246.
- LAEMMLI U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, 227, 680-685.

- MALATIALY M. L. (1990). Croissance et floraison chez *Spinacia oleracea* L. cv. Nobel: Etude, *in vitro*, du transport du calcium et de l'interaction avec le tonoplaste et le plasmalemme. *Ph. D. diss. Geneva University*.
- OLIVARI C., PUGLIARELLO M. C., RASI-CALDOGNO F. & DE MICHELIS M. I. (1992). Characteristics and regulatory properties of the H+ATPase in a plasma membrane fraction purified from *Arabidopsis* thaliana. Bot. Acta, 106: 13-19.
- PARDO J. M. & SERRANO R. (1989). Structure of a plasma membrane H+ATPase gene from the plant Arabidopsis thaliana. J. Biol. Chem., 264: 8557-8562.
- PEDERSON P. L. & CARAFOLI E. (1987). Ion motive ATPases.I. Ubiquity, Properties and Significance to Cell Function. TIBS, 12: 146-150.
- RASI-CALDOGNO F., PUGLIARELLO M. C., DE MICHELIS M. I. (1985). Electrogenic transport of protons driven by the plasma-membrane ATPase in membrane vesicles from Radish. *Plant Physiol.* 77: 200-205.
- RAYLE D.L. & CLELAND R.E. 1992. The acid growth theory of auxin-induced cell elongation is alive and well. *Plant physiol.* 99, 1271-1274).
- Santoni V., Vansuyt G. & Rossignol M. (1991). The changing sensitivity to auxin of the plasma membrane H+ATPase: Relationship between plant development and ATPase content of membranes. *Planta*, 185: 227-232.
- SERRANO R. (1989). Structure and function of plasma membrane ATPase. Annu. Rev. Plant Physiol. Plant Mol. Biol., 40: 61-94.
- SERRANO R. (1988). Structure and function of proton translocating ATPase in plasma membranes of plants and fungi. *Bioch. Bioph. Acta*, 947: 1-28.
- TOWBIN H., STAEHELIN T. & GORDON J., (1979). Electrophoretique transfer of proteins from polyacrylamide gels to nitrocellulose sheets: Procedure and some applications. *Proc. Natl. Acad. Sci. USA*, 76, 4350-4354.