

Zeitschrift: Archives des sciences et compte rendu des séances de la Société
Herausgeber: Société de Physique et d'Histoire Naturelle de Genève
Band: 51 (1998)
Heft: 3

Artikel: Oribatids from Singapore II. (Acari, Oribatida) : new and interesting mites from the Geneva Museum LXXIII
Autor: Mahunka, Sándor
DOI: <https://doi.org/10.5169/seals-740158>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 01.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

ORIBATIDS FROM SINGAPORE II. (ACARI: ORIBATIDA) (NEW AND INTERESTING MITES FROM THE GENEVA MUSEUM LXXIII)

BY

Sándor MAHUNKA¹

(Ms. reçu le 2.12.1998, accepté le 14.12.1998)

ABSTRACT

Oribatids from Singapore II. (Acari: Oribatida). New and interesting mites from the Geneva Museum LXXIII. - A new genus and species is described from the tropical rain-forest Reserve of Bukit Timah in Singapore: *Singabodes rarus* gen. n., sp. n. (Carabodidae). Additional remarks on *Bukitritia timah* Mahunka, 1990 are given.

Key-words: Acari, Oribatida, Taxonomy, New genus, new species, Singapore.

INTRODUCTION

Dr Bernd Hauser, former Head of the Arthropod Department of the Muséum d'histoire naturelle, Geneva, visited Singapore during most of his 8 expeditions concerned with a research programme on soil fauna (especially Microarthropoda) of the rain forests of South-East Asia and surrounding areas (MAHUNKA, 1995). These studies were organized together with Dr Charles Lienhard, Research Officier at the same Department, the opportunity was taken to collect soil samples at different points on the island.

Drs Hauser and Lienhard were generously helped in this project by Prof. Dr D.H. Murphy, Department of Zoology, National University of Singapore, who not only accompanied them but also gifted soil samples collected by himself (MURPHY, 1973) to the Geneva team.

This is the second contribution on this very interesting Orbatid material from Singapore, conserved in the collections of the Geneva Museum (MAHUNKA, 1989). A special focus is the study of the fauna of the famous nature reserve of Bukit Timah, "perhaps the oldest small rain-forest reserve in the world" (CORLETT, 1988). The high diversity of the fauna of moss mites of this reserve is a evidence for the value of conserving small fragments of lowland tropical rain forest (TURNER & CORLETT, 1996).

DESCRIPTION AND REMARKS

Euphthiracaridae JACOT, 1930

Bukitritia timah MAHUNKA, 1990
(Fig. 1)

¹ Zoological Department, Hungarian Natural History Museum, Baross utca 13, H-1088 Budapest, Hungary.




FIG. 1
Bukitrititia timah Mahunka, 1990 - 1: body in lateral view.

This taxon was described by the author in a paper which summarized the relationships of the superfamily Euphthiracaroidea (MAHUNKA, 1990). The description (both of genus and species) remains valid except for the number of lyrifissures: the correct number is four (and not five).

The complete data of the material examined are: Holotype: Singapore: Bukit Timah Nature Reserve, forest litter, sandy dry slope lower JFV; 25.VI.1969; leg. D.H. Murphy (D 11-2); 1 paratype: from the sample; 1 paratype: Singapore: Bukit Timah Nature Reserve, forest litter, streamside litter wet; 2.VII.1969; leg. D.H. Murphy (D 11-9). Holotype and 1 paratype: MHNG², 1 paratype (1374-PO-89): MNHM³.

Derivation nominis: The names of the genus and species refer to the hill of Bukit Timah which, 162'5 m, is the highest point of the island, and to the Nature Reserve of 75 hectares.

Carabodidae C.L. Koch, 1837

Singabodes gen. n.

D i a g n o s i s : Family *Carabodidae*. Prodorsum simple, interlamellar setae arising on the lamellae. Sensillus long and clavate. Notogaster with a wide, smooth, projecting median field, and with a deep hollow posteriorly. Fourteen pairs of modified notogastral setae present, five pairs arising around the hollow, are long and fringed, two pairs originate in the humeral region, the other four pairs in a posteromarginal position, one pair in front of the hollow and two pairs near to the anterior margin of the notogaster.

² MHNG = deposited in the Muséum d'Histoire naturelle, Geneva.

³ HNHM = deposited in the Hungarian Natural History Museum, Budapest, with identification number of the specimens in the Collection of Arachnida.

Epimeral setal formula: 3-1-3-3. Anogenital region well sclerotized, ornamented by rugae, hollows and crests. The leg chaetotaxy is of the "reduced type".

Type species: *Singabodes rarus* sp. n.

R e m a r k s : The characteristic notogastral structure, combined with the shape of setae and sensillus, makes it impossible to place this new species in any of the so far known genera of Carabodidae (MAHUNKA, 1986).

D e r i v a t i o n o n o m i n i s : allusion to the name of Singapore: Singapura in Sanskrit (= Lion City), this name was given at the end of the 13th century by the prince Sang Nila Utama from Sumatra who landed in Temasek (now Fort Canning Hill), the first settlement on the island (HAENLEIN, 1985).

Singabodes rarus sp. n.

(Figs 2-14)

M a t e r i a l e x a m i n e d : Holotype: Singapore: Bukit Timah Nature Reserve, forest litter; 9.VII.1969; leg. D.H. Murphy (D 12-7); 1 paratype: from the same sample. Holotype: MHNG², paratype (1377-PO-89): NHNM³.

M e a s u r e m e n t s . - Length: 344-384 µm, width: 127-153 µm.

P r o d o r s u m : Rostrum elongated, rostral setae arising anterior of the lamellar setae (Fig. 2). Some large and round alveoli visible in this region. Lamellae narrow, anterior part ornamented by transversal striations (Fig. 2). Lamellar setae originating on the inner margin of lamellae, approximately lanceolate; the two other pairs of prodorsal setae phylliform, their surface well spiculate. Interlamellar region with some irregular spots medially. Sensillus long, gradually widened distally, clavate. Tutorium strong, some other crests also observable on the lateral part of prodorsum (Fig. 5).

N o t o g a s t e r : Dorsosejugal region slightly concave. Median part of notogaster smooth, lateral surface granulate, enclosed between these two parts is a clearly polygonate surface with tubercles. Median field with an elevation posteriorly (in which the hollow is situated) (Fig. 7). The hollow consists of two parts (Fig. 6) and some uninterpreted structures are visible at a deeper level. Among the notogastral setae nine pairs are small, approximately round, or phylliform, five pairs, around the hollow, large, with long fringes.


V e n t r a l s i d e (Fig. 8): Mentum coarsely foveolate. Setae *h* simple and short, originating near to each other. Coxisternal region wide, the widest anteriorly, near to the very large pedotecta 1. Apodemes and epimeral borders conspicuous. Between the epimeres a wide median field. All epimeral setae minute. Anogenital region divided and ornamented by ribs and crests. Genital and anal apertures removed far from each other. Anogenital setal formula: 4-1-2-3. Genital and aggenital setae widened basally, with pointed distal ends. Anal setae simple, adanal setae round, phylliform. Setae *ad*₁ and *ad*₂ in postanal, setae *ad*₃ in preanal position.

L e g s : Chaetotaxy of legs:

I: 1-4-3+1-4+2-15+2-1 (Figs 9, 10)


II: 1-4-3+1-3+1-15+2-1! (Figs 11, 13)

IV: 1-2-2-2+1-12-1 (Fig. 14)



FIGS 2-7

Singabodes rarus gen. n., sp. n. - 2: body in dorsal view, 3: rostrum, 4: lateral part of prodorsum, 5: prodorsum in lateral view, 6: median hollow of notogaster, 7: posterior part of notogaster in lateral view.



FIGS 8-14

Singabodes rarus gen. n., sp. n. - 8: body in ventral view, 9: tibia and tarsus of leg I, 10: genu of leg I, 11: tibia and genu of leg II, 12: femur and genu of leg III, 13: leg II, 14: leg IV.

Surface of some joints of legs: granulate or pustulate, femora and trochanters of legs III (Fig. 12) and IV (Fig. 14) with strong dorsal protuberances or a wide blade-like ventral formation. Ventral setae (*pv, a*) of tarsi and tibiae of legs I and II (Figs 9, 10) modified, spiniform; setae *u* very wide, blunt at tip. Some solenidia (of legs II and IV) sharply pointed.

Derivation nominis: allusion to the rarity of this taxon.

ACKNOWLEDGEMENTS

I am very grateful to Dr Malcolm Luxton (National Museum of Wales, Cardiff) for his careful reading of the manuscript and his comments.

RÉSUMÉ

ORIBATES DE SINGAPOUR II (ACARI: ORIBATIDA) (ACARIENS NOUVEAUX ET INTÉRESSANTS DU MUSÉUM DE GENÈVE LXXIII)

Le travail contient la description d'un genre nouveau et d'une espèce nouvelle de la Réserve de forêt pluviale Bukit Timah de Singapour: *Singabodes rarus* gen. n., sp. n. (*Carabodidae*) et des remarques complémentaires sur *Bukitritia timah* MAHUNKA, 1990.

Mots-clés: Acariens, Oribates, taxonomie, nouveau genre, nouvelle espèce, Singapour.

REFERENCES

- CORLETT, R.T. 1988. Bukit Timah: the history and significance of a small rain-forest Reserve. *Envir. Conserv.* 15: 37-44.
- HAENLEIN, R. 1985. Singapur. *Polyglott-Reiseführer* 168, *Polyglott-Verlag, München*, 64 pp.
- MAHUNKA, S. 1986. A survey of the family Carabodidae C.L. Koch, 1836 (Acari: Oribatida). *Acta zool. hung.* 32: 73-135.
- MAHUNKA, S. 1989. New and interesting mites from the Geneva Museum LXIV. Oribatids from Singapore (Acari: Oribatida). *Revue suisse Zool.* 96: 381-402.
- MAHUNKA, S. 1990. A survey of the superfamily Euphthiracaroidea Jacot, 1930 (Acari: Oribatida). *Folia ent. hung.* 51: 37-80.
- MAHUNKA, S. 1995. Oribatids from Brunei I (Acari: Oribatida). New and interesting mites from the Geneva Museum LXXV. *Revue suisse Zool.* 102: 913-942.
- MURPHY, D.H. 1973. Animals in the Forest Ecosystem, pp. 53-73. In: S.H. CHUANG (ed.). *Animal Life and Nature in Singapore*. *Singapore University Press*.
- TURNER, I.M. & R.T. CORLETT. 1996. The conservation value of small, isolated fragments of lowland tropical rain forest. *Trends Ecol. Evolut.* 11: 330-333.