

Zeitschrift: Archives des sciences et compte rendu des séances de la Société
Herausgeber: Société de Physique et d'Histoire Naturelle de Genève
Band: 53 (2000)
Heft: 3

Artikel: NADH oxidase activities of intact leaf discs of spinach
Autor: Barr, Rita / Penel, Claude / Greppin, Hubert
DOI: <https://doi.org/10.5169/seals-740507>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 16.11.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Communication présentée à la séance du 17 mai 2000

NADH OXIDASE ACTIVITIES OF INTACT LEAF DISCS OF SPINACH

BY

Rita BARR¹, Claude PENEL^{3*}, Hubert GREPPIN³ & D. James MORRÉ²

ABSTRACT

NADH oxidase activities of intact leaf discs of spinach. - Leaf disks, 1 cm in diameter, were punched from leaves of market spinach and used for assay of NADH oxidase. The disks exhibited a steady state rate of NADH oxidation of about 1 nmole/min/disk. The activity was proportional to time of incubation and to the number of disks. The activity was stimulated by the synthetic auxin 2,4-D to a maximum near 1 μ M 2,4-D. The disks provide a convenient system to measure the cell surface NADH oxidase in leaves.

Abbreviations: NADH, reduced nicotinamide adenine dinucleotide; NOX, plasma membrane-associated NADH oxidase; 2,4-D, 2,4-dichlorophenoxyacetic acid; IAA, indole-3-acetic acid.

INTRODUCTION

Ongoing work in the laboratory has identified an external cell surface NADH oxidase of soybean hypocotyls (HICKS and MORRE, 1998) that can be readily assayed using intact hypocotyl sections (BARR *et al.*, 2001).

The activity correlated well with findings from right side-out isolated plasma membrane vesicles (DEHAHN *et al.*, 1997). The cell surface NADH oxidase of soybean assayed either with intact sections or with isolated right side-out vesicles was stimulated by active auxins 2,4-dichlorophenoxyacetic acid (2,4-D) or indole-3-acetic acid (IAA), but was unaffected by inactive or weak auxins 2,3-dichlorophenoxyacetic acid (2,3-D) or β -naphthaleneacetic acid (β -NAA) (MORRÉ *et al.*, 1988). Additionally the auxin-stimulated components of the surface NADH oxidase and of growth were blocked by thiol reagents such as N-ethylmaleimide (BARR *et al.*, 2001). Similar results were obtained with plasma membrane vesicles where the auxin-stimulated component of the activity was blocked by thiol reagents (MORRÉ *et al.*, 1995b).

In this report, the stimulation of NADH oxidase activity by auxin already reported for plasma membrane vesicles and intact soybean stem sections was extended to leaf disks of spinach. The system provides a convenient method for the study of the surface NADH oxidase of leaves.

¹ Department of Biological Sciences and ² Department of Medicinal Chemistry and Molecular Pharmacology, Hansen Life Sciences Research Building, Purdue University, West Lafayette, Indiana 47907, USA. ³ Laboratoire de Biochimie et Physiologie Végétale, Université de Genève, 3 Place de l'Université, CH-1211 Geneva 4, Switzerland.

*Author for correspondence and reprints. E-mail: Claude.Penel@bota.unige.ch

MATERIALS AND METHODS

Plant material

Commercial spinach (*Spinacia oleracea* L.) was used. One cm diam leaf disks were punched using a sharp cork borer. Disks from 10 to 20 different spinach leaves were collected directly into 0.1 M sucrose, 5 mM CaCl₂, 10 mM NaCl and 10 mM KCl (sucrose-salts). The sections were maintained at room temperature before and during all experiments.

NADH oxidase activity

The conditions selected were to incubate 10 to 20 of the 1 cm diam. disks in 30 ml of sucrose-salts containing 100 μ M NADH with mild initial shaking. The rate of NADH oxidation was determined on a 6 ml sample from the decrease in A₃₄₀ using a Sequoia-Turner Model 340 spectrophotometer. After determining the optical density, the solution was returned to the flask and the incubation continued. The blank rate in the absence of disks was negligible (0.1 O.D. units per 180 min).

RESULTS

Based on the decrease in absorbance at 340 nm, 1-cm leaf disks cut from leaves of spinach oxidize NADH (Fig 1A). The rate in the absence of 2,4-D or KCN calculated from the disappearance of NADH (decrease in absorbance at 340 nm) was 9 nmol/min/10 1-cm disks (Fig. 1). Following addition of 2,4-D, the rate of NADH disappearance was increased by about 50% to 13 nmol/min/10 1 cm diam. leaf disks (Fig. 1). The blank rate in the presence of NADH but in the absence of tissue (<0.5 nmol/min) was subtracted. In the presence of tissue but in the absence of NADH, the absorbance increased slightly. No rates were observed with tissue segments boiled for 10 min prior to assay. Distilled water or 50 mM Tris-Mes, pH 7.0, gave results equivalent to sucrose-salts.

The rate of NADH oxidation was proportional to number of disks (Figs. 2 and 3). The 2,4-D-stimulated component of the cell surface NADH oxidase was resistant to 1 mM potassium cyanide (Fig. 1B). This resistance was seen only with the auxin-stimulated component of the activity. Both in the presence or absence of 2,4-D, the rate of NADH oxidation by the leaf disks was greatly reduced by the addition of 1 mM cyanide. However in the presence of 2,4-D, a 2,4-D-induced increment in activity in the presence of cyanide (Fig. 1B) similar to that observed in the absence of cyanide (Fig. 1A) was observed. The cyanide-resistant component was proportional to the number of leaf disks in the cuvette (Fig. 3). The degree of auxin stimulation (proportion of total) was enhanced considerably over that in the absence of cyanide because the cyanide-susceptible component of NADH oxidation was reduced.

The response of NADH oxidation to the logarithm of 2,4-D concentration revealed an optimum at 1 μ M with the 1-cm disks (Fig. 4). The weak auxin 2,3-dichlorophenoxyacetic acid, at 1 μ M, elicited no auxin-stimulated rate (not shown). With 2,4-D, the total

2,4-D-induced increment of activity was increased by about 10 nmoles/min/30 sections both in the absence and presence of cyanide and the ratio of plus 1 μM 2,4-D to no 2,4-D was greatly augmented due to the reduced background in the presence of cyanide (Fig. 4).




FIG. 1.

Time course of NADH oxidation catalyzed by 10 1-cm diameter spinach leaf disks as determined by the decrease in absorbance at 340 nm in the absence (solid symbols) or presence (open symbols) of 1 μM 2,4-D. A: no KCN. B: 1 mM KCN.

DISCUSSION

Since an extracellular source of NADH in plants sufficient to reach concentrations at or near the K_m for the surface NADH oxidase is highly unlikely, some constituent endogenous to the plant plasma membrane, such as hydroquinones or protein thiols, has been assumed to be the physiological electron donor for the 2,4-D-stimulated cell surface NADH oxidase activity. Both protein thiols (CHUEH *et al.*, 1997) and reduced phyloquinone (BRIDGE *et al.*, 2000) have been implicated. In addition to oxygen (MORRÉ, 1994), protein disulfides appear to serve as acceptors of protons and electrons coming




FIG. 2.

Cell surface plasma membrane NADH oxidase is proportional to number of 1-cm diameter spinach leaf disks in the absence (solid symbols) or presence (open symbols) of 1 μ M 2,4-D and in the absence of KCN. Values are averages of 3 experiments \pm standard deviations among experiments.




FIG. 3.

Cell surface plasma membrane NADH oxidase is proportional to number of 1-cm diameter spinach leaf discs in the absence (solid symbols) or presence (open symbols) of 1 μ M 2,4-D but in the presence of 1 mM KCN. Values are averages of 3 experiments \pm standard deviations among experiments.




FIG. 4.

Response of NADH oxidase of 10 1-cm diameter leaf disks of spinach to the logarithm of 2,4-D concentration in the absence and presence of 1 mM KCN. The 2,4-D-responsive cell surface plasma membrane NADH oxidase is cyanide resistant. Values are averages of 2 or 3 experiments \pm standard deviations among experiments. A: no KCN. B: 1 mM KCN.

from NADH (CHUEH *et al.*, 1997) or from the physiological electron donor, reduced phyloquinone (BRIDGE *et al.*, in press). A model has been developed where the protein disulfide-thiol interchange portion of the activity (MORRÉ *et al.*, 1995) could function in physical membrane displacements important to the enlargement growth of plant cells (MORRÉ, 1994).

An external location of the growth-related NADH oxidase was first raised for plants from studies comparing the 2,4-D-responsive oxidation of NADH by right side-out and inside-out plasma membrane vesicles (DEHAHN *et al.*, 1997): Both populations of vesicles contained an NADH oxidase activity but the component of the activity stimulated by 2,4-D was accessible only to NADH supplied to right side-out vesicles and was, therefore,

located at the external cell surface. The possibility of an external NADH oxidase in plants was raised early by LIN (1984) for plant roots. However, until recently (MORRÉ, 1998), the functional significance of an external NADH oxidase was unclear.

A similar situation of having NADH oxidase activity at both the internal and external surfaces of the plasma membrane exists for cultured mammalian cells as well. That at least one of the NADH oxidase activities associated with the mammalian cell surface was external was indicated from studies of the antitumor sulfonylurea-inhibited NADH oxidase activity of cervical carcinoma (HeLa) derived human cell line (MORRÉ, 1995). The drug-inhibited oxidase was observed only with vesicles of right side-out orientation (MORRÉ, 1995). Intact HeLa cells subsequently were shown to exhibit the drug-responsive activity (MORRÉ *et al.*, 1997) as would be expected for a drug-inhibited NADH oxidase associated with the cell surface. That both the drug and the NADH site of the drug-responsive NADH oxidase were external was demonstrated subsequently by inhibition of NADH oxidase activity using impermeant conjugates of an antitumor sulfonylurea (KIM *et al.*, 1997).

The resistance of external NADH oxidase to inhibition by cyanide was not unexpected since the assay procedure for the auxin-responsive NADH oxidase of plants (BRIGHTMAN *et al.*, 1988, MORRÉ *et al.*, 1986) as well as that for the drug inhibited activity of mammalian membranes (MORRÉ *et al.*, 1995) has included cyanide in the assay mixture. Clearly with the intact hypocotyl sections of the soybean, the 2,4-D responsive activity was cyanide resistant as was the total NADH oxidase activity of right-side-out plasma membrane vesicles (BRIGHTMAN *et al.*, 1988). What was unexpected considering only assays of isolated plasma membrane vesicles was that the intact sections appeared to exhibit an NADH oxidase activity that was cyanide sensitive. Such an activity normally has been low or absent with plasma membrane vesicles. It has previously been attributed to mitochondrial contamination (MOYA-CAMARENA *et al.*, 1995). In the work with intact soybean sections, it was unlikely that the cyanide sensitive activity was from cut surfaces since neither washing the sections nor sealing the ends of the sections with lanolin eliminated the activity. The explanation proposed is that the cyanide-sensitive activity was the result of a peroxidase that was either loosely associated with the cell surface or was present in the apoplast and normally lost during the preparation of plasma membranes. A similar phenomenon was observed for intact soybean sections (BARR *et al.*, 2001) and this explanation may be equally applicable to leaf disks of spinach.

RÉSUMÉ

ACTIVITÉS NADH OXYDASE DE DISQUES INTACTS DE FEUILLE D'ÉPINARD

Des disques sont prélevés sur des feuilles d'épinard et utilisés pour une mesure directe de l'activité NADH oxydase. Mis en présence de NADH, les disques oxydent le NADH de façon continue au taux d'environ 1 nmole/min/disque. Cette activité est linéaire dans le temps et proportionnelle au nombre de disques de feuilles. Elle est stimulée par l'auxine synthétique 2,4-D, avec un maximum à 1 μ M. L'utilisation de disques de feuilles permet l'étude de la NADH oxydase présente sur la surface des cellules.

REFERENCES

- BARR, R., D.J. MORRÉ & F.L. CRANE. 2001. Oxidation of NADH by intact segments of soybean hypocotyls is stimulated by 2,4-D. *Plant Physiol. Biochem.* 38: 739-745.
- BRIDGE, A., R. BARR & D.J. MORRÉ. 2000. The plasma membrane NADH oxidase of soybean has vitamin K₁ hydroquinone oxidase activity. *Biochim. Biophys. Acta* 1463: 448-458.
- BRIGHTMAN, A.O., R. BARR, F.L. CRANE & D.J. MORRÉ. 1988. Auxin-stimulated NADH oxidase purified from plasma membrane of soybean. *Plant Physiol.* 86: 1264-1269.
- CHUEH P-J, D.M. MORRÉ, C. PENEL, T. DEHAHN & D.J. MORRÉ. 1997. The hormone-responsive NADH oxidase of the plant plasma membrane has properties of a NADH:protein disulfide reductase. *J. Biol. Chem.* 272: 11221-11227.
- DEHAHN T, R. BARR & D.J. MORRÉ DJ. 1997. NADH oxidase activity present on both the external and internal surfaces of soybean plasma membranes. *Biochim. Biophys. Acta* 1328: 99-108.
- HICKS C. & D.J. MORRÉ. 1998. Oxidation of NADH by intact segments of soybean hypocotyls and stimulation by 2,4-D. *Biochim. Biophys. Acta* 1375: 1-5.
- KIM C, W.C. MACKELLAR, N. CHO, S.R. BYRN & D.J. MORRÉ. 1997. Impermeant antitumor sulfonylurea conjugates that inhibit plasma membrane NADH oxidase and growth of HeLa cells in culture for identification of binding proteins from sera of cancer patients. *Biochim. Biophys. Acta* 1324: 171-181.
- LIN, W. 1984. Further characterization on the transport property of plasmalemma NADH oxidation system in isolated corn root protoplasts. *Plant Physiol.* 74: 219-222.
- MORRÉ, D.J. 1994. The hormone- and growth factor-stimulated NADH oxidase. *J Bioenerg. Biomembr.* 26: 421-433.
- MORRÉ, D.J. 1995. NADH oxidase activity of HeLa plasma membranes inhibited by the antitumor sulfonylurea *N*-(4-methylphenylsulfonyl)-*N'*-(4-chlorophenyl)urea (LY181984) at an external site. *Biochim. Biophys. Acta* 1240: 201-208.
- MORRÉ, D.J. 1998. NADH oxidase: A multifunctional ectoprotein of the eukaryotic cell surface, in: *Plasma Membrane Redox Systems and their Role in Biological Stress and Disease*. H. Asard, A Berczi, RJ Caubergs, eds., Kluwer, Dordrecht, pp 121-156.
- MORRÉ, D.J., P. NAVAS, C. PENEL & F.J. CASTILLO. 1986. Auxin-stimulated NADH oxidase (semidehydroascorbate reductase) of soybean plasma membrane: Role in acidification of cytoplasm? *Protoplasma* 133: 187-193.
- MORRÉ DJ, A.O. BRIGHTMAN, L.-Y. WU, R. BARR, B. LEAK & F.L. CRANE. 1988. Role of plasma membrane redox activities in elongation growth in plants. *Physiol. Plant.* 73: 187-193.
- MORRÉ DJ, A.O. BRIGHTMAN, A. HIDALGO & P. NAVAS. 1995a. Selective inhibition of auxin-stimulated NADH oxidase activity and hypocotyls by thiol reagents. *Plant Physiol.* 107: 1285-1291.
- MORRÉ DJ, R. DE CABO, E. JACOBS & D.M. MORRÉ. 1995b. Auxin-modulated protein disulfide-thiol-interchange activity from soybean plasma membranes. *Plant Physiol.* 109: 573-578.
- MORRÉ DJ, L.-Y. WU & D.M. MORRÉ. 1997. Inhibition of NADH oxidase activity and growth of HeLa cells by the antitumor sulfonylurea *N*-(4-methylphenylsulfonyl)-*N'*-(4-chlorophenyl)urea (LY 181984) and response to epidermal growth factor. *Biochim. Biophys. Acta* 1355:114-120.
- MORRÉ DJ, D.M. MORRÉ, C. PENEL & H. GREPPIN. 1998. Auxin modulated protein disulfide-thiol interchange activity from plasma membranes of spinach leaves responds to photoperiod and NADH. *Int. J. Plant Sci.* 159:105-109.
- MOYA-CAMARENA, S.Y., D.J. MORRÉ & D.M. MORRÉ. 1995. Response of NADH oxidation and growth in K-562 cells to the antitumor sulfonylurea, *N*-(4 methyl-phenylsulfonyl)-*N*-(C4-chlorophenyl)urea (LY181984). *Protoplasma* 188: 151-160.