

**Zeitschrift:** Archives des sciences et compte rendu des séances de la Société  
**Herausgeber:** Société de Physique et d'Histoire Naturelle de Genève  
**Band:** 54 (2001)  
**Heft:** 3

**Artikel:** Prebiotic phosphoramidation of nucleobases by Mg<sup>2+</sup> -triggered decyclization of trimetaphosphate  
**Autor:** Turian, Gilbert / Rivara-Minten, Elisabeth  
**DOI:** <https://doi.org/10.5169/seals-740526>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 02.05.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

Archs Sci. Genève	Vol. 54	Fasc. 3	pp. 233-238	Décembre 2001
-------------------	---------	---------	-------------	---------------

Communication présentée à la séance du 6 décembre 2001

## PREBIOTIC PHOSPHORAMIDATION OF NUCLEOBASES BY $Mg^{2+}$ -TRIGGERED DECYCLIZATION OF TRIMETAPHOSPHATE

BY

Gilbert TURIAN\* & Elisabeth RIVARA-MINTEN\*\*

### ABSTRACT

**Prebiotic phosphoramidation of nucleobases by  $Mg^{2+}$ -triggered decyclization of trimetaphosphate.** – Nucleobases can be phosphoramidate-bonded on trimetaphosphate decyclized by the nucleophilic attack of their reversibly tautomered units in stringent synergy with  $Mg^{2+}$  asymmetrically shielding the repellent  $OH^-$  charges of trimetaphosphate. The produced linear triphosphates are thermopolymerizable, after splicing, into polybasediphosphates.

**Key-words:** phosphoramidation, nucleobases, trimetaphosphate, decyclization.

### INTRODUCTION

In the realm of our interest in the prebiotic origin of the genetic code, we have postulated the prenucleic role of phosphoramidic (N~P) bonding of nucleobases recognizing amino acids of primal peptides (TURIAN, 1996-2000). This has incited us to experimentally check our hypothesis by  $^{31}P$  NMR identification of decyclized products of trimetaphosphate (TMP), essentially linear triphosphate (TP), presumably freed by the nucleophilic attack of thereby P~N bonded nucleobases (TURIAN *et al.*, 1999). Such decyclizations were obtained in 2,5 mM  $Mg^{2+}$  – containing mineral water and found to be enhanced by complements of 5mM  $Mg^{2+}$  ions. It was therefore interesting to further study the possible stringency of  $Mg^{2+}$  ions in this decycling process by incubation of the mixture in plain  $H_2O$  as solvent enriched or not (controls) in  $Mg^{2+}$  ions.

### MATERIALS AND METHODS

We have followed the recently described procedure (TURIAN *et al.*, 1999), but with the major change to 2 ml distilled  $H_2O$  (pH adjusted to 8.0 with molar NaOH) instead of mineral water (pH 7.7) as solvent containing 10 (5) mM of trimetaphosphate (TMP Sigma, grade III) enriched or not with 5 (10) mM of  $MgCl_2 \cdot 6H_2O$ . The solutions (2 ml)

\* Microbiologie générale, Département de Biologie végétale, Université de Genève, CH-1211 Genève 4.

\*\* Chimie organique pharmaceutique, Sciences II, Université de Genève, CH-1211 Genève 4.

were incubated in the presence of the chosen nucleobase (5 mM adenine or cytosine, Sigma products), from 18 to 23 days in shaken Pyrex capped miniflasks at 25°C.

The amounts of triphosphate (TP) decyclized from TMP have been compared by the intensity of their doublet peaks determined by  $^{31}P$ -NMR at 81 MHz on a AC200F Bruker NMR spectrometer using  $H_3PO_4$  as an external reference. These measures have been quantified by the increase of the  $^{31}P$  NMR signal of TP (at -6.0 ppm according to van WAZER *et al.* (1956), CALLIS *et al.* (1957) et VOGEL (1984)).

To ascertain the amidation of one of the phosphates of linear TP, we have thermopolymerized (120° C, 1h) the incubated mixtures of presumed polyphosphates and centrifuged their white precipitates, washed with slightly alkaline (pH 8)  $H_2O$  to insure the robustness of the P~N bond and lyophilized into powder (1) checked for its polyphosphate nature by metachromatic staining with toluidine blue (BRACHET, 1956, KORNBERG *et al.*, 1999) and (2) tested for putative nucleobase bonding by the UV (254 m $\mu$ ) quenching of the blue fluorescence of filter paper (SCHLEICHER & SCHÜLL) supports at the level of spots of wetted (distilled  $H_2O$  pH 8) powder .

## RESULTS AND DISCUSSION

Decyclization of TMP by hydrolytic opening of one of its phosphorus anhydride bonds (P – O – P – ...) into a linear TP did not occur in plain distilled  $H_2O$  (controls Fig. 1a) but required the addition of 5 mM  $Mg^{2+}$  as evidenced by the NMR detection of the double peak at ~ -6 ppm characteristic of TP (Figs 1b and 2a).

Nucleobase (adenine or cytosine) alone could not decyclize TMP into linear TP but only hydrolytically split it into traces of pyrophosphate signaled at ~ -8 ppm (Fig. 1d). Both required complementation by  $Mg^{2+}$  ions while overpowering the efficiency of these ions alone (Figs 1c and 2b).

Theoretically, the stringent role of  $Mg^{2+}$  in the process of TMP decyclization could be explained by their asymmetric shielding of the  $OH^-$  charges of the 3P atoms (Fig. 3a): the two positive charges of one  $Mg^{2+}$  would balance the  $OH^-$  charges of two P atoms while a second  $Mg^{2+}$  ion could only half devote one of its 2 positive charges to balance the charge of the third P, keeping its second positive charge free to ionize  $H_2O$  to  $H^+ + OH^-$ . This would thus provide an hydroxyl nucleophile (see WESTHEIMER, 1987) as “opener” of the cyclic bond of TMP while maintaining the  $OH^-$  group completing the fifth valence of the P atom at the tip of the now linear triphosphate (Fig. 3a). The additional presence of a tautomerized nucleobase endowed with an “aggressive” nucleophilic N group would overpower that of the nucleophile  $OH^-$  of  $H_2O$  while benefiting of the  $Mg^{2+}$  shield (Fig. 3b). This protonic tautomerization of the nucleobases by the low pH locally produced by  $H^+$  ions from ionized  $H_2O$  is thus a stringent condition for the transition of the NH group (1' in pyrimidines, 9' in purines) to the  $NH^+$  provider of the pair of free electrons nucleophilically attacking one of the electrophilic P atoms of TMP.

The N~P bonding of a nucleobase on decyclized TMP requiring the prototropic tautomerization of the base is only the initial step of the process. It must then be secondly enforced by an alkaline (pH>4) – induced tautomeric “back-shift” of its double bond from




FIG. 1.

$^{31}\text{P}$  - NMR spectra from 18-day incubation mixtures compared for doublet signals at  $\sim -6$  ppm (at  $-7.8$  for the pyrophosphate singlet\*) of triphosphate (TP) decyclized from a trimetaphosphate (TMP): a) TMP alone; b) TMP +  $\text{Mg}^{2+}$ ; c) TMP +  $\text{Mg}^{2+}$  + the nucleobase adenine; d) TMP + nucleobase alone.

the N1 atom to position 6-5 (pyrimidines) or from N9 to N8-7 (purines). Such shifts as proposed by Vogel (1984) for imidazole can thus also concern nucleobases to produce adenyl/cytosine triphosphates.

To experimentally check such locking of the nucleobases on the TPs, we separately thermopolymerized  $\text{Mg}^{2+}$ -TP and  $\text{Mg}^{2+}$  nucleobase triphosphates. To confirm their poly-




FIG. 2.

Idem to Fig. 1 (23-day incubation): a) TMP +  $Mg^{2+}$ ; b) TMP +  $Mg^{2+}$  + the nucleobase cytosine.




FIG. 3.

Decyclization of TMP permitted by  $Mg^{2+}$  shielding of  $PO^-$  groups allowing nucleophilic attack of one of the three P-O-P anhydride bonds by (a)  $Mg^{2+}$  - ionized  $H_2O$  synergistically overpowered by (b) the  $\geq HN:$  group of any P-N bonded nucleobase (*Nb*).




FIG. 4.

Slices of filter paper with its UV excited blue fluorescence: left, shadowed by nude polyphosphate; right, quenched by adenylyl polyphosphate.

phosphate nature, we positively tested the switching of their toluidine blue color into pink. In this thermoprocess, the 3P must have been spliced into 2P as revealed by the accumulation of  $P_{inorg}$  (signal at  $\sim 0$  ppm) and PP (signal at  $-7 - 8$  ppm) in the supernatant of the precipitated polyphosphates. Additionally, their adenine N-bonding was ascertained by the UV (254 m $\mu$  absorption test) showing a strong quenching of fluorescence restricted to the powder produced from the original nucleobase-TP (Fig. 4).

These preliminary results then lead us to consider the phosphoramidate linkage of adenine into adenyldiphosphate and, presumably, of the other nucleobase cytosine on diphosphates as nucleophosphate units of riboseless prenucleic polymers. A sequential bonding of nucleobases lined up on a polyphosphate backbone might well be relevant for pregenetic coding while highlighting Kornberg's (1995) prediction of a vicariant role of polyphosphates in the prebiotic evolution.

### RÉSUMÉ

**Phosphoramidation prébiotique de nucléobases par décyclisation du trimétaphosphate activée par les ions  $Mg^{2+}$ .** - Les nucléobases peuvent être phosphoramidolées sur des trimétaphosphates décyclisés par l'attaque nucléophile de leurs unités réversiblement tautomérisées en synergie obligatoire avec des ions  $Mg^{2+}$  protégeant les charges répulsives  $OH^-$  des trimétaphosphates. Les triphosphates linéaires produits sont thermopolymérisables, après épissure, en polybasediphosphates.

**Mots-clés:** phosphoramidation, nucléobases, trimétaphosphate, décyclisation.

### ACKNOWLEDGEMENTS

We are grateful to Dr P.-Y. Morgantini (Physical Chemistry Lab., Prof. J. Weber) for the graphics modeling and to Mrs Arlette Cattaneo for technical help.

### REFERENCES

- BRACHET J. 1956. *Biochemical Cytology*. Academic Press. Inc. Publ. New York, 516 p.
- CALLIS, C.F., J.R. VAN WAZER, J.R. SHOOLERY & W.A. ANDERSON. 1957. Principles of phosphorus chemistry. III. Structure proofs by nuclear magnetic resonance. *J. Amer. Chem. Soc.* 79: 2719-2726.

- KORNBERG, A. 1995. Inorganic polyphosphate: towards making a forgotten polymer unforgettable. *J. Bacteriol.* 177: 491-496.
- KORNBERG, A., N. N. RAO & D. AULT-RICHÉ, 1999. Inorganic polyphosphate: a molecule of many functions. *Annu. Rev. Biochem.* 68: 69-125.
- TURIAN, G. 1996. Polarity at onset of genetic coding. I. Bipolar bondings in the two-step takeover of peptide templates by prenucleic-ribonucleic acids. *Archs Sci. Genève* 49: 213-227.
- TURIAN, G. 1997. Polarity at onset of genetic coding. II. Primary recognition of amino acids by base doublets of prenucleic sugarless polymers secondarily taken over by ribonucleic acids. *Archs Sci. Genève* 50: 95-104.
- TURIAN, G. 2000. Model of prenucleic replication cyclically coupling encoding to decoding of peptide templates. *Archs. Sci. Genève* 53, 239-245.
- TURIAN, G., E. RIVARA-MINTEN & A. CATTANEO. 1999. Further  $^{31}P$ -NMR evidence of phosphoramidate bonding of nucleobases by  $Mg^{2+}$  enhanced nucleophilic attack on cyclic triphosphate. *Archs Sci. Genève* 52: 209-216.
- VAN WAZER, J. R., C.F. CALLIS, J. N. SHOOLERY & R.C. JONES. 1956. Principles of phosphorus chemistry. II. Nuclear magnetic resonance measurements. *J. Amer. Chem. Soc.* 78: 5715-5726.
- VOGEL, H.J., 1984.  $^{31}P$ -NMR studies on phosphoproteins, in *Phosphorus-  $^{31}P$  NMR Principles and Application*, Academic Press, Inc., New York, p. 105-153.
- WESTHEIMER, F.H. 1987. Why Nature chose phosphates. *Science* 235: 1173-1177.