

Zeitschrift: Archives des sciences et compte rendu des séances de la Société
Herausgeber: Société de Physique et d'Histoire Naturelle de Genève
Band: 55 (2002)
Heft: 2

Artikel: Adenine phosphoramide as a possible precursor of ATP
Autor: Turian, Gilbert
DOI: <https://doi.org/10.5169/seals-740295>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 01.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

ADENINE PHOSPHORAMIDE AS A POSSIBLE PRECURSOR OF ATP

BY

Gilbert TURIAN*

(*Ms. received 13.09.2002, accepted 15.10.2002*)

ABSTRACT

Adenine phosphoramide as a possible precursor of ATP. - The acid-opened phosphoramide bond of adenine triphosphate can possibly be bridged by amphihydroxylated ribose precursors and thereby evolve into adenosine triphosphate.

Key-words: phosphoramide, adenine, triphosphate, ribose, adenosine.

INTRODUCTION

“The prebiotic synthesis of nucleotides remains an unsolved challenge” (BRACK, 1998), because of the scarcity of ribose produced among a larger number of other sugars (SHAPIRO, 1988; JOYCE, 1989; JOYCE & ORGEL, 1999; LAHAV, 1999). The complex biotic synthesis of the “modern energy currency” adenosine triphosphate (ATP) which involves ribonucleotide formation, would therefore have been impeded in prebiotic conditions. In 1965, FRITZ LIPMANN, the founder of bioenergetics, had already proposed that simple condensed phosphates such as pyrophosphate (PPi), triphosphate or polyphosphates from primitive environmental sources might have been the forerunners of ATP, as further enforced by BALTSCHEFFSKY (1993) with his prebiotic “PPi world”.

However, consideration of the chemical evolution from a bottom-to-top point of view has led to predict a progressive arisal of affinity of the available phosphates for the early synthesized nucleobase adenine (5 HCN, ORÓ, 1960), in anticipation of their linking by ribose precursors ($C_2 + C_3$) produced by the well-known formose reaction (see SCHWARTZ, 1998).

Our finding of a prebiotic-like direct phosphoramidic (P~N) bonding (TURIAN *et al.* 1999; TURIAN & RIVARA-MINTEN, 2001), bypassing the double bonding – phosphoesterasic and glycosylic – of ribose, has recently suggested a stepwise insertion of ribose precursors between acid-induced opened P~N bonds of pre-RNA (TURIAN, 2001a) as also proposed below for the transition from sugarless adenine triphosphate to the adenosine triphosphate of ATP.

* Laboratoires de Microbiologie et Bioénergétique, Département de Biologie végétale, Université de Genève, CH-1211 Genève 4.

MODEL

It rests on the contrast between the robustness of our nucleobase P~N bondings at high pH (~8 TURIAN *et al.*, 1999), also known for imidazole (LOHRMANN & ORGEL, 1973), and other phosphoramides (CORBRIDGE, 1978) and the lability of their P~N bonds in acidic pH (< 4) conditions. This opens the interesting possibility to fill the gap — opened by acidification to pH 4-2 — between the nucleobase and the first phosphate group by ribosylating compounds (Fig. 1).





FIG. 1. Final step in the prebiotic origin of adenosine triphosphate implies acid-opening (pH < 4) P~N of the bonding between adenine (A) and triphosphate and further bridging by amphihydroxylated ribose precursors (Rp). For the initial steps of P~N bonding on “low tautomerized” adenine, see Fig. I in TURIAN (2001a,b).

This model therefore involves the median, symmetric insertion of the hydroxylated precursors of ribose between the two acid-opened poles ^+N and P^+ of the adenine phosphoramide. These $\text{C}_1 \rightarrow \text{C}_2 + \text{C}_3$ compounds could secondarily condense into the pentose C_5 according to the following symmetric-sequence of ribosylation:



Of evolutive significance, it should be emphasized that the two high-energy bonds generated by a gradient of protons in the biotic synthesis of ATP from AMP were automatically present in prebiotic adenine triphosphate which incorporates the two terminal P anhydride bonds of its cyclic triphosphate precursor (see Fig. 1).

CONCLUSION

To the question "How to make a nucleotide?", be it ATP or a monomer of RNA, it has been theoretically and experimentally answered by *asymmetric* processes, either covalent glycosylation of the nucleobases on ribose (poly)phosphate (KORNBERG, 1954-55, PITSCHE et al., 1995; MOJZSIS et al., 1999) or phosphorylation by trimetaphosphate of first heat-produced ribonucleosides, as pioneered by LOHRMANN (1977) and ETAIX & ORGEL, (1978). Recently, ZUBAY & MUI (2001) have favored the first alternative according to which ribose was probably phosphorylated prior to becoming linked to the nucleobase in such a way that "the nucleoside formation step was bypassed in favor of nucleotide formation".

However, in the early prebiotic conditions, when ribose was hardly available, nucleotide synthesis such as that of ATP, would necessarily have occurred in a two-step process modelized above: first by the bypass of a direct phosphoramide bonding of adenine and phosphate groups, secondarily, by *symmetric* insertion of ribose precursors.

RÉSUMÉ

Adénine phosphoramide comme possible précurseur de l'ATP ? - La liaison phosphoramide de l'adénine triphosphate, ouverte par acidification, pourrait être pontée par des précurseurs amphihydroxylés du ribose qui la ferait évoluer en adénosine triphosphate.

Mots-clés: phosphoramide, adénine, triphosphate, ribose, adénosine.

ACKNOWLEDGEMENTS

We are very grateful to Dr P.-Y. Morgantini (Physical Chemistry Lab., Prof. J. Weber) for the graphics modeling, Prof. Reto J. Strasser (Director Bioenergetics-Microbiology Lab.) for reading the manuscript, and Ms M. L. Manelli for her typing skills.

REFERENCES

- BALTSCHEFFSKY, H., 1993. Chemical origin and early evolution of biological energy conversion. In: Ponnamperuma, and J. Chela-Flores (eds). Chemical evolution. Hampton, Virginia: A. Deepak Publishing.
- BRACK, A. (Ed.). 1998. The Molecular Origins of Life, Cambridge University Press, Cambridge, U.K.
- CORBRIDGE, D.E.C. 1978. Phosphorus. An Outline of its Chemistry, Biochemistry and Technology. Elsevier Sci. Publ. Co.
- ETAIX, E. & L.E. ORGEL. 1978. Phosphorylation of nucleosides in aqueous solution using trimetaphosphate: formation of nucleoside triphosphates. *J. Carboh. Nucleosides-Nucleotides* 5: 91-110.
- JOYCE, G.F. & L.E. ORGEL. 1999. Prospects for understanding the origin of the RNA world. In: The RNA World, 2nd edit., Chapt 2: 49-77. Gesteland, R.F., T.R. Cech & J.F. Atkins, Cold Spring Harbor, Labor. Press.
- JOYCE, G.F. 1989. RNA evolution and the origin of life. *Nature* 338: 217-224.
- KORNBERG, A. et al., 1954-55. In J. N. Davidson, The Biochemistry of the Nucleic Acids, 1957, Methuen & Co Ltd., London.
- LAHAV, N. 1999. Biogenesis: Theories of Life's Origins. Oxford University Press.

- LIPMANN, F., 1965. Projecting backward from the present stage of evolution of biosynthesis. In: S.W. Fox (ed.) *The origin of prebiological systems and of their molecular matrices*. New York: Academic Press. pp. 212-226.
- LOHRMANN, R. & L. E. ORGEL. 1973. Prebiotic Activation Processes. *Nature* 244: 418-420.
- LOHRMANN, R. 1977. Formation of Nucleoside-5'-Phosphoramidates under Potentially Prebiological Conditions. *J. Mol. Evol.* 10: 137-154.
- MOJZSIS, S. J., R. KRISHNAMURTI & G. ARRHENIUS, 1999. Before RNA and after: Geophysical and geochemical constraints on molecular evolution. 1: 1-47 in *The RNA World*, 2nd edit., Gesteland, R.F., T.R. Cech & J.F. Atkins, Cold Spring Harbor, Labor. Press.
- ORÓ, J., 1960. Synthesis of adenine from ammonium cyanide. *Biochem. Biophys. Res. Commun.* 2: 407-412.
- PITSCH, S., R. KRISHNAMURTHY, M. BOLLI, S. WENDEBORN, A. HOLZER, M. MINTON, C. LESUEUR, I. SCHLÖNGOVT, B. JAUN & A. ESCHENMOSER. 1995. Pyranosyl-RNA ('p-RNA'): Base-pairing selectivity and potential to replicate. *Helvetica Chimica Acta* 78: 1621-1635.
- SCHWARTZ, A.W. 1998. Origins of the RNA world. In: *The Molecular Origins of Life*. Pp. 237-254. A. Brack (Ed.), Cambridge University Press, Cambridge, U.K. in A. Brack (ed.), *The Molecular Origins of Life*, Cambridge University Press, Cambridge, U.K.
- SHAPIRO, R. 1988. Prebiotic ribose synthesis: A critical analysis. *Origins of Life Evol. Biosphere* 18: 71-85.
- TURIAN, G. 2001a. Phosphoramide-bonded polynucleobase phosphates as riboseless precursors of RNA? J. Chala-Flores *et al.* (Eds). *First Steps in the Origin of Life in the Universe*. 107-110. Kluwer Acad. Publ., Netherlands.
- TURIAN, G. 2001b. Prenucleic nucleobase phosphoramidates in the transition pre- RNA - RNA ? *Archs Sci. Genève* 54: 239-247.
- TURIAN, G. & E. RIVARA-MINTEN. 2001. Prebiotic phosphoramidation of nucleobases by Mg²⁺-triggered decyclization of trimetaphosphate. *Archs. Sci. Genève*, 54: 233-238.
- TURIAN, G., E. RIVARA-MINTEN E. & A. CATTANEO. 1999. Further ³¹P-NMR evidence of phosphoramide bonding of nucleobases by Mg²⁺ enhanced nucleophilic attack on cyclic triphosphate. *Archs Sci. Genève* 52: 209-216.
- ZUBAY, G. Y. & T. MUI. 2001. Prebiotic synthesis of nucleotides. *Orig. of Life*. 31(2): 87-102.