

**Zeitschrift:** Archives des sciences et compte rendu des séances de la Société  
**Herausgeber:** Société de Physique et d'Histoire Naturelle de Genève  
**Band:** 56 (2003)  
**Heft:** 3

**Artikel:** Descriptions of three new species and two new records of Raphignathus dugés (Acari: Raphignathidae) from Turkey  
**Autor:** Doan, Salih  
**DOI:** <https://doi.org/10.5169/seals-740437>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 16.09.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

Archs Sci. Genève	Vol. 56	Fasc. 3	pp. 143-153	Décembre 2003
-------------------	---------	---------	-------------	---------------

## DESCRIPTIONS OF THREE NEW SPECIES AND TWO NEW RECORDS OF *RAPHIGNATHUS* DUGÉS (ACARI: RAPHIGNATHIDAE) FROM TURKEY

BY

**Salih DOĞAN**

*(Ms. received the 20.11.2003, accepted 19.12.2003)*

### ABSTRACT

Seven *Raphignathus* species are reported from Bursa, Çorum, Denizli, Erzincan, Erzurum, Giresun and Kırıkkale, Turkey. *Raphignathus erzincanica*, *R. giresuniensis* and *R. vahiti* are described and illustrated as new species. *Raphignathus atyeoi* Meyer and Ueckermann, previously known only from southern Africa and *R. hecmatanaensis* Khanjani and Ueckermann known only from Iran, are now also recorded from Turkey. New localities for *Raphignathus collegiatus* Atyeo, Baker and Crossley and *R. gracilis* (Rack), from Turkey are given. A key to the species of Turkey is provided.

**Key-words:** Acari, Raphignathidae, *Raphignathus*, new species, new records, Turkey.

### INTRODUCTION

The genus *Raphignathus* has a worldwide distribution containing more than 50 described species to date. They are found on tree barks, litter, moss, lichens, soil, in stored products, house dust, bird's nests, intestine of a wedded seal and the urine of a human (FAN & YIN, 2000; KHANJANI & UECKERMANN, 2003). This genus in Turkey is represented by four species: *Raphignathus collegiatus* Atyeo, Baker and Crossley, *R. gracilis* (Rack), *R. fani* Doğan and Ayyıldız and *R. kuznetzovi* Doğan and Ayyıldız (KOÇ & AYYILDIZ, 1996; DOĞAN & AYYILDIZ, 2003). Seven species are dealt with here, including three new species: *Raphignathus atyeoi* Meyer and Ueckermann, *R. hecmatanaensis* Khanjani and Ueckermann, *R. collegiatus*, *R. gracilis*, *R. erzincanica* n. sp., *R. giresuniensis* n. sp. and *R. vahiti* n. sp. Of these; *Raphignathus atyeoi* and *R. hecmatanaensis*, are new for Turkey, *Raphignathus collegiatus* and *R. gracilis* are recorded from new localities.

Dorsal and leg setal designations follow KETHLEY (1990), MEYER & UECKERMANN (1989) and GRANDJEAN (1944). All measurements are given in micrometers ( $\mu\text{m}$ ). Length of idiosoma excluding gnathosoma and width of idiosoma are measured at the widest breadth of idiosoma. Length of legs is measured from tip of claw to base of coxae. Length of palpi is measured from tarsi to base of trochanters. Known and type materials are deposited in the Zoological Museum of Ataturk University, Erzurum, Turkey.

### *Raphignathus* Dugés

**Type species** – *Raphignathus ruberrimus* Dugés, 1834.

Cheliceral bases fused to form a conical stylophore; peritremes arising from mid-basal part of stylophore, projecting to anterior margin of idiosoma; palptibial claw small; subcapitulum with 2 pairs of subcapitular setae and 2 pairs of adoral setae; podosoma with 3 shields, opisthosoma with a large shield; dorsum with 11-12 pairs of setae; one pair of eyes on lateral propodosomal shields; 2 pairs of aggenital setae, genital and anal shields each with 3 pairs of setae.

#### Key to the Turkish species of *Raphignathus*

- 1 Interscutal membrane dorsomedially with 1 pair of setae ..... 2
- Interscutal membrane dorsomedially with 2 pairs of setae ..... 4
- 2 Dorsum without small shields behind median prodorsal shield  
..... *R. fani* Doğan & Ayyıldız, 2003
- Dorsum with small shields behind median prodorsal shield ..... 3
- 3 Setal formula of femora 6-5-3-2 ..... *R. kuznetzovi* Doğan & Ayyıldız, 2003
- Setal formula of femora 6-6-4-4 .. *R. collegiatus* Atyeo, Baker & Crossley, 1961
- 4 Femur IV with 2 setae ..... 5
- Femur IV with 3 setae ..... 6
- 5 Palpfemur with 2 setae, coxa II with 2 setae ..... *R. erzincanica* n. sp.
- Palpfemur with 3 setae, coxa II with 1 seta  
..... *R. atyeoi* Meyer & Ueckermann, 1989
- 6 Palpfemur with 2 setae ..... 7
- Palpfemur with 3 setae ..... *R. giresuniensis* n. sp.
- 7 Dorsum without small shields behind median prodorsal shield ..... 8
- Dorsum with small shields behind median prodorsal shield  
..... *R. hecmatanaensis* Khanjani & Ueckermann, 2003
- 8 Setae  $e_1$  reach to opisthosomal shield,  $d_1-d_1 < e_1-e_1$  .... *R. gracilis* (Rack, 1962)
- Setae  $e_1$  not reaching opisthosomal shield, the distances between  $d_1-d_1$  and  $e_1-e_1$  subequal ..... *R. vahiti* n. sp.

#### *Raphignathus collegiatus* Atyeo, Baker & Crossley

*Raphignathus collegiatus* Atyeo, Baker & Crossley, 1961: 17; Atyeo, 1963: 185; Kuznetzov, 1976: 44; Zaher & Gomaa, 1979: 198; Vainstein & Kuznetzov, 1978: 150; Kuznetzov & Petrov, 1984: 100; Meyer & Ueckermann, 1989: 42; Fan & Yin, 2000: 91; Doğan & Ayyıldız, 2003: 146; Khanjani & Ueckermann, 2003: 305.

*Raphignathus guiyuanensis* Hu, Jing & Liang, 1995: 21.

**Material examined** – One female from moss, Güney, Denizli, 05 March 2002.

**Distribution** – China, Egypt, Iran, former USSR, Turkey and USA (ATYEO *et al.*, 1961; ATYEO, 1963; KUZNETZOV, 1976; VAINSTEIN & KUZNETZOV, 1978; ZAHER & GOMAA, 1979; KUZNETZOV & PETROV, 1984; MEYER & UECKERMANN, 1989; HU *et al.*, 1995; FAN & YIN, 2000; DOĞAN & AYYILDIZ, 2003; KHANJANI & UECKERMANN, 2003).

***Raphignathus gracilis* (Rack)**

*Acheles gracilis* Rack, 1962: 281

*Raphignathus gracilis* (Rack) Atyeo, 1963: 181; Gerson, 1968: 434; Vainstein & Kuznetzov, 1978: 150; Zaher & Gomaa, 1979: 198; Ehara, 1980: 248; Kuznetzov, 1976: 43; Charlet & McMurtry, 1977: 200; Kuznetzov & Petrov, 1984: 99; Meyer & Ueckermann, 1989: 39; Koç & Ayyıldız, 1996: 210; Khaustov & Kuznetzov, 1997: 80; Fan & Yin, 2000: 90; Kazmierski, 2000: 324; Mehrnejad & Ueckermann, 2001: 6.

**Materials examined** – Two females from soil under *Astragalus* sp., Akveren Village, Hinis, Erzurum, 19 March 2000; twelve females from moss, Ahmediyeh, Erzincan, 26 May 2001; one female from moss, Güney, Denizli, 05 March 2002; six females from moss, Åskale, Erzurum, 05. VIII. 2002.

**Distribution** – China, Egypt, former USSR, Germany, Iran, Israel, Japan, Poland, South Africa, Turkey and USA (RACK, 1962; ATYEYO, 1963; GERSON, 1968; KUZNETZOV, 1976; CHARLET & McMURTRY, 1977; VAINSTEIN & KUZNETZOV, 1978; ZAHER & GOMAA, 1979; EHARA, 1980; KUZNETZOV & PETROV, 1984; MEYER & UECKERMANN, 1989; KOÇ & AYYILDIZ, 1996; KHAUSTOV & KUZNETZOV, 1997; FAN & YIN, 2000; KAZMIERSKI, 2000; MEHRNEJAD & UECKERMANN, 2001).

***Raphignathus atyeoi* Meyer & Ueckermann  
(Figs 1-4)**

*Raphignathus atyeoi* Meyer & Ueckermann, 1989: 32.

**Materials examined** – Five females from soil under *Astragalus* sp., Akveren Village, Hinis, Erzurum, 19 May 2000; one female from soil, Akbayır Village, Hinis, Erzurum, 19 May 2000.

**Differential diagnosis** – This species can be readily distinguished from other known species in this genus in that median shield bearing or not bearing  $c_1$ , coxisternal shields absent, coxa II with one seta, legfemur IV with two setae and palpifemur with three setae. In all Turkish specimens dorsum without small shields behind median prodorsal shield.


**Distribution** – South Africa (MEYER & UECKERMANN, 1989) and Turkey (current paper).

***Raphignathus hecmatanaensis* Khanjani & Ueckermann  
(Figs 5-6)**

*Raphignathus hecmatanaensis* Khanjani & Ueckermann, 2003: 301.

**Materials examined** – Fifty-one females from soil in the nut grove, Delice, Kırıkkale, 16 May 2002; one female from moss, Güney, Denizli, 05 March 2002.

**Differential diagnosis** – This species closely resembles *R. summersi* Robaux, 1976 but differs in that (1) the setae  $h_3$  on opisthosomal shield, (2)  $e_1$  not reaching opistho-



FIGS 1-2.

*Raphignathus atyeoi* Meyer & Ueckermann (female) – 1. Dorsal view, 2. Ventral view.


somal shield and (3) only median shields partly striated. This species also resembles *R. aciculatus* Fan, 2000 but differs in that the palp femur bears two setae. General features of Turkish specimens like that given by KHANJANI & UECKERMANN (2003) but setae  $c_1$  not reaching bases of setae  $d_1$ .

**Distribution** – Iran (KHANJANI & UECKERMANN, 2003) and Turkey (current paper).

*Raphignathus erzincanica* n. sp.  
(Figs 7-8)

**FEMALE** – Length of body 360, width 180.

**Gnathosoma** – Length of gnathosoma 60. Ventral with two pairs of adoral setae and two pairs of subcapitular setae, about equal in length, 34. Distances between  $n-n$  37,  $m-m$  16,  $n-m$  10. Dorsal stylophore with striae, 54 long, movable digits 33 long. Palpi 110, number of setae and solenidion on palpi (femur to tarsus): 2, 2, 3 + 1 claw, 4 + 1  $\omega$  + 4 eupathidia.



FIGS 3-4.

*Raphignathus atyeoi* Meyer & Ueckermann (female) – 3. Leg I, 4. Legfemur IV.

**Dorsum** – With one median and two lateral shields, without small shields behind median prodorsal shield. Opisthosoma with one opisthosomal shield. Median shield bearing three pairs of setae. Lateral shields each bearing three pairs of setae, one pair of eyes and one pair of cupules. Four pairs of setae and one pair of cupules on opisthosomal shield. Two pairs of setae,  $d_1$  and  $e_1$ , and one pair of cupules, located on interscutal membrane. All dorsal shields punctuated and only median shield finely and partly striated. Body surface striated between propodosomal and opisthosomal shields. Dorsal setae simple. Dimensions of setae as follows:  $vi=ve=sce\ 30$ ,  $sci=c_2=f_1=h_1=h_2=h_3\ 27$ ,  $c_1=d_1=e_1\ 23$ . Distance between setae:  $vi-vi\ 17$ ,  $sci-sci\ 38$ ,  $vi-sci\ 36$ ,  $ve-sce\ 20$ ,  $ve-c_2\ 57$ ,  $sce-c_2\ 45$ ,  $c_1-c_1\ 17$ ,  $sci-c_1\ 80$ ,  $d_1-d_1\ 64$ ,  $e_1-e_1\ 70$ ,  $d_1-e_1\ 63$ ,  $f_1-f_1\ 40$ ,  $f_1-h_1\ 50$ ,  $h_1-h_1\ 16$ ,  $h_2-h_2\ 44$ ,  $h_1-h_2\ 17$ ,  $h_3-h_3\ 67$  and  $h_2-h_3\ 10$ .


**Venter** – Venter striated, with coxisternal shield adjacent to coxae III and IV. Genital and anal shields each are bearing three pairs of setae. A pair of cupules located laterally on genital shields.

**Legs** – Leg I 323, leg II 267, leg III 277, leg IV 353. Number of setae (solenidia in parentheses) on leg segments as follows: tarsi 21(φp,ω)-16(ω)-14(ω)-13, tibiae 6(φp)-6(φp)-6(φp)-5(φp), genua 6(k)-6(k)-4-4, femora 6-5-3-2, trochanters 1-1-2-1, coxae 2-2-2-1.

**MALE** – Unknown.

**Type material** – Holotype female from moss, AhmedİYE, Erzincan, 26 May 2001.

**Etymology** – The species name refers to the type locality, Erzincan.



FIGS 5-6.


*Raphignathus hecmatanaensis* Khanjani & Ueckermann (female) – 5. Dorsal view, 6. Ventral view.

**Differential diagnosis** – This new species is similar to *R. hexeris* Chaudhri, Akbar and Rasool, 1979, *R. scutatus* Kuznetzov, 1976, *R. rarus* Kuznetzov, 1976, *R. tumidus* Kuznetzov, 1976, *R. atyeoi* Meyer and Ueckermann, 1989, *R. africanus* Meyer and Ueckermann, 1989 and *R. kamiensis* Meyer and Ueckermann, 1989 in that leg femur IV bears two setae and two pairs setae on interscutal membrane but can be distinguished from affine species by the following combination of characters: (1) dorsum without small shields near podosomal shields, (2) only median shield finely and partly striated, (3) palp-femur with two setae, (4) leg femur I with six setae and (5) leg femur II with five setae.

*Raphignathus giresuniensis* n. sp.  
(Figs 9-10)

**FEMALE (Holotype)** – Length of body 340, width 170.

**Gnathosoma** – Length of gnathosoma 60. Ventral with two pairs of adoral setae, two pairs of subcapitular setae, about equal in length 37, distances between *n-n* 37, *m-m*




FIGS 7-8.

*Raphignathus erzincanica* n. sp. (female) – 7. Dorsal view, 8. Ventral view.

17, n-m 9. Dorsal stylophore with striae, 57 long, movable digits 37 long. Palpi 107, of setae and solenidion on palpi (femur to tarsus): 3, 2, 3 + 1 claw, 4 + 1 ω + 4 eupathidia.

**Dorsum** – Propodosoma with one median and two lateral shields separated by striae, without small shields behind median prodorsal shield. Opisthosoma with a large opisthosomal shield. Median shield bearing three pairs of setae. Lateral shields bearing three pairs of setae, one pair of eyes and one pair of cupules. Four pairs of setae and one pair of cupules on opisthosomal shield. Two pairs of setae,  $d_1$  and  $e_1$ , and one pair of cupules located on interscutal membrane. All dorsal shields punctuated and finely striated. Body surface striated between propodosomal and opisthosomal shields. Dorsal setae simple. Dimensions of setae as follows:  $vi=ve=sci=sce=f_1=h_1=h_2=h_3$  27,  $c_1=c_2=d_1=e_1$  24. Distance between setae:  $vi-vi$  20,  $sci-sci$  43,  $vi-sci$  33,  $ve-sce$  23,  $ve-c_2$  53,  $sce-c_2$  33,  $c_1-c_1$  17,  $sci-c_1$  76,  $d_1-d_1$  63,  $e_1-e_1$  75,  $d_1-e_1$  47,  $f_1-f_1$  53,  $f_1-h_1$  31,  $h_1-h_1$  30,  $h_2-h_2$  56,  $h_1-h_2$  30,  $h_3-h_3$  77 and  $h_2-h_3$  13.

**Venter** – Setae  $1a$  near coxa I. Coxae III and IV flanked by coxisternal shields with  $3a$ . Venter striated and with three pairs of setae. Three pairs of setae on genital and anal shields. One pair of cupules located lateral to genital shields.



FIGS 9-10.

*Raphignathus gireshuniensis* n. sp. (female) – 9. Dorsal view, 10. Ventral view.

**Legs** – Leg I 313, leg II 240, leg III 257, leg IV 303. Number of setae (solenidia in parentheses) on leg segments as follows: tarsi 21( $\varphi p, \omega$ )-16( $\omega$ )-14( $\omega$ )-13, tibiae 6( $\varphi p$ )-6( $\varphi p$ )-6( $\varphi p$ )-5( $\varphi p$ ), genua 6(k)-6(k)-4-4, femora 6-5-3-3, trochanters 1-1-2-1, coxae 2-2-2-1.

**MALE** – Unknown.

**Type materials** – Holotype female and one paratype female from moss, Gülburnu Village, Espiye, Giresun, 30 August 2002.

**Etymology** – The species name refers to the type locality, Giresun.

**Differential diagnosis** – This species resembles *R. scutatus* Kuznetzov, 1976 in having two pairs of setae on interscutal membrane, the same setation on palp femur and on leg femur IV, but can be recognized by (1) tibiae III and IV bearing 6 and 5 setae, respectively, (2) tarsus II bearing 16 setae, (3) *ia* on lateral shields, (4) *im* on interscutal membrane, (5) setae *e*<sub>1</sub> not reaching opisthosomal shield and (6) dorsal shields with striae. This species also resembles *R. orientalis* Fan and Li, 1993 and *R. aciculatus* Fan, 2000 but can be distinguished by the following characters: (1) the distance between *f*<sub>1</sub>-*f*<sub>1</sub>

clearly shorter than the distance between  $e_1-e_1$ , (2) setae  $e_1$  not reaching opisthosomal shield, (3) dorsum not bearing small shields behind median propodosomal shields and (4) venter with coxisternal shields. This species further resembles *R. neogracilis* Robaux, 1976 but can be distinguished by the following characters: (1) dorsal shields with striae, (2) setae  $e_1$  not reaching opisthosomal shield, (3) coxa I without coxisternal shields, (4) opisthosomal shield without lateral extensions and (5) tarsus III with solenidion  $\omega$ .

***Raphignathus vahiti* n. sp.**  
**(Figs 11-12)**

**FEMALE (Holotype)** – Length of body 393, width 217.

**Gnathosoma** – Length of gnathosoma 53. Ventral with two pairs of adoral setae, two pairs of subcapitular setae, about equal in length 31, distances between  $n-n$  36,  $m-m$  17,  $n-m$  10. Dorsal stylophore with striae, 53 long, movable digits 33 long. Palpi 107, numbers of setae and solenidion on palpi (femur to tarsus): 2, 2, 3 + 1 claw, 4 + 1  $\omega$  + 4 eupathidia.

**Dorsum** – Covered by one median, a pair of lateral and one opisthosomal shield. No small shields behind median prodorsal shield. Median shield bearing three pairs of setae. Lateral shields reduced, bearing three pairs of setae, one pair of eyes and one pair of cupules. Four pairs of setae on reduced opisthosomal shield. Two pairs of setae and two pairs of cupules located on the interscutal membrane. Dorsal shields finely punctuated and striated. Body surface striated between podosomal and opisthosomal shields. Dorsal setae simple. Dimensions of setae as follows:  $vi=ve=sci=sce=c_2=f_1=h_1=h_2=h_3$  27,  $c_1=d_1$  20,  $e_1$  23. Distance between setae:  $vi-vi$  20,  $sci-sci$  37,  $vi-sci$  40,  $ve-sce$  23,  $ve-c_2$  60,  $sce-c_2$  43,  $c_1-c_1$  13,  $sci-c_1$  83,  $d_1-d_1=e_1-e_1$  73,  $d_1-e_1$  70,  $f_1-f_1$  39,  $f_1-h_1$  24,  $h_1-h_1$  23,  $h_2-h_2$  50 and  $h_3-h_3$  80.

**Venter** – Setae  $1a$  near coxa I. Coxae III and IV flanked by coxisternal shields, with  $3a$ . Venter striated and with three pairs of setae. Three pairs of setae on genital and anal shields. One pair of cupules located lateral to genital shields.


**Legs** – Leg I 317, leg II 263, leg III 270, leg IV 343. Number of setae (solenidia in parentheses) on leg segments as follows: tarsi 21( $\varphi p, \omega$ )-16( $\omega$ )-14( $\omega$ )-13, tibiae 6( $\varphi p$ )-6( $\varphi p$ )-6( $\varphi p$ )-5( $\varphi p$ ), genua 6( $k$ )-6( $k$ )-4-4, femora 6-5-3-3, trochanters 1-1-2-1, coxae 2-2-2-1.

**MALE** – Unknown.

**Type materials** – Holotype female and one paratype female from the hollow of *Quercus* sp., Sungurlu, Çorum, 16 May 2002; one paratype female from moss, Ahmediye, Erzincan, 26 May 2001; two paratype females from litter under *Cupressus* sp., in the campus of Uludağ University, Görükle, Bursa, 10 September 2003.

**Etymology** – The species is named in honour of Dr. Vahit Bilaloglu, Atatürk University, Erzurum.

**Differential diagnosis** – This species resembles *R. gracilis* (Rack, 1962) and *R. giselae* Meyer and Ueckermann, 1989, in having two pairs of setae on interscutal membrane, the same setation on the palpi and legs, but can be distinguished by the



FIGS 11-12.

*Raphignathus vahiti* n. sp. (female) – 11. Dorsal view, 12. Ventral view.

following combination of features: (1) lateral and opisthosomal shields clearly smaller, (2) setae  $e_1$  not reaching opisthosomal shield and (3) distances between  $d_1-d_1$  and  $e_1-e_1$  subequal.

#### ACKNOWLEDGEMENTS

I wish to offer cordial thanks to Dr. Eddie A. Ueckermann, ARC-Plant Protection Research Institute, South Africa for reviewing on the manuscript, comparing some materials and confirming my identification. I thank to Dr. Nusret Ayyıldız, Erciyes University, Turkey and Dr. Sabina Fajardo Swift, University of Hawaii, USA for critically reading this paper and checking the manuscript for use of English. I also thank to Dr. Shozo Ehara, emeritus, Tottori University, Japan for help with references.

#### REFERENCES

- ATYEO, W. T., E. W. BAKER and D. A. CROSSLEY. 1961. The genus *Raphignathus* Dugès (Acarina, Raphignathidae) in the United States with notes on the old world species. *Acarologia* 3 (1): 14–20.  
 ATYEO, W. T. 1963. New and redescribed species of Raphignathidae (Acarina) and a discussion of the chaetotaxy of the Raphignathoidea. *Journal of Kansas Entomological Society* 36 (3): 172–186.

- CHARLET, L. D. & J. A. McMURTRY. 1977. Systematics and Bionomics of Predaceous and Phytophagous Mites Associated with Pine Foliage in California. *Hilgardia* 45 (7): 173–236.
- CHAUDHRI, W. M., S. AKBAR & A. RASOOL. 1979. Studies on the predatory leaf in habiting mites of Pakistan. Rep. Proj. No. PA-ARS30, Univ. Agric. Faisalabad, Pakistan, 234 pp.
- DOĞAN, S. & N. AYYILDIZ. 2003. Mites of the genus *Raphignathus* (Acari: Raphignathidae) from Turkey. *New Zeal. J. Zool.* 30 (2): 141–148.
- EHARA, S. 1980. Illustrations of the mites and ticks of Japan. *Zenkoku Noson Kyoiku Kyokai*, pp. 248–249.
- FAN, Q-H. & L-S. LI. 1993. Descriptions of three new species of superfamily Raphignathoidea (Acari: Actinedida). *Journal of Fujian Agricultural University* 22 (3): 321–324.
- FAN, Q-H. & X-M. YIN. 2000. The genus *Raphignathus* (Acari: Raphignathidae) from China. *Systematic and Applied Acarology* 5: 83–98.
- GERSON, U. 1968. Some raphignathoid mites from Israel. *J. Nat. Hist.* 2: 429–437.
- GRANDJEAN, F. 1944. Observations sur les acariens de la famille des Stigmeidae. *Arch. Sci. Physiques et Natur.* 26: 103–131.
- HU, C-Y., Z-Q. JING & L-R. LIANG. 1995. Two new species and one new record of the genus *Raphignathus* Duges (Acari: Raphignathidae). *Journal of Suzhou Railway Teachers College* 12 (3): 21–26.
- KAŻMIERSKI, A. 2000. Prostigmatic mites (Acari: Actinedida) from the Nature Reserve Slonsk. Part I: The families Stigmeidae, Raphignathidae, Caligonellidae and Camerobiidae. *Biological Bulletin of Poznan* 37 (2): 317–325.
- KETHLEY, J. 1990. Acarina: Prostigmata (Actinedida). In: *Soil Biology Guide*. Ed. D. L. Dindal. John Wiley and Sons, New York, pp. 667–756.
- KHANJANI, M. & UECKERMANN, E. A. 2003. Two new species of the genus *Raphignathus* Dugés (Acari: Raphignathidae) from Iran. *Acarologia* 43: 299–306.
- KHAUSTOV, A. A. & N. N. KUZNETZOV. 1997. Raphignathoid Mites (Acariformes, Raphignathoidea) of North-Eastern Ukraine, with the Description of a New Species of the Genus *Caligonella*. *Vestnik Zoologii* 31: 80–83.
- KOÇ, K. & N. AYYILDIZ. 1996. Türkiye faunası için yeni iki *Raphignathus* Dugès (Acari, Prostigmata, Raphignathidae) türü. *Turk. J. Zool.* 20: 209–214.
- KUZNETZOV, N. N. 1976. Fauna of mites of the family Raphignathidae Kramer 1877. *Nauchnye Doklady Vyshei Shkoly Biologicheskie Nauki* 8: 37–44.
- KUZNETZOV, N. N. & V. M. PETROV. 1984. Predaceous mites of the Baltic region (Parasitiformes: Phytoseiidae, Acariformes: Prostigmata). *Riga Zinatne* 1–142.
- MEHRNEJAD, M. R. & E. A. UECKERMANN 2001. Mites (Arthropoda, Acari) associated with pistachio trees (Anacardiaceae) in Iran (I). *Systematic and Applied Acarology Special Publications* 6: 1–12.
- MEYER, M. K. P. & E. A. UECKERMANN. 1989. African Raphignathoidea. *Entomology Mem. Dep. Agric. Wat. Supply Repub. S. Afr.* 74: 1–58.
- RACK, G. 1962. Milben aus Taubennestern mit Beschreibung einer neuen Art, *Acheles gracilis* (Acarina, Raphignathidae). *Zool. Anz.* 168 (7–8): 275–292.
- ROBAUX, P. 1976. Observation sur quelque Actinedida (=Prostigmates) du sol d'Amérique du Nord. VII. Sur deux espèces nouvelles de Raphignathidae (Acari). *Rev. Ecol. Biol. Sol.* 13 (3): 505–516.
- VAINSTEIN, B. A. & N. N. KUZNETZOV. 1978. Identification key of soil inhabiting mites, Trombidiformes. In: *Gilyarov, M. S. ed. Moscow, Nauka. Pp. 147–171.*
- ZAHER, M. & E. A. GOMAA 1979. Three new species of the genus *Raphignathus* in Egypt (Prostigmata: Raphignathidae). *Acarologia* 21: 197–203.

