Zeitschrift:	Berichte des Geobotanischen Institutes der Eidg. Techn. Hochschule, Stiftung Rübel
Herausgeber:	Geobotanisches Institut der Eidg. Techn. Hochschule, Stiftung Rübel
Band:	42 (1972-1973)
Artikel:	Biosystematic investigations in Cardamine pratensis L.s.I : 1. Diploid taxa from Central Europe and their fertility relationships
Autor:	Urbanska-Worytkiewicz, Krystyna / Landolt, Elias
Kapitel:	2: Material and methods
DOI:	https://doi.org/10.5169/seals-377678

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 24.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Acknowledgements

A number of persons have contributed to our collections; they have mostly sampled the polyploid populations of *Cardamine pratensis*. The authors should like to express their gratitude to Dr. F. SPETA, H. RANNER, R. STEIN-WENDTNER (Linz), F. GRIMS (Taufkirchen), Dr. J. PFADENHAUER (Munich), Dr. E. SULGER-BÜEL (+), PROF. Dr. H. HESS, Dr. E. LANDOLT-STADLER, W. ENGET-SCHWILER (Zurich) as well as to our colleagues and friends from the Geobotanical Institute: E. BRÄM, J. BURNAND, P. BOLLIGER, E. BROUILLET, Dr. A. GIGON, Dr. F. GROSSMANN, Dr. M.HAUSER, A. HEITZMANN-HOFFMANN, PD Dr. F. KLÖTZLI, Ch. LEUTHOLD, CH. ROTH, H. SIGG and S. ZÜST.

The floristic information given to us by Dr. A. BECHERER (Lugano) has been of great benefit. Thanks are also due to the Directors and the Keepers of the Herbaria for loan of specimens.

Our ample material could not have been fully investigated without help of the technical staff of our Institute. Special attention is due to the gardener, Mr. E.BROUILLET and to Mrs M. JONSEN who have taken care of the collected plants and kept them in experimental garden. Technical assistance of Mrs M. SIEGL, Mrs A. HEGI and Miss U. BODMER is greatly appreciated. Drawings and graphs were faultlessly made by Miss E. BRAM to whom the authors present their sincere thanks. The photos were made by Mr. H. SIGG.

2. Material and methods

Plants for the present study have been sampled in 1969 - 1974, mostly by the authors. A number of other persons have also contributed to the collection which actually comprises 900 populations. The diploids were found in 224 populations out of which 208 (1637 plants) are dealt with in the present paper, (Tables 1-6). A relatively high percentage of the diploids doesn't exactly reflect the pattern of geographical distribution of the whole group of *Cardamine pratensis*; at the present phase of our investigations a special attention has been payed to the diploid taxa.

Plants for the investigations were chosen at random within their populations. As a representative sample, 8-15 plants were collected; subsequently they were transferred to the greenhouse and, later on, into the experimental field of the Geobotanical Institute.

The studied region was principally confined to the Alps; however, for comparative purposes, the neighbouring areas were as well included in our program. In general, the boundaries of the investigated area can be traced along the following lines: Landsberg (Bavaria) - Innsbruck - Bolzano -Lecco - Turin - Chambéry - Nantua - Besançon - Epinal - Colmar - Rottweil -Landsberg. Rather intensive collections were made within this area and one can assume that a general pattern of distribution of the diploid taxa has

44

been followed there.

As far as other parts of the Alps are concerned, some plants were sent to us from surroundings of Salzburg and from Oberösterreich; they represented higher polyploid levels and therefore are not included in the present paper. It should be noted, however, that LÖVKVIST (1956) has previously dealt with some diploid taxa from the region of Vienna and from Styria. The Swedish author has also studied an ample material of *C. pratensis* s.l. from various parts of its area of distribution.

Southwestern Alps are not represented in our collections. According to the bibliographic data, the *C. pratensis* complex is there either extremely scarce or totally absent.

Not only living plants, but also an ample herbarium material was studied in the course of the present work; numerous specimens were tested for the pollen quality. The following collections were obtained on loan; Austria: Graz (GZU), Innsbruck (IB,IBF), Klagenfurt (KL), Linz (LI) France: Grenoble (GR)

Germany: Munich (M)

Italy: Florence (FI), Genoa (GE), Padua (PAD), Pavia (PAV), Verona (VER)
Switzerland: Berne (BERN), Geneva (G), Lausanne (LAU), Lugano, Zurich (Z,
ZT,RUEB)

Yugoslavia: Ljubliana (LJU)

As far as some herbarium collections from Vienna (W, WU) and those from Turin (T) are concerned, we refer to the data of LÖVKVIST (1956).

Figures in the description of the taxa are based upon a combination of data from measurements of the herbarium material collected in natural conditions and those of the cultivated plants. Maps and lists comprise as well the data obtained from the loan specimens.

The methods used in cytological investigations as well as technical details concerning the experimental crosses are given in the respective chapters.

3. Nomenclature problems and species delimination

Six taxa have been distinguished among the diploids occurring within the studied area: Cardamine granulosa, C. Matthioli, C. udicola, C. rivularis, C. pratensis and C. nemorosa. C. udicola, very rare and particularly isolated in its geographical distribution, was but partly investigated in the

45