

Zeitschrift: Berichte des Geobotanischen Institutes der Eidg. Techn. Hochschule, Stiftung Rübel

Herausgeber: Geobotanisches Institut der Eidg. Techn. Hochschule, Stiftung Rübel

Band: 44 (1975-1976)

Artikel: Reproduction in natural triploid hybrids ($2n=24$) between *Cardamine rivularis* Schur and *C. amara* L.

Autor: Urbanska-Worytkiewicz, Krystyna

Kapitel: 2: Materials and methods

DOI: <https://doi.org/10.5169/seals-377685>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 01.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

2. Materials and methods

The material for the present study was collected during 1972-1977. Fixations were made both in the field as well as the experimental greenhouse of the Geobotanical Institute. The flower buds were fixed in acetic alcohol (1:3) with a small addition of ferric acetate and acetocarmine. Micro- and macrosporogenesis as well as the seed development after experimental crosses were mostly studied on microtome sections stained with HEIDENHAIN's heamatoxylin. In addition, some lacto-propionic squashes were investigated. The viability of pollen was tested in acetocarmine smears.

The technique used in experimental crosses was the same as previously described for diploid taxa of the *Cardamine pratensis* group (URBANSKA-WORYTKIEWICZ and LANDOLT 1974) and so were the methods used for the seed germination tests and raising of young plants.

3. Results

3.1. *Vegetative multiplication*

The triploid hybrids from Urnerboden achieve their vegetative multiplication by various means. New clones are partly formed by stolons; the most characteristic, however, are small daughter plants developing at post-flowering stages. They are observable on flowering shoots, being localized very often at every node (Fig. 3) and sometimes even at the base of inflorescence. Later on, the shoots begin to bend and eventually are laying on the ground; the daughter plants have then an ample opportunity to root in and so 4 - 5 new plants per shoot may appear, separated by a distance of about 5-6 cm from each other. Complete daughter plants develop also at the upper surface of the rosette leaves (Fig. 1); their number may reach