Zeitschrift: Berichte des Geobotanischen Institutes der Eidg. Techn. Hochschule,

Stiftung Rübel

Herausgeber: Geobotanisches Institut der Eidg. Techn. Hochschule, Stiftung Rübel

Band: 44 (1975-1976)

Artikel: Reproduction in natural triploid hybrids (2n=24) between Cardamine

rivularis Schur and C. amara L.

Autor: Urbanska-Worytkiewicz, Krystyna

Kapitel: 4: Evaluation of the breeding behaviour of the triploid hybrids

DOI: https://doi.org/10.5169/seals-377685

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 28.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

the functioning of polarized R and RA gametes in the 24chromosomic hybrids, individuals from the wild as well as those experimentally reconstructed behaving alike. Furthermore, the experiments indicate that diploid backcross plants may replace a genuine *C. rivularis* in crosses with the triploids.

4. Evaluation of the breeding behaviour of the triploid hybrids

The results obtained in the course of the present study strongly suggest a non-Mendelian segregation occurring in the 24chromosomic plants from Urnerboden. It is conceivable that two complementary phases i.e. a selective sporogenesis and a zygotic selection are involved, the former mechanism eliminating all aneuploid spores or most of them, whereas the latter one may favourize only some of euploid zygotes.

No positive proof was obtained as to the functioning of truly reduced, 12chromosomic gametes in the triploid hybrids; the 24chromosomic plants obtained in our experimental crosses might as well have arisen from an union of polarized gametes with the respective chromosome numbers 8 and 16. On the other hand, as much as 21 plants out of the total 25, might only have been formed with a part of one or both polarized gametes (Table 13).

Table 13. Plants obtained in various experimental crosses with triploid(s) used as one or both parents; the latter cases marked by a double asterisk (**).

Chromosome number	Genomic structure	Possible way of origin*	Number of plants
2n=16	RR	R + R	15
2n=24	RRA	RA + R	5
2n=24**	RRA	RA + R or R1/2A + R1/2A	5
2n=32**	RRRA	RRA + R	1
2n=32	RRRA	RRA + R	2
2n=40	RRRRA	RRA + RR	2

^{*}putative gametes of triploids listed first.

It should be kept in mind that the sexual reproduction in the triploid hybrids is apparently limited; just these limitations, however, stress the importance of an anomalous breeding behaviour as a factor of stabilizing the hybridity, structural and numerical one alike.

Formation of gametes containing whole genomes and subsequently functioning in a complementary way might fortuitously appear in any sexual hybrid; by contrast, only a few well-documented cases are known where this mechanism represents a principal factor influencing patterns of variation and evolution. For the purpose of comparison, three classic examples are characterized below.

The anomalous breeding behaviour of the Oenothera biennis group (2n=14) and, in particular, that of O. Lamarckiana was extensively investigated (DE VRIES 1917, RENNER 1917, 1925 etc.; for detailed references see CLELAND 1972). These diploid taxa are permanent heterozygotes for successive translocations on all or nearly all chromosomes of the complement. The formation of chromosome rings followed by a polarized segregation results in two differentiated genomes ("the RENNER complexes") being reassembled intact at the end of meiosis; all spores produced by a given Oenothera taxon have accordingly the same reduced chromosome number n=7, but in fact represent two genetically distinct classes, both on the male and the female side. However, the plants breed true for their specific heterozygous structure, all other combinations being eliminated in selfings or sib crosses by a system of balanced lethals that operates either in gametes (e.g. O. muricata, DARLINGTON 1937) or zygotes (e.g. O. Lamarckiana). On the other hand, crosses between various true breeding lines often produce segregating progeny in the F_1 ; one, two or four classes of progeny may appear depending whether one or both parents produce one or two kinds of functional sperms and eggs (CLEVELAND 1962). A comparable behaviour was recorded in some Oenothera species of temperate South America (HAGEN 1950, HECHT 1950), in Gayophytum heterozygum (LEWIS et al. 1958, LEWIS and SZWEYKOWSKI 1964), Gaura biennis and G. triangulata (RAVEN and GREGORY 1972), Rhoeo discolor (SAX 1931), Paeonia californica and P. Brownii (STEBBINS and ELLERTON 1939, WALTERS 1942) and partly in Isotoma petraea (JAMES 1956, 1970); however, in most of these cases it is not known whether the translocation heterozygosity is associated with balanced lethals.

The Rosa canina group represents a different aspect. Most taxa are pentaploid (2n=35). At meiosis seven bivalents and twenty-one univalents occur both in PMCs and EMCs; however, the distribution of chromosomes at I anaphase is quite different in anthers and ovules. A random segregation of univalents occurring in PMCs results mostly in aneuploid, abortive microspores; the only functioning pollen grains are those with a basic set of 7 chromosomes derived from a normal distribution of bivalents. On the other hand, a polarized segregation occurring in the ovules results in the formation of two uneven groups comprising 7 and 28 chromosomes, respectively; the only functioning macrospores are 28chromosomic. Pentaploid chromosome number is subsequently restored through the fertilization of a 28chromosomic female gamete with 7chromosomic sperm. Thus, the heterogamic representatives of the Rosa canina group breed true for their odd polyploid condition (TäCKHOLM 1922, HURST 1931, DARLINGTON 1937, FAGERLIND 1940).

Leucopogon juniperinus is, to our knowledge, the only case comparable to the Rosa canina group although a random segregation does not occur; furthermore, the pattern of microspore selection being very special, there is no reduction in the total output of a good pollen. In this heterogamic allotriploid (2n=12, the genomic constitution AAB), both PMCs and EMCs form four bivalents and four univalents. As a result of a polarized segregation at I anaphase, 8 and 4 chromosomes occur in the respective groups. A subsequent elimination of some micro- and macrospores taking place, the embryo sacs in Leucopogon always carry eight chromosomes corresponding to a genomic constitution AB, whereas the functioning microspores contain only four chromosomes representing the genome A. Fertilization restores the original chromosome number and the proportion of the parent genomes (SMITH-WHITE 1948, 1955, 1959b).

As far as the *Cardamine* hybrids are concerned, eight bivalents and eight univalents occurred rather frequently during meiosis; it may be supposed that the bivalents corresponded to the two genomes RR of *C. rivularis*, whereas the univalents represented the only genome A of *C. amara*. A polarized segregation at I anaphase resulted in two uneven groups comprising respectively eight and sixteen chromosomes; the meiotic behaviour observed in

some part of the studied material seems thus comparable both to *Leucopogon* as well as the *Rosa canina* group. However, gametes of either sex formed in the above mentioned taxa are of a single kind, whereas both polarized types of sperms and eggs i.e. R and RA apparently occur in the *Cardamine* hybrids; in this respect, the studied plants correspond to the *Oenothera Lamarckiana* group.

Recurrent patterns appearing in experimental crosses strongly suggest three categories of gametes operating in the 24chromosomic hybrids from Urnerboden: apart from the polarized gametes, unreduced RRA ones are functional. However, not all theoretically possible combinations were realized.




Fig. 41. Breeding behaviour of the triploid Cardamine hybrids compared to the Oenothera Lamarckiana type. RRRA - tetraploid found in the wild as well as obtained experimentally; RRRAA = pentaploid obtained from crosses between triploid hybrids and autoallohexaploids (URBANSKA-WORYTKIEWICZ, in press); RRRRAA = natural autoallohexaploid (URBANSKA-WORYTKIEWICZ and LANDOLT 1974, 1977, URBANSKA-WORYTKIEWICZ 1976, op. cit.). GGV and GVV = O. semigigas (STOMPS 1912, DE VRIES and BOEDIJN 1924, RENNER fide EMERSON 1935); GGVV = O. gigas (LUTZ 1907, 1908, EMERSON 1935).

Crosses between various triploid individuals resulted mostly in likewise triploid RRA plant; in addition, a single tetraploid of RRRA type was

obtained. We are inclined to think that the triploid offspring derived from polarized gametes R and RA functioning in a complementary way; on the other hand, the tetraploid plant might have been issued from the union of an unreduced RRA gamete and a polarized R one. The triploid hybrids from Urner-boden appear thus to breed true for their particular genomic constitution, bar the cases where unreduced gametes occasionally contribute to some new combinations. This unusual breeding behaviour might be double-controlled by a gametophytic and a zygotic selection, the latter mechanism comparable to the balanced lethals system operating in *Oenothera Lamarckiana* (Fig. 41). It seems therefore that several ordinarily deleterious mechanisms add up in the *Cardamine* hybrids to a system with high survival value in which plants are able not only to bred true but also to profit from maximum hybrid vigour and to multiply prolifically.

5. Discussion

The rarity of hybrids between Cardamine rivularis and C. amara in the wild apparently is a measure of an isolation occurring between the two species (URBANSKA-WORYTKIEWICZ and LANDOLT 1972). The present investigations revealed that internal barriers of the reproductive isolation play the main rôle, strong incompatibility blocks operating both at pre- and postmating stages.

Natural hybrids born from crosses at the homoploid level between sexual species usually have the same chromosome number as their parents, particularly in diploid taxa. The large triploid population at Urnerboden issued from the diploid parents represents thus an exceptional case. To some extent comparable are the peculiar "lolioid" and "festucoid" hybrids (2n=21, the genomic constitution LLF and FFL, respectively) between Lolium perenne (2n=14) and Festuca pratensis (2n=14), studied by a succession of authors (e.g. PETO 1934, JENKIN 1933, 1955, WIT 1959, 1963, 1964, WHITTINGTON and HILL 1961, ESSAD 1962, 1964, 1966, 1968, GYMER and WHITTINGTON 1973a, 1973b, 1975a, 1975b). A single triploid hybrid (2n=33) between Primula