

Zeitschrift: Bulletin of the Geobotanical Institute ETH
Herausgeber: Geobotanisches Institut, ETH Zürich, Stiftung Rübel
Band: 68 (2002)

Artikel: Research Project : odor communication and the evolution pollination syndromes in orchids
Autor: Schiestl, Florian P.
DOI: <https://doi.org/10.5169/seals-377847>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 23.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

RESEARCH PROJECT

Odor communication and the evolution of pollination syndromes in orchids

FLORIAN P. SCHIESTL

Geobotanisches Institut ETH, Zollikerstrasse 107, CH-8008 Zürich, Switzerland;
Schiestl@geobot.umnw.ethz.ch

Summary

1 Whereas most orchid species offer a food-reward for their pollinators, some, without offering any reward, mimic the signals of rewarding flowers (food deception) or females of the pollinator species (sexual deception). Floral fragrances are important for attracting pollinators in both rewarding and deceptive species and are therefore significant for the reproductive success of a plant.

2 This project will investigate, in a comparative approach, the role of floral odor in orchid pollination ecology in species representing the pollination syndromes of food reward, food deception, and sexual deception. It will use a powerful combination of analytical chemistry, electrophysiology, and behavioral experiments.

3 The following aims will be targeted: (i) Identify the floral odor compounds that are detected by the pollinator insects. (ii) Determine the variability in active and non-active odor compounds within orchid populations. (iii) Investigate the attractiveness of specific odor compounds for pollinators and the consequence of their variability for influencing pollinator behavior. (iiii) Investigate the changes in odor communication in relation to the evolution of various orchid pollination syndromes.

Keywords: electroantennographic detection, floral odor, gas chromatography, orchid pollination, pollinator behavior

Bulletin of the Geobotanical Institute ETH (2002), 68, 55–62

Introduction

Orchid pollination mechanisms range among the most intriguing insect-plant relationships. The majority of species offer a reward, typically nectar, which is exploited by a number of insect pollinators e.g. bees, butterflies and moths (Dressler 1981; Nilsson 1992; van der Cingel 1995). However, a great number of orchids are deceptive, i.e. they do not offer any food for their pollinators. So-called food-de-

ceptive species imitate a range of floral attractants, i.e. floral shape, color, and scent that are associated by the pollinators with an edible reward (Nilsson 1983, 1992; Dafni 1987, 1984, Vogel 1983; Roy & Widmer 1999). Most food-deceptive orchids are pollinated by social or solitary bees (Nilsson 1983, 1992; van der Cingel 1995). Sexually deceptive orchids mimic female sex pheromones and at-

tract only male insects, which pollinate the flowers in an attempted copulation or pre-copulatory routine (Pouyanne 1917; Kullenberg 1961; Peakall & Beattie 1996; Schiestl *et al.* 1999).

Floral odor is an important cue for many plants to attract their pollinators (Vogel 1983). Most rewarding and some food-deceptive orchids show obvious floral odor emission (Nilsson 1983; Tollsten & Bergström 1989; Knudsen *et al.* 1993; Kaiser 1993). In sexually-deceptive orchids, the odor, although not detectable for humans, is the most important cue to attract pollinator males and release mating behavior (Kullenberg 1961, Schiestl *et al.* 2000a). Although numerous studies have identified floral odor compounds in various plant species (e.g. Knudsen *et al.* 1993; Kaiser 1995), we know very little about which compounds a pollinator may actually detect (Schiestl & Marion-Poll 2001).

Floral odor influences the behavior of pollinators in multiple ways. Pollinators are attracted to flowers by their scent and color over long distances (Vogel 1983). More subtle odor cues may also be used by insects at close range to decide which flower to visit (Dobson *et al.* 1999; Schiestl & Ayasse 2001). Bees learn to associate floral cues with food reward and scent is learned faster and is more prominent than color (Bogdany 1978). Since odor enables bees to discriminate species of food-providing plants it may also play a role in maintaining flower constancy of pollinators (Waser 1996). Assortative mating in plant populations may be mediated by preferential pollinator visitation of certain odor types (Pellmyr 1996). In food-deceptive orchids, high degrees of odor variability have been found, which was interpreted as an adaptive trait for reducing the learning and avoidance of flowers by a pollinator (Ackerman 1986; Ackerman *et al.* 1997; Ferdy *et al.* 1998; Roy

& Widmer 1999). Odor variability may be maintained by negative frequency dependent selection, since pollinators will avoid a common phenotype more readily than a rare one. In rewarding orchid species, however, odor variability has also been found to be high (Patt *et al.* 1989; Tollsten & Bergström 1989; Kaiser 1993). Both in rewarding and in food-deceptive species behavioral consequences for pollinators are unknown. In the sexually-deceptive *Ophrys sphegodes* it has been demonstrated, by focusing on biologically active compounds, that intra-individual variability leads to visitation of multiple flowers of one plant by a single pollinator hence enhancing reproductive success of a plant (Ayasse *et al.* 2000).

Food reward is thought to be the ancestral mode of pollination within the orchids and food deception has probably evolved from it (Dressler 1981; Dafni 1987; Aceto *et al.* 1999). Since very little is known about functions of odor in pollinator attraction in rewarding and food-deceptive orchids, the consequences of this shift for odor communication are unknown. Sexual deception has most likely evolved from food deception (Dafni 1984; Kores *et al.* 2001). Whereas food-deceptive orchids may attract their pollinators with general floral odor compounds, a switch to sexual deception requires the emission of few specific sex pheromone compounds, like in the sexually-deceptive genus *Chiloglottis* (Schiestl *et al.* 2000b,c) or specific blends of common compounds, like in *Ophrys sphegodes* (Schiestl *et al.* 1999, 2000a). The Australian orchid genus *Caladenia* contains both food and sexually-deceptive species, which may share pollinators, which is the optimal prerequisite to investigate the changes in odor communication that accompany a switch in pollination syndrome.

Goals

This project will investigate the evolutionary ecology of odor communication within different orchid pollination syndromes by focusing on the following goals:

(1) Identification of biologically active and non-active floral odor compounds

To identify volatiles that are smelled by the pollinator insect, odor will be sampled from flowers and biologically active compounds, detected and identified. Biologically active compounds are here referred to as the ones that elicit electroantennographic responses in the olfactory receptors of an insect antenna. The following specific questions will be asked:

- Which compounds emitted from the flowers are detected by olfactory receptors in the pollinators and which are not detected?
- Is there a difference in detection by different pollinator species of one orchid species?

(2) Determination of odor variability

All samples will be analysed quantitatively to answer the following specific questions:

- What degree of odor variability can be found within a plant population?
- How does the variability differ between active and non-active compounds?
- Is there a difference in variability between orchids with different pollination syndromes?

(3) Pollinator behavior

Behavioral tests will be done to investigate the role of biologically active odor compounds in attracting orchid pollinators. Further it will be investigated how odor variability influences the pollinators' behavior when visiting rewarding or deceptive flowers. The following questions will be asked:

- Which compounds attract the pollinator?
- How is the visitation behavior of the pollinator influenced by odor variability?

(4) Evolutionary changes in odor communication

Since orchids either reward or cheat their pollinators, thereby exploiting food-seeking or mating behavior, different specialisation for attracting pollinators are likely to have evolved. However, the availability of bio-synthetic pathways for odor compounds as a pre-adaptation may have greatly influenced the evolution of different pollination modes. The comparative approach will help to test the following questions:

- Is there an evolutionary trend in odor communication between different pollination syndromes, i.e. do rewarding and deceptive species attract their pollinators with different compounds or different amounts of compounds, and are patterns of variability different between rewarding and deceptive species?
- Is there an indication that energetic costs of odor emission influenced the evolution of pollination syndromes?
- Are there any pre-adaptations for the evolution of sexual deception (i.e. the synthesis of compounds similar to sex pheromones) in food-deceptive species?

Methods

STUDY SPECIES

Gymnadenia conopsea

This rewarding species is widely distributed and easily accessible around Zürich. Its flowers show strong odor emission, which has been shown to vary considerably between individuals (Kaiser 1993). Most emitted odor compound have been identified by Kaiser (1993), which will speed up our investigations. This orchid is visited by a number of insects, but only long-tongued lepidoptera are able to access the nectar in the flower spur (van der Cingel 1995).

Orchis mascula

This species is fairly common around Zürich. Like most species of the genus *Orchis* (Dafni 1987), it is a food-deceptive orchid, which does not provide any reward for its pollinators. The most important pollinators of most *Orchis* spp. are bumble bees (*Bombus* spp.) and honey bees (van der Cingel 1995). Little is known about *Orchis* floral odor and its influence on pollinators (Nilsson 1983).

Caladenia tentaculata, *C. arenaria*, *C. gracilis*

These species grow sympatrically in the surrounding of Canberra/Australia. *C. tentaculata* (Fig. 1) is a sexually-deceptive, whereas *C. arenaria* and *C. gracilis* are food-deceptive species. Hybridisation between *C. tentaculata* and *C. arenaria* has been recorded (Bower, pers.




Fig. 1. *Caladenia tentaculata* (Orchidaceae), one of the study species (Canberra, November 2000).

comm.) and indicates that the two species share pollinators. Pollinators of these orchids occur in the surrounding of Canberra.

ANALYTICAL TECHNIQUES

Collection of odor samples

To collect highly volatile odor compounds, single flowers are placed in glass tubes. Air is pulled over the flowers by a battery-operated membrane pump. Air coming into the glass chamber is cleaned by an adsorbent tube filled with 120 mg activated charcoal (SKC). The air exits through a cartridge that contains a small amount of Porapak Q or activated charcoal (CLSA Filter), onto which the volatiles are trapped. The volatiles are eluted using pentane or carbon disulphide according to the technique described in Grob *et al.* (1984) and Schiestl *et al.* (1997). These samples are subsequently analysed on a gas chromatograph.

Low volatile odor compounds of orchid-labellum will be sampled by extracting these flower parts in pentane for 12 h, according to Schiestl *et al.* (2000). Again, gas chromatographic analyses will be performed.

Detection of biologically active compounds

In the electroantennogram (EAG) technique, developed by Schneider (1957), slow olfactory receptor potentials can be recorded from an isolated insect antenna positioned between two glass capillary micro-electrodes. By blowing puffs of various odor fractions over the antenna, their potential to elicit electroantennographic responses can be determined (Fig. 2). Combined gas chromatography and electroantennogram recording (GC-EAD) allows the analysis of multi-component volatile samples and enables the importance of single compounds to be tested (Arn *et al.* 1975; Wadhams 1990). It allows an extremely sensitive and selective detection of physiologically active compounds within a multi-component sample.




Fig. 2. Schematic representation of the GC-EAD method used to detect and identify biologically active compounds in odor samples (Schiestl & Marion-Poll 2002). Odour samples are first separated into single compounds by a gas chromatograph (GC). The effluent from the GC column is then split: half the amount of each compound is directed to a flame ionisation detector (FID), and the other half into a stimulus delivery tube. Compounds in the stimulus delivery tube remain airborne and are blown over an insect antenna. Reactions from olfactory receptors in the insect antenna (electroantennographic detector EAD), i.e. differences in electric potential between the base and the tip of the antenna, are recorded with micro-electrodes and amplified. The simultaneous recording of EAD and FID traces makes it possible to judge which compound can be detected by an insect.

Chemical identification of odor compounds

The identification of semiochemicals by means of gas chromatography – mass spectrometry (GC-MS) will be done in collaboration with Dr. Roman Kaiser (Givaudan Dübendorf).

Relevance of the project

Although floral odor has fascinated scientists for centuries (Sprengel 1793; Darwin 1862; Vogel 1990), and recent investigations have accumulated a reasonable amount of data on its chemical composition in different plant

species (e.g. Knudsen *et al.* 1993; Kaiser 1993), our knowledge of its ecological significance remains small. One problem for linking floral odor to pollination ecology is our limited knowledge about the ability of pollinators to smell specific compounds emitted by a flower, and the potential of these compounds to influence the pollinators' behavior. The proposed study is an interdisciplinary approach and will use modern analytical methods combined with electrophysiology and behavioral experiments. This combination has proven to be a powerful approach to in-

investigate floral odor and its link to pollinator behavior (summarised in Schiestl & Marion-Poll 2002). Since odor communication is important for pollinator attraction, we need to understand its mechanisms to learn about the evolution of different pollination modes, which make the orchids one of the most fascinating plant families.

Orchids are ideal for studying the evolution of floral traits (Darwin 1862) because many species have pollinia, show a high degree of specialisation in their utilisation of pollinators, and are often pollinator limited in their reproductive success (Nilsson 1992). Further, the evolution of different pollination syndromes within the family allows a comparison of floral specialisations in response to different pollinators and their behavioral patterns. The Australian genus *Caladenia* is unique in its containing both food - and sexual deceptive species, which provides the exclusive opportunity to investigate odor communication and its evolutionary transition between these two pollination syndromes within closely related species. The family of orchids also contains many threatened species, therefore, the expected data on odor variability and pollinator behavior will be useful for question related to conservation biology.

Funding

This project is funded by a grant from the Swiss Federal Institute of Technology (ETH) Zürich.

References

- Aceto, S., Caputo, P., Cozzolino, S., Gaudio, L. & Moretti, A. (1999) Phylogeny and evolution of *Orchis* and allied genera based on ITS DNA variation: Morphological gaps and molecular continuity. *Molecular Phylogenetics and Evolution*, **131**, 67–76.
- Ackerman, J.D. (1986) Mechanisms and evolution of food-deceptive pollination systems in orchids. *Lindleyana*, **1**, 108–113.
- Ackerman, J.D., Meléndez-Ackerman E.J. & Salguero-Faria J. (1997) Variation in pollinator abundance and selection on fragrance phenotypes in an epiphytic orchid. *American Journal of Botany*, **84**, 1383–1390.
- Arn, H., Städler, E. & Rauscher, S. (1975) The electroantennographic detector – a selective and sensitive tool in the gas chromatographic analysis of insect pheromones. *Zeitschrift für Naturforschung*, **30 c**, 722–725.
- Ayasse, M., Schiestl, F.P., Paulus, H.F., Löfstedt, C., Hansson, B. S., Ibarra, F. & Francke, W. (2000) Evolution of reproductive strategies in the sexually deceptive orchid *Ophrys sphegodes*: How does flower-specific variation of odor signals influence reproductive success? *Evolution*, **54**, 1995–2006.
- Bogdany, F.J. (1978) Linking of learning signals in honey bee orientation. *Behavioral Ecology and Sociobiology*, **3**, 323–336.
- Dafni, A. (1987) Pollination in *Orchis* and related genera: evolution from reward to deception. *Orchid Biology, reviews and perspectives IV* (ed. J. Arditti), pp. 79–104. Cornell University Press, Ithaca.
- Dafni, A. (1984) Mimicry and deception in pollination. *Annual Reviews of Ecology and Systematics*, **15**, 259–278.
- Darwin, C. (1862) *On the various contrivances by which British and foreign orchids are fertilized by insects*. Murray London.
- Dobson, H.E.M., Danielson E.M. & Van Wesep, I.D. (1999) Pollen odor chemicals as modulators of bumble bee foraging on *Rosa rugosa* Thunb. (Rosaceae) *Plant Species Biology*, **14**, 153–166.
- Dressler, R.L. (1981) *The orchids: natural history and classification*. Harvard University Press: Cambridge.
- Ferdy, J.B., Gouyon, P.H., Moret, J. & Godelle, B. (1998) Pollinator behavior and deceptive pollination – learning process and floral evolution. *American Naturalist*, **152**, 696–705.
- Grob, K., Grob, G. & Habich, A. (1984) Overcoming background contamination in closed loop stripping analysis (CLSA). *Journal of High Resolution Chromatography and Chromatography Communications*, **7**, 340–342.

- Kaiser, R. (1993) *The scent of orchids*. Roche, Basel.
- Kaiser, R. (1995) New or uncommon volatile compounds in floral scents. *13th International congress of flavours, fragrances and essential oils*, Istanbul, pp 1–34.
- Knudsen, J.T., Tollsten, L. & Bergström, G. (1993) Floral scents—a checklist of volatile compounds isolated by headspace techniques. *Phytochemistry*, **332**, 253–280.
- Kores, P., Molvray, M., Hopper, S., Weston, P. H., Brown, A., Cameron, K. & Chase, M. (2001) A phylogenetic analysis of Diurideae (Orchidaceae) based on plastid DNA sequence data. *American Journal of Botany*, **88**, 1903–1914.
- Kullenberg, B. (1961) *Studies in Ophrys pollination*. Zool. Bidrag. Uppsala.
- Nilsson, L.A. (1983) Anthecology of *Orchis mascula*. *Nordic Journal of Botany*, **3**, 157–179.
- Nilsson, L.A. (1992) Orchid pollination biology. *Trends in Ecology and Evolution*, **7**, 255–259.
- Patt, J.M., Merchant, M.W., Williams, D.R.E. & Meeuse D.J.W. (1989) Pollination biology of *Platanthera stricta*. *Phytochemistry*, **27**, 91–95.
- Peakall, R. & Beattie, A. J. (1996). Ecological and genetic consequences of pollination by sexual deception in the orchid *Caladenia tentaculata*. *Evolution*, **50**, 2207–2220.
- Pellmyr, O. (1996) Three pollination morphs in *Cimicifuga simplex*: incipient speciation due to inferiority in competition. *Oecologia*, **78**, 304–307.
- Pouyanne, A. (1917) La fécondation des *Ophrys* par les insectes. *Bulletin de la Société d'Histoire Naturelle l'Afrique du Nord*, **8**, 6–7.
- Roy, B.A. & Widmer, A. (1999) Floral mimicry: a fascinating yet poorly understood phenomenon. *Trends in Plant Sciences*, **48**, 325–330.
- Schiestl, F.P. & Marion-Poll F. (2002) Detection of physiologically active flower volatiles using gas chromatography coupled with electroantennography. *Molecular Methods of Plant Analysis, Volume 21: Analysis of Taste and Aroma* (eds. J.F. Jackson, H.F. Linskens & R. Inman), pp 173–198. Springer, Berlin.
- Schiestl, F.P. & Ayasse, M. (2001) Post-pollination production of a repellent compound in a sexually-deceptive orchid: a new mechanism for maximising reproductive success? *Oecologia*, **126**, 531–534.
- Schiestl, F.P., Ayasse, M., Paulus, H.F., Löfstedt, C., Hansson, B. S., Ibarra F. & Francke, W. (2000a) Sex pheromone mimicry in the Early Spider Orchid (*Ophrys sphegodes*): patterns of hydrocarbons as the key mechanism for pollination by sexual deception. *Journal of Comparative Physiology, A* **186**, 567–574.
- Schiestl, F.P., Mant, J., Peakall, R., Ibarra, F. & Francke, W. (2000b) Pollinator attraction by pheromone-mimicking compound in Australian and European sexually deceptive orchids. *Proceedings of the XXI International Congress of Entomology, Foz do Iguassu, Vol I*: pp 183.
- Schiestl, F.P., Ayasse, M., Paulus, H.F., Löfstedt, C., Hansson, B. S., Ibarra F. & Francke, W. (1999) Orchid pollination by sexual swindle. *Nature*, **399**, 421–422.
- Schiestl, F.P., Ayasse, M., Paulus, H.F., Erdmann, D. & Francke, W. (1997) Variation of floral scent emission and post pollination changes in individual flowers of *Ophrys sphegodes* subsp. *sphogodes* (MILLER). *Journal of Chemical Ecology*, **23**, 2881–2895.
- Schneider, D. (1957) Elektrophysiologische Untersuchungen von Chemo- und Mechanorezeptoren der Antenne des Seidenspinners (*Bombyx mori* L.). *Zeitschrift für Vergleichende Physiologie*, **40**, 8–41.
- Sprengel, C. K. (1793) Das entdeckte Geheimnis in der Natur, im Bau und in der Befruchtung der Blumen. *Floral Biology* (eds. D.G. Lloyd & S.C.H. Barrett), pp 3–43. Chapman & Hall, New York.
- Tollsten, L. & Bergström, J. (1989) Variation and post-pollination changes in floral odours released by *Platanthera bifolia*. *Nordic Journal of Botany*, **9**, 359–362.
- Van der Cingel, N.A. (1995) *An atlas of orchid pollination*. Balkema, Rotterdam.
- Vogel, S. (1983) Ecophysiology of zoophilic pollination. *Encyclopedia of plant physiology, New Series Vol. 12C*. (eds. O.L. Lange, P.S. Nobel, C.B. Osmond & H. Ziegler), pp 560–611. Springer, Berlin.
- Vogel, S. (1990) *The role of scent glands in pollination*. Smithsonian Institutions Libraries, Washington D.C.
- Wadhams, L.J. (1990) The use of coupled Gas Chromatography: electrophysiological techniques in the identification of insect pheromones. *Chromatography of insect hormones and pheromones* (eds A.R. McCaffery & I.D. Wilson), pp 289–298. Plenum Press, New York.

Waser, N.M. (1996) Flower constancy: definition, cause and measurement. *American Naturalist*, **127**, 593–603.

Received 21 March 2002

Revised version accepted 22 May 2002