**Zeitschrift:** Berichte der Schweizerischen Botanischen Gesellschaft = Bulletin de la

Société Botanique Suisse

Herausgeber: Schweizerische Botanische Gesellschaft

**Band:** 60 (1950)

**Artikel:** Synopsis of the genus Catalpa (Bignoniaceae) II

Autor: Paclt, Jii

**DOI:** https://doi.org/10.5169/seals-42141

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF: 21.10.2025** 

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# Synopsis of the genus Catalpa (Bignoniaceae) II<sup>1</sup>

Chapters on physiology and biochemistry by Jiří Paclt, Bratislava, Czechoslovakia

### 1. Growth

The influence of certain physical factors and chemicals on the germination of *Catalpa* has been studied by Montemartini (1914), who used seeds of two *Catalpa* species. According to this author, ether and chloroform, as well as increasing gravity have a stimulating effect on the germination of *Catalpa*. In addition, one seed was observed by Montemartini which contained two embryos.

The stimulation of the growth of Catalpa planted in association with black locust (Robinia pseudo-acacia), has been treated by Ferguson (1922) and McIntyre and Jeffries (1932). However, it appears to have been W. F. Cook (cp. Garman, 1912, p. 209), who first drew attention to that effect. The black locust increases soil fertility through the nitrifying bacteria of the root nodules. On the other hand, it is well known to European botanists, how many herbs, with the possible exception of Leguminosae, are destroyed when growing in the neighbourhood of Robinia.

#### 2. Movements

The phenomenon of thigmonasty in the stigmatic lobes of *Catalpa* was observed by authors in the second half of the last century; e.g., Clos (1869), Heckel (1874) and Meehan (1874). Because of the presence of only one vascular cylinder in the stigmata, the movements, being due to a change in turgescence, are relatively shortlived (c. 60 seconds per reaction). They are characterized by the approach and then the separation of the lobes of the stigma.

### 3. Chemical Substances

# a) Inorganic Constituents

An analysis of the leaf of  $Catalpa\ speciosa\ shows\ (McHargue\ and\ Roy, 1932)$ :

<sup>&</sup>lt;sup>1</sup> The main part, Part I, dealing with morphology and taxonomy, appears in «Blumea» (1951).

Weight of dry substance used for the analysis

			1000,0	gm	Mg		7.		5,1 gm
N	•		24,7	gm	Fe				$0.56~\mathrm{gm}$
Ash			73,0	gm	P				3,1 gm
K			14,5	gm	S			/•	2,2 gm
Na			1,2	gm	Si				3,9 gm
Ca			18,5	gm	Mn				0,1 gm

# b) Organic Constituents

Sardo (1884) described in his paper a new acid, catalpic acid. The substance, however, was later proven to be nonhomogenous by Piutti and Comanducci (1902). According to these authors it comprises a mixture of two acids, namely: the p-oxybenzoic [I] and p-oxybenzoic + protocatechuic acids [II].

(OH) 
$$C_6H_4$$
  $CO_2H$  (4,1) [I]  
(OH)  $C_6H_4$   $CO_2H$  (4,1) + (OH<sub>2</sub>)  $C_6H_3$   $CO_2H$  (1, 2, 4) + 2H<sub>2</sub>O [II]

These acids may arise chemically by hydrolysis of the heteroside catalposide (the "catalpin" of Claassen), which was first described in 1888. They represent, no doubt, the principal constituents of the aglycone of that heteroside. In addition, Plouvier (1947) discovered another heteroside which he called catalpinoside.

The components above mentioned occur more or less specifically in certain parts of *Catalpa*. There exist, however, a number of constituents which cannot be considered as specific in regard to *Catalpa*. According to Hiramoto and Watanabe (1939), these are sitosterol, p-coumaric acid and isoferulic acid and, according to Chollet (1946 a), stachyose.

It may be of further interest to note that no antibiotic principle has been found in Catalpa (Cannizzaro, 1946). On the other hand, it may be mentioned, that Catalpa, containing catalposide in various parts, shows some therapeutic characteristics such as the tonic, stimulating, antipyretic and diuretic properties, which were ascribed to it long ago by some of the earlier authors; e.g., Thunberg (1784), Descourtilz (1833) and Nuttall (1855). Whether tannin may occur in the bark of Catalpa, especially C. longissima, has yet to be determined.

Another phenomenon is the toxicity of the pollen of Catalpa. Bureau (1894) questioned this as he wrote: «D'après Nuttall [1855], le miel recueilli sur les fleurs de ce Catalpa [C. bignonioides] serait vénéneux. C'est peu probable. » Now, it is well known that the pollen of Catalpa can cause allergic hay fever. Swineford and Pipes tested 87 patients, who exhibited their major symptoms during April or May, with Catalpa pollen extract. Of these, 64 were negative, 13 reacted "weakly but definitely", and 10 had strongly positive

reactions (Swineford, 1940). The dermatitis venenata of White (1887) was also described as being due to Catalpa pollen.

# c) Anthocyanins and other Pigments

As is known, the anthocyanins are chemically related to the heterosides. They may have their origin in the oxidized chromogens which probably arise from the latter:

Heteroside 
$$+$$
 H<sub>2</sub>O (hydr.)  $+$  Hydrolase  $\rightarrow$  Chromogen  $+$  Monosaccharide (6)  $+$  Energy

Since the chromogens and anthocyanins are easily oxidized and reduced, respectively, they may both act as oxygen carriers in the life of the plant. The oxidation of chromogens, is often due to an injury, resulting in an increased penetration of oxygen into the plant tissues. It was Combes (1912) who experimented in this connection with Catalpa bignonioides. According to his observations, the petioles of that species become reddish beyond the trauma, but the respective blades do not. However, it must be noted that this is only a special case of the reaction which, generally, results in the typical change in colour of the leaves when the plant is injured.

The anthocyanins may also occur in apparently undamaged specimens. According to my observations, these pigments are often abundant in the young leaves of Catalpa ovata, C. ovata × C. bignonioides, and are present sometimes also in those of C. bignonioides. On reaching maturity, such leaves lose anthocyanins. It seems that the red colouring matter serves to make the energy from sunlight available to the immature leaves, as a result of which their internal temperature may rise by several degrees.

There exists a horticultural form, *C. erubescens* f. *purpurea*, the leaves of which conserve their dark purple colouration during a much longer period of development than those of typical forms of the species.

In order to determine the group of the colouring matters of flowers of *Catalpa*, I tested both the purple and yellow (orange) pigments occurring in the corolla tube, with the conclusion that the former belong to the anthocyanins, and the latter to the carotinoids.

## 4. Metabolism

The excess of sugars usually existing in leaves when anthocyanins are abundant may favour the metabolism of the respective plant. Also, in *Catalpa*, it is known that the form with anthocyanins (*C. erubescens* f. purpurea), exhibits higher values in assimilation than the normally green forms (Plester, 1912). On the other hand, the value of assimilation

ation of the yellow-leaved ("chlorina-blättrig") C. bignonioides f. aurea was estimated to be about one half of that of C. ovata. However, it is impossible to interpret these data properly, since the process of respiration was disregarded in Plester's experiments.

# 5. Biophysical Data

The osmotic pressure (P) of the cell sap of leaves of Catalpa (collected August  $30^{th}$  (was measured cryoscopically by Dixon (cp. Grafe, 1925):

C. bignonioides . .  $\triangle = 1,905$ ; P = 22,92 atm. C. speciosa . . .  $\triangle = 1,724$ ; P = 20,73 atm.

### Literature

- Bureau, E., 1894. Révision du genre Catalpa. Nouv. Arch. Mus., Paris (sér. 3), 6, 182.
- Cannizzaro, G., 1946. Ricerche critico sperimentali sulle cosiddette sostanze antibiotiche esistenti nei vegetali superiori. Boll. Soc. ital. Biol. sperim., 22, 894/895.
- Chollet, M.-M., 1946 a. Le stachyose dans Catalpa bignonioides. C. R. Acad. Sci., Paris, 222, 242—244.
  - 1946 b. Sur quelques constituants du *Catalpa bignonioides*. I. Catalposide et acide catalpique. Bull. Soc. Chim. biol., **28**, 668—671.
- Claassen, E., 1888. Catalpin, a bitter principle. Amer. Chem. J., 10, 328-330.
- Clos, D., 1869. L'irritabilité du stigmate est-elle un caractère physiologique ordinal des Bignoniacées? Bull. Soc. bot. France, 16, 114/115.
- Colin, H., Tanret, G., et Chollet, M.-M., 1943. Sur le catalposide, hétéroside des fruits du *Catalpa*. C. R. Acad. Sci., Paris, 216, 677.
- Combes, R., 1912. Formation de pigments anthocyaniques déterminée dans les feuilles par la décortication annulaire des tiges. Ann. Sci. natur. Bot. (sér. 9), 16 (4—6), 26, 35, 38—41, 44 et 49.
- Descourtilz, 1833. Flore pittoresque et médicale des Antilles. Ed. 2, vol. 1, 87, Paris.
- Ferguson, J. A., 1922. Influence of locust on the growth of catalpa. J. Forestry, 20, 318/319.
- Freeman, R.D., 1944. L.E. Wise, Wood Chemistry, 574, 662, 668, New York.
- Garman, H., 1912. The catalpas and their allies. Kentucky Agr. Exp. Sta. Bull., 164, 203—223.
- Grafe, 1925. Kryoskopische, osmotische Werte (usw.). Tab. biol. Hague, 1, 431.
- Heckel, E., 1874. Du mouvement dans les stigmates bilobiés des *Scrophulariaceae*, des *Bignoniaceae* et des Sésamées. C. R. Acad. Sci., Paris, **79**, 702—704.
- Hiramoto, M., et Watanabe, K., 1939. On the constituents of Catalpa ovata G. Don. J. Pharm. Soc. Japan, 59, 261—264.
- Höfker, 1939. Tabakersatz (Catalpa bignonioides). Mitt. deutsch. dendr. Ges., 52, 173.

- McHargue, J.S., and Roy, W.R., 1932. Mineral and nitrogen content of the leaves of some forest trees at different times in the growing season. Bot. Gaz., 94, 381—393.
- McIntyre, A.C., and Jeffries, C.D., 1932. The effect of black locust on soil nitrogen and growth of catalpa. J. Forestry, 30, 22—28.
- Meehan, 1874. On the movement in the stigmatic lobes of Catalpa. Proc. Amer. Assoc. Adv. Sci. (1873), 72/73.
- Montemartini, L., 1914. Note di biologia dei semi. Atti Ist. Bot. Univ. Pavia (ser. 2), 13, 213—222.
- Nuttall, T., 1855. Michaux and Nuttall, The North American Sylva. Vol. 3, 77, Philadelphia.
- Piutti, A., et Comanducci, E., 1902. Sur les acides du Bignonia Catalpa. Bull. Soc. chim., Paris (sér. 3), 27, 615—620.
- Plester, W., 1912. Kohlensäureassimilation und Atmung bei Varietäten derselben Art, die sich durch ihre Blattfärbung unterscheiden. Beitr. Biol. Pflanz., 11, 249—304.
- Plouvier, V., 1947. Sur les hétérosides du *Catalpa bignonioides* Walt. (Bignoniacée) et du *Paulownia tomentosa* C. Koch (Scrofulariacée). C. R. Acad. Sci., Paris, 224, 670—672.
- Puriewitsch, K., 1914. Untersuchungen über Photosynthese. Jahrb. wiss. Bot., 53, 210—254.
- Sardo, S., 1884. Prime ricerche sulla *Bignonia Catalpa*. Acido catalpico. Gazz. chim. ital., 14, 134—139.
- Swineford, O., 1940. Catalpa as a cause of hay fever. J. Allergy, 11, 398—401. Thunberg, C.P., 1784. Flora japonica. Lipsiae.
- White, 1887. Dermatitis venenata. Boston, Mass.