

Zeitschrift: Botanica Helvetica
Herausgeber: Schweizerische Botanische Gesellschaft
Band: 93 (1983)
Heft: 1

Artikel: Polarized acidification at germ tube outgrowth from fungal spores (Morchella ascospores, Neurospora conidia)
Autor: Turian, Gilbert
DOI: <https://doi.org/10.5169/seals-65230>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 11.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Polarized acidification at germ tube outgrowth from fungal spores (*Morchella* ascospores, *Neurospora* conidia)

par Gilbert Turian

Département de Biologie végétale, Université de Genève

Manuscrit reçu le 21 septembre 1982

Abstract

The cytosol sorted out from the mitochondria-containing cytoplasm in a polar fashion at the onset of germ tube emergence from conidia of *Neurospora* or from ascospores of *Morchella* turns pH-indicator dyes such as bromocresol green or alizarin S to a yellow acid tinge indicating a pH < 5. Such localized acidification is confirmed by the quenching of the fluorescence of acridine orange in the emerging buds and tips of germ tubes outgrowing from *Neurospora* conidia.

Three types of structural changes forming part of the same process can be recognized during germination of fungal spores: spore swelling, germ tube emergence and germ tube elongation (Allen 1965, Gottlieb 1978). However, the swelling is not a simple consequence of the initial «reawakening» hydration of the spore but very soon involves synthesis of new proteins, and should more adequately be called isometric (isotropic) growth (Bartnicki-Garcia 1981). Emergence or outgrowth of germ tube(s) would then correspond to the first stage of anisometric or anisotropic growth which is of fundamental importance for the normal developmental sequence leading to new hyphal growth of the fungus. Unfortunately, the mechanism of its inception has as yet remained elusive even though its initial event could be ultrastructurally described as the appearance of a vesicle-rich polar zone (Hemmes and Hohl 1971, Grove and Bracker 1978). This zone which becomes the so-called exclusion zone (amitochondrial) at the tips of elongating hyphae has previously been cytochemically characterized by both its reducing capacity (Turian 1978) and its relatively low pH (Turian 1979a). Such apical acidity detected with colorimetric indicators has also been found in the germ tubes outgrowing from *Neurospora* conidia and suggested to be of physiological significance (Turian 1980).

It was therefore of interest to confirm the localized acidification not only in extending the semi-vital color reactions to wider germ tubes such as those outgrown from *Morchella* ascospores, but in applying a more refined technique, that of the fluorescent probe acridine orange (Nuccitelli and Deamer 1982), to the germ tubes of *Neurospora*.

Material and Methods

Ascospores of *Morchella conica* Pers. (var. *costata* Vent.) were scratched from commercial, dry carpophores and dispersed into an aqueous, 1% solution of malt (Berthet 1964). After 8-9 h at 20°C, drops of the suspension of ascospores in early germination were flattened below coverslips; pH-indicators (10^{-3} w/v in distilled water) were added laterally to obtain gradients of concentration.

Conidia of *Neurospora crassa* were obtained from 10-15 d old slant cultures on solid Vogel's medium (1956). They were dispersed in Vogel's medium ($\sim 10^6$ /ml) and drops of the suspension were allowed to germinate on slides in humid chambers at 20°C. After 3 h incubation, the early germinating conidia were spread below coverslips and stained with the pH-indicators gradientially added as above.

As pH-indicator dyes we used bromocresol green, bromocresol purple, and alizarin yellow S (corresponding reagent for Ca^{2+} ions) while chlorophenol red served for additional checkings. The internal pH was estimated by comparison of the tinges in spores chosen in nonoverstaining zones of the color gradient with the color scales of Langeron (1934) and Drawert (1968). Color photomicrographs were made on a Wild-M20 microscope with Agfa color (400 Asa) film for *Neurospora* conidia and Fujicolor F-II film for the ascospores of *Morchella*.

The fluorescent pH-probe acridine orange (Merck products) was freshly dissolved in distilled water (10^{-4} M). One drop of it was added to one drop of conidial suspension and the flattened preparation immediately screened for color photomicrographs with Ektachrome film on a Leitz Dialux 20 epifluorescence microscope.

Results

With germinating ascospores of *Morchella* we obtained our clearest results with alizarin yellow S but could not avoid reddish precipitates in the medium (glycoprotein exsudates?). Optimal focusing on a young germ tube just emerging in a clear yellow color from a pole of an ascospore (Plate 1,1) revealed both its acidity ($\text{pH} < 5$) and low Ca^{2+} content by contrast with the red color of the less acidic but calcium-rich cytoplasm of the ascospore body.

Plate I.

Semi-vital stainings of germinating spores with pH-indicators (1-6).

A. *Morchella conica* ascospores:


1. In alizarin yellow S, emergent (arrow) yellow acid bud, $\times 1200$.
2. Bipolar germination in bromocresol purple, $\times 3000$.

B. *Neurospora crassa* conidia:

3. In alizarin yellow S, yellow germ tube (arrow), $\times 800$.
4. In bromocresol green, outgrowing acidic bud (side view), $\times 1000$.
5. Idem, frontal view, $\times 1000$.
6. Ultrastructural-topological diagram of differential pHs in a germinating conidium as indicated by bromocresol green switched to yellow ($\text{pH} < 4.5$) in the outgrowing bud, bluish-green ($\text{pH} > 5$) in the general cytosol and blue ($\text{pH} > 6$) in the mitochondria. Protons vectorially dissipated into the emerging tip (+) attract the vesicles fusing their negatively (-) charged content into the extending tip wall.

pH-fluorescence probes (7-9).

7. Fluorescence of acridine orange quenched at the presumptive germination site (arrow), $\times 1500$.
8. Idem in the germ bud, $\times 1500$.
9. Idem at the tip of the germ tube, $\times 1500$.



Bromocresol purple having a low permeability rate was slower to reveal the contrast between the yellowish germ tubes and the purplish staining of the ascospore body, especially noticeable at the periplasmic level (Plate 1,2). With bromocresol green, the ascospores appeared dark green, with a bluish hue in the peripheral layers and a greenish-yellow tinge in the outgrowing buds.

On germinating conidia of *Neurospora*, we had to add a drop of a nearly saturated solution of alizarin yellow S in distilled water to observe a sharp contrast between the yellow outgrowing germ tube and the reddish content of the conidial body (Plate 1,3). Here also, cloudy reddish precipitates formed in the dispersion fluid, probably in contact with conidial exsudates.

In bromocresol green, germ buds outgrowing from conidia stained yellow (pH ~ 4.5), in sharp contrast to the general bluish-green staining (pH > 5) of the conidial body (Plate 1,4) in which, at high focusing, a few blue mitochondrial granules could be visualized. The yellow tinge of the buds observed laterally was maintained in frontal views (Plate 1,5), a fact indicating that the switch from green to yellow was not due to an optical effect of restriction of the light absorption path in the narrow bud structures. In bromocresol purple or in chlorophenol red, the outgrowing buds appeared yellowish, in contrast to the general red hue of the conidial bodies. However, permeation of these stains through the conidial walls was generally poor and the increased concentration required, rather than improving the internal, semi-vital staining, often directly led to a lethal acidification of the whole conidial contents.

With acridine orange, the yellowish-green fluorescence conferred to the cytoplasm of the conidial bodies was repeatedly observed to be quenched to an orange tinge in the germinating structures. Such orange fluorescence due to local acidification signalled the initial cytosolic bumps (Plate 1,7), filled the young buds (Plate 1,8) and «capped» the elongating germ tubes (Plate 1,9).

Discussion

Local cytoplasmic acidification parallels germ tube outgrowth from two types of fungal spores, as shown by the yellow switch at their level of all pH-indicator dyes tested. This has been further evidenced by the use of the fluorescent pH-probe acridine orange which, as a weak base, accumulates in acidified cell compartments; there, its yellowish-green, monomer fluorescence is more or less intensively quenched to a polymeric fluorescence appearing red in some highly acidified animal cells (Plates in Nuccitelli and Deamer 1982) or orange as in our more moderately acidified fungal structures. With another fluorescent pH-probe, 4-methyl-esculetin, we had already observed quenching of the fluorescence but to darkness in yeast buds and hyphal tips of *Allomyces*, indicative of an internal pH value of not more than 5 in such emergent structures (Turian 1981); such estimation is in good agreement with the average pH value of 4.5 estimated now from the yellow tinge observed with indicator-dyes in outgrowing buds of *Morchella* and *Neurospora*. Even admitting that this low value of pH could be upgraded to 5.0 by taking into account the possible «color shift» of the indicators due to salts and protein effects (Drawert 1968), such pH remains clearly lower than those measured in a few vegetative fungal structures. They span from the still relatively acidic values of 5.8-6.0 in yeast cells as measured by nuclear magnetic resonance technique (Gillies *et al.* 1981) to 7.1 in *Neurospora* hyphae as determined with microelectrodes (Sanders and Slayman 1980).

With our simple microscopic method of the pH-indicator dyes, we have observed that the general cytoplasm, mainly the cytosol, of the sporal bodies stained bluish-green with bromocresol green (reddish-purple with bromocresol purple) corresponding to an average pH value of 5.5 while its mitochondrial granules appearing clear blue have a pH of at least 6. When compared to the values obtained with the sophisticated techniques, ours are therefore under-estimated by about 1 pH unit. However, they do not detract from the fact that the cytosol of the germ buds, stained yellow with the same reagents, is the most acid. We cannot therefore escape the main conclusion that the polarized inception of fungal germ tubes is paralleled by the onset of a downhill pH gradient spanning from the perimitochondrial cytosol of the sporal bodies to the cytosol sorted out into the expanding germ buds. A diagrammatic color representation illustrates such downhill gradiential distribution of the internal pHs in an early germinating conidium of *Neurospora* (Plate 1,6).

The first event in the inception of the pH gradient has been hypothesized as a vectorial dissipation of protons from randomly back-positioned mitochondria (Turian 1979b). Such vectorial protonation of a cytosolic zone would locally depolarize the plasmalemma, while determining the presumptive site of germ tube emergence.

Continuity of the outgrowth process would require polar sorting out of acidified cytosol from the mitochondria-containing cytoplasm self-entrained by ionic exchanges ($H^+ - K^+$ especially) through the apically depolarized plasmalemma functioning as a proton sink for the vectorially deprotonated frontal mitochondria (Turian 1980).

We are grateful to Prof. H.-J. Huggel (Laboratoire de Physiologie comparée de l'Université) for making available his Leitz microscopy facilities and to Dr. E. Kolod for advice on fluorescence techniques.

Résumé

Le cytosol séparé polairement du cytoplasme peuplé de mitochondries lors de l'initiation de l'émergence du tube germinatif à partir des conidies de *Neurospora* ou des ascospores de *Morchella* fait virer les colorants-indicateurs de pH tels que le vert de bromocrésol ou l'alizarine S à leur teinte acide jaune ($pH < 5$). Une telle acidification localisée est confirmée par le changement de fluorescence de l'orange d'acridine dans les bourgeons en émergence et les extrémités de tubes germinatifs sortant des conidies de *Neurospora*.

Literature

- Allen P.J. (1965). Metabolic aspects of spore germination in fungi. *Ann. Rev. Phytopath.* 3: 313-334.
- Bartnicki-Garcia S. (1981). Cell wall construction during spore germination in *Phycomycetes*, 533-556, in *The Fungal Spore: Morphogenetic Controls*, eds. G. Turian and H.R. Hohl, Academic Press, London.

- Berthet P. (1964). Essai biotaxonomique sur les Discomycètes. Thèse No 340, Faculté Sciences, Université de Lyon.
- Drawert H. (1968). Vitalfärbung und Vitalfluorochromierung pflanzlicher Zellen und Gewebe. Protoplasmatologia. Band II, D, 3, 749 pp. Springer Verlag, Wien.
- Gillies, R.J., K. Ugurbil, J.A. den Hollander and R.G. Shulman (1981). ^{31}P NMR studies of intracellular pH and phosphate metabolism during cell division cycle of *Saccharomyces cerevisiae*. Proc. Natl. Acad. Sci. USA 78: 2125-2129.
- Gottlieb D. (1978). The germination of fungus spores. 166 pp. Meadowfield Press, Shildon, England.
- Grove S.N. (1978). The cytology of hyphal tip growth. Chapt. 3: 28-50, in The Filamentous Fungi III, eds. J.E. Smith and D.R. Berry, Edward Arnold, London.
- Grove S.N. and C.E. Bracker (1978). Protoplasmic changes during zoospore encystment and cyst germination in *Pythium aphanidermatum*. Exp. Mycol. 2: 51-98.
- Hemmes D.E. and H.R. Hohl (1971). Ultrastructural aspects of encystation and cyst-germination in *Phytophthora parasitica*. J. Cell Sci. 9: 175-192.
- Langeron M. (1934). Précis de Microscopie, 1205 pp. Masson et Cie, Paris.
- Nuccitelli R. and D.W. Deamer (1982). Intracellular pH, its measurement, regulation, and its utilization in cellular functions. Kroc Foundation Series, Vol. 15, Alan R. Liss, Inc., New York.
- Sanders D. and C.L. Slayman (1980). Measurement of internal pH in *Neurospora* with pH-sensitive microelectrodes. Neurospora Newsletter 27: 13.
- Turian G. (1978). The 'Spitzenkörper', centre of the reducing power in the growing hyphal apices of two septomycetous fungi. Experientia 34: 1277-1279.
- Turian G. (1979a). Cytochemical gradients and mitochondrial exclusion in the apices of vegetative hyphae. Experientia 35: 1164-1166.
- Turian G. (1979b). Polarity of elongation growth generated and sustained by anisotropic distribution of the protons ejected from mitochondria into the cytosol of hyphal apices (*Neurospora* model) ? Arch. Sc. Genève 32: 251-254.
- Turian G. (1980). Germination of conidia (*Neurospora*, *Trichoderma*): evidence of an electrogenic sink of protons mitochondrially-extruded into emerging tubes. Ber. Schweiz. Bot. Ges. 90: 203-212.
- Turian G. (1981). Low pH in fungal bud initials. Experientia 37: 1278-1279.
- Vogel H.J. (1956). A convenient growth medium for *Neurospora*. Microbial Genet. Bull. 13: 42.

Prof. Dr. G. Turian
 Département de Biologie végétale
 Université de Genève
 CH-1211 Genève 4