

Zeitschrift:	Botanica Helvetica
Herausgeber:	Schweizerische Botanische Gesellschaft
Band:	93 (1983)
Heft:	2
Artikel:	Chromosome numbers, scutellarin and iridoid patterns in the genus Galeopsis L. (Labiatae)
Autor:	Wieffering, J.H.
DOI:	https://doi.org/10.5169/seals-65250

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 22.02.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Chromosome numbers, scutellarin and iridoid patterns in the genus *Galeopsis* L. (Labiatae)

by J. H. Wieffering

Laboratorium voor Experimentele Plantensystematiek, University of Leiden
(Netherlands)

Manuscript received May 24, 1983

Abstract

J. H. Wieffering 1983. Chromosome numbers, scutellarin and iridoid patterns in the genus *Galeopsis* L. (Labiatae). *Bot. Helv.* 93: 239–253. New Chromosome counts for all species of *Galeopsis* are reported (table 4). No deviations from previous counts (table 3) were found. Scutellarin was isolated from *G. pubescens* and *G. tetrahit*. A large number of specimens (table 5; fig. 1) was screened for the presence of this flavone 7-glucuronide. By including other 7-glucuronides (chrysanthemum 7-glucuronide, baicalin) in the microchemical screening a fairly satisfactory specificity of the Molisch-test for scutellarin was demonstrated. The presence or absence of scutellarin proved to be useful for the discrimination between the closely related allotetraploid taxa *G. tetrahit* (+) and *G. bifida* (–). This character is, however, of no use for the phylogenetic interpretation of these taxa. Literature pertaining to iridoid patterns of all species of *Galeopsis* (table 6, fig. 3) is critically evaluated and its bearing on the phylogenetic interpretation of the genus is discussed.

Introduction

Eight years ago we discussed the evolution and relationships within the genus *Galeopsis* L. (Wieffering and Fikenscher 1974b) making use mainly of the extensive biosystematic studies of Müntzing (1927–1943). The need of additional information in order to obtain full understanding of the differentiation within the genus was stressed. The main subject of the 1974 paper (Wieffering and Fikenscher 1974b) was an evaluation of the systematic potentialities of iridoid patterns in the leaves of the various species of *Galeopsis* (classification and nomenclature according to Townsend 1972). By comparative chromatographic investigations of leaf extracts, and on a more limited scale of root and seed extracts, the main iridoid glucosides of all 9 species were traced and tentatively identified with compounds whose structures were known in 1973 (compare table 6 and fig. 3).

Travail dédié au professeur Claude Favarger, à l'occasion de son 70^e anniversaire

It was concluded that acetylharpagide, as a leaf constituent, was typical for the species of subgenus *Ladanum* and galiridoside for those of subgenus *Galeopsis*. Moreover two unknown ester glucosides were found in the genus. "Ester Rf 0.65–0.70" (= reptoside, see also note c, table 6) was observed mainly in species of subgenus *Ladanum* and in *G. pubescens*. The so-called "bifida-Stoff" (= ajugoside, see also note d, table 6) seemed to be restricted to the leaves of *G. bifida* and *G. tetrahit* and to accumulate in the roots of all four species of subgenus *Galeopsis*. The two subgenera seemed to be connected by *G. pubescens* which belongs to subgenus *Galeopsis*, but resembles the species of subgenus *Ladanum* in several respects, including the iridoid pattern of the leaves. It was also remarked that the iridoid pattern of the leaves of *Galeopsis* are very similar to those of the leaves of *Lamiastrum* Heister ex Fabr., but are quite distinct from those of the investigated species of *Lamium* L. sensu Ball (1972) (Wieffering and Fikenscher 1974a, b).

The purpose of the present paper is threefold:

- 1). To report chromosome counts for all species of *Galeopsis*, mainly from localities in the Netherlands, Switzerland, and France and to discuss their systematic meaning.
- 2). To check the usefulness of the character "presence of scutellarin in leaves" for the taxonomy of *Galeopsis* and for the identification of the taxa within the polyploid aggregate *G. bifida* – *G. tetrahit*.
- 3). To reevaluate the systematic meaning of iridoid patterns in the genus *Galeopsis*. This became desirable after Sticher and his group (see table 6) had described the isolation and identification of a total of eleven iridoid glucoside from 4 species of *Galeopsis*.

Material and methods

Plants

Plants and/or seeds were collected in nature or received from Botanical Gardens. Seeds were sown and grown to maturity in an experimental garden. Mature specimens from all acquisitions were carefully identified and documented by voucher specimens (voucher numbers in tables 4 and 5; classification and nomenclature according to Townsend 1972). Vouchers are kept in the herbarium of the Laboratorium voor Experimentele Plantensystematiek.

Karyological techniques

Young flower buds were fixed according to Östergren and Heneen (1962) or in Carnoy's fluid (EtOH-CHCl₃-AcOH = 6:3:1 v/v/v). Aceto-carmine staining was used. Satisfactory squash preparations were documented by photographs or by transforming them into permanent mounts according to Zeilinga and Kroon (1965). However, euparal was used for mounting instead of canada balsam. Chromosome counts were accomplished from meiotic meta- or telophases. In the few cases where no satisfactory meiotic divisions could be found, mitotic divisions from actively growing parts of the flower buds were used instead.

Detection of scutellarin

Small fragments of several leaves of each plant were placed in a vial containing a few ml of 5% hydrochloric acid. This transforms the soluble salts of scutellarin into the highly insoluble free acid form (see fig. 2: carboxylic group of glucuronic acid) and thus induces crystallization of the compound. After at least 24 hours the leaf-fragments were transferred to a large drop of aqueous chloralhydrate (7 + 3) on a slide and covered with a coverslip. After clearing by gently warming the preparation is ready for microscopic examination.

The presence of scutellarin (the 7-glucuronide of scutellarein) is indicated by large, yellow spherocrystals (fig. 1). This test was also applied to *Scutellaria altissima* (known to contain scutellarin), *S. columnae* (containing baicalin (the 7-glucuronide of baicalein), fig. 2), and *S. galericulata* (containing the 7-glucuronide of chrysin, fig. 2). Since spherocrystals were formed only in *S. altissima*, the HCl-test already used by Molisch and Goldschmiedt (1901) and by Strecker (1909) must be rather specific for scutellarin. The presence of scutellarin, baicalin, and chrysin 7-glucuronide in the above mentioned species of *Scutellaria* was proven by isolation (tables 1 and 2).

Isolation and identification of flavone 7-glucuronides (reference substances)

Glucuronides (present in plants as salts) were extracted from fresh plants by boiling water and subsequently precipitated from the aqueous extracts by hydrochloric acid. Purification was performed by crystallization from methanol (98%) or from methyl cellosolve, which is a much better solvent for scutellarin than methanol. The compounds were identified by their melting points and U.V. spectroscopy (Marsh 1955; Ruygrok unpublished). Moreover scutellarin was hydrolyzed and characterized by its aglycon, scutellarein. The relevant facts are reported in tables 1 and 2.




Fig. 1. Spherocrystals of scutellarin in the leaf-epidermis of *Galeopsis tetrahit* L.




Fig. 2. Scutellarin and related glucuronides occurring in *Galeopsis* and *Scutellaria*.

Scutellarin: $R_1 = R_3 = OH$; $R_2 = H$; $R_4 = GlA$

Baicalin: $R_3 = OH$; $R_1 = R_2 = H$; $R_4 = GlA$

Chrysin glucuronide: $R_1 = R_2 = R_3 = H$; $R_4 = GlA$

Scutellarein (= 6-hydroxyapigenin): $R_1 = R_3 = OH$; $R_2 = R_4 = H$

Baicalein: $R_1 = R_2 = R_4 = H$; $R_3 = OH$

Chrysin: $R_1 - R_4 = H$

6-Hydroxyluteolin: $R_1 - R_3 = OH$; $R_4 = H$

GlA = glucuronic acid

Table 1. Yields and properties of glucuronides isolated from fresh aerial parts of some species of *Scutellaria* and *Galeopsis*

Taxon and voucher no	Amount of extracted material	mg crude compound	mg purified compound	M.P. (°C) (uncorr.)	Rf-values ^b		
					A	B	C
<i>S. altissima</i> 12099	400 g	5200	c. 3000 (S) ^a	300	.40-.45	.32-.38	.42
<i>S. columnae</i> 17840	50 g	1016	400 (B)	218	.54	.56	.61
<i>S. galericulata</i> 18319	70 g	1188	437 (CH)	221	.52	.54	.61
<i>G. tetrahit</i> 14820	10 g	—	4 (S)	300	.43	.30	.42
<i>G. pubescens</i> 12013	65 g	874	75 (S)	300	.45	.33	.42

^a (S) = scutellarin (m.p. in lit. 300 °C; “fusion instantanée” of hydrated form according to Charaux 235–240 °C).

(B) = baicalin (m.p. in lit. 223 °C).

(CH) = chrysin 7-glucuronide (m.p. in lit. 225–226 °C):

^b (A) = phenol, water-saturated; Whatman 1 paper, ascending.

(B) = borax – borid acid buffer, pH 8,6 (Tabellenboekje Kon. Ned. Chem. Ver., 18^e ed. 1962); Whatman 1 paper, ascending.

(C) = pentanol – AcOH – water = 20:12:10 (Bose and Fröst 1967); Cellulose – Fertigplatten (Merck), ascending.

Table 2. UV spectra of glucuronides isolated from *Scutellaria* and *Galeopsis*: absorption peaks^a

Source	Scutellarin	Baicalin	Chrysin 7-glucuronide
Marsh 1955	285; 335	246; 279; 314	270; 306 (shoulder)
present paper:			
<i>S. altissima</i>	286; 336	—	—
<i>S. columnae</i>	—	247; 280; 316	—
<i>S. galericulata</i>	—	—	272; 310 (shoulder)
<i>G. tetrahit</i>	286; 336	—	—
<i>G. pubescens</i>	287; 337	—	—

^a major peak (λ_{max}) *underlined*.

Table 4. Chromosome counts in the genus *Galeopsis*. Personal counts

Sub-genus	Taxon	2n ^a	Origin ^b	Voucher number ^c
<i>Ladanum</i>	<i>G. angustifolia</i> ^d	16	Switzerland, Ticino Bot. Garden	12396 8637
	<i>G. ladanum</i> ^d	16	Bot. Garden	8606, 8635, 8636, 12343
	<i>G. pyrenaica</i> ^d	16	France, Pyrénées Orientales	17076
		16	Bot. Garden	8599, 8607, 12019, 12379, 15592, 18012
	<i>G. reuteri</i> ^d	16	France, Alpes Maritimes	17217 (see Table 3)
<i>Galeopsis</i>	<i>G. segetum</i> ^d	16	Netherlands, N-Brabant	18020
		16	Bot. Garden	8605, 8608, 8638, 8639, 8640, 12012, 12017, 12022, 12380, 12381
		32	Netherlands, N-Brabant	19492
	<i>G. bifida</i> ^d	32	Netherlands, Overijssel	8758
		32	Bot. Garden	8603, 8609, 14862
		16	Netherlands, Gelderland	8755
	<i>G. pubescens</i> ^e	16	Switzerland, Ticino	8304, 12395, 12397
		16	Bot. Garden	8759, 15581, 15583
		16	Eastern Germany, Sachsen-Anhalt	17593
	<i>G. speciosa</i> ^d	16	Netherlands, Groningen	20766
		16	Netherlands, Overijssel	22872
		16	Bot. Garden	8760, 11533, 12344
		32	Italy, Pavia	15852
<i>tetrahit</i> ^e		32	Netherlands, Drenthe	12365, 12368, 14810, 14817, 14818, 14820
<i>G. tetrahit</i> ^e	32	Netherlands, Gelderland	16684	
	32	Netherlands, Groningen	12370, 12372, 21454	
	32	Netherlands, N-Brabant	19494	
	32	Netherlands, N-Holland	19491	
	32	Netherlands, Utrecht	18005	
	32	Netherlands, Zeeland	12340	
	32	Netherlands, Z-Holland	18011, 23589 (see Table 3)	
	32	Switzerland, Glarus	6466, 10566, 10581	
	32	Switzerland, Graubünden	10663, 20326	
	32	Switzerland, Neuchâtel	8105, 8299, 20455	
	32	Switzerland, St. Gallen	19858	
	32	Switzerland, Ticino	12399, 12400, 15843, 18010	

^a With the exception of two counts published earlier (see Table 3 – v.d. Brand et al.; Kliphuis and Wieffering) and a few specimens where no meiotic metaphases could be found, only meiotic metaphases from PMC's were studied. For the sake of convenience the haploid numbers were doubled.

^b Only country and provinces (cantons, départements) are given. Plants grown from seeds procured by Botanical Gardens were carefully identified.

^c Each voucher number represents a separate sample (acquisition, collection).

^d All samples, *G. bifida* 19492 excepted, gave a negative reaction for scutellarin (see also Table 5).

^e All samples, *G. tetrahit* 23589 excepted, gave a positive reaction for scutellarin (see also Table 5).

Results

New chromosome counts in Galeopsis

According to Müntzing (1930a) the genus *Galeopsis* comprises three coenospecies; *Ladanum* (all 5 ecospecies of subgenus *Ladanum*), the diploid pair *G. pubescens* and *G. speciosa* ($2n = 16$), and the tetraploid pair *G. bifida* and *G. tetrahitz* ($2n = 32$). The latter pair is to be interpreted as an allopolyploid complex consisting of two ecospecies, each made up of a rather large number of genoecodemes (= ecotypes sensu Turesson) (for terminology see e.g. Stace 1980).

Müntzing (1930a, b) found a highly sterile triploid plant among the F_2 of a hybridization experiment with *G. pubescens* and *G. speciosa*. On backcrossing with *G. pubescens* this plant produced only one viable seed. From this seed a tetraploid self-fertile plant was grown. By selfing and selecting in the descendant generations Müntzing obtained plants indistinguishable from, and interfertile with, naturally occurring *G. tetrahitz*. As early as 1932 Müntzing formulated his conviction that natural polyploids in most cases arise via unreduced gametes, especially egg cells (Müntzing 1932b p. 136 ff.). In recent years this view is rapidly gaining ground (e.g. Harlan and deWet 1975; deWet 1980; Lewis 1980b).

There remain, however, still some questions to be answered before the tetraploid complex within the genus *Galeopsis* is fully understood (Wieffering and Fikenscher 1974b). A very intriguing question concerns the possibility of a polytopic origin of both species, *G. bifida* and *G. tetrahitz*. This pattern of evolution becomes still more likely if tetraploid cytodesmes do exist within the diploid parent species. Intraspecific polyploidy is known from many plant species (e.g. Lewis 1980a) and seems to be restricted in some species to marginal populations (Sieber and Murray 1980). In any case it seemed highly desirable to extend considerably the karyological investigations of *Galeopsis* and to cover, as far as possible, new localities of all species of the genus. My results are reported in table 4 and compared with chromosome counts of other scientists which are summarized in table 3.

Tables 3 and 4 demonstrate clearly that chromosome numbers are constant in each species of *Galeopsis*. All new counts reported in table 4 confirm former counts. This makes it highly probable that there is only one way to tetraploidy in *Galeopsis*, viz. hybridization of diploid species and polyploidization by production of non-reduced gametes in F_1 - and F_2 -plants, i.e. the process already described by Müntzing. Two questions, however, remain unsolved: (1) Did *G. tetrahitz* arise only once? (2) Did *G. bifida* arise independently from *G. tetrahitz* or did it arise by ecological specialization within allopolyploid *G. tetrahitz*? Much work is still needed to procure an unambiguous answer to both questions.

Scutellarin as a genetic and taxonomic marker in Galeopsis

Molisch and Goldschmiedt (1901) described scutellarin as a new flavonoid constituent of several species of *Scutellaria*. Scutellarin is slowly hydrolyzed by strong acids to scutellarein and a sugar-like compound. The latter was later shown to be glucuronic acid (Goldschmiedt and Zerner 1910). Marsh (1955) showed scutellarin to be the 7-glucuronide of 5:6:7:4'-tetrahydroxyflavone (= scutellarein = 6-hydroxyapigenin) (fig. 2). Molisch (Molisch and Goldschmiedt 1901) described also some microchemical reactions which are highly characteristic of scutellarin (and perhaps closely related glucuronides). If leaf fragments are placed in a cold aqueous solution of a strong acid

Table 5. Occurrence of scutellarin in leaves of *Galeopsis*

Taxon	Origin	Number of tested samples (voucher number ^a)	Scutellarin test ^b
Subgenus <i>Ladanum</i>			
<i>G. angustifolia</i>	France	3	0
	Italy	1	0
	Switzerland	5	0
	Western Germany	1	0
	Bot. Garden	2	0
<i>G. ladanum</i>	France	2	0
	Switzerland	6	0
	Bot. Garden	4	0
<i>G. pyrenaica</i>	France	2	0
	Bot. Garden	8	0
<i>G. reuteri</i>	France	1	0
<i>G. segetum</i>	France	5	0
	Luxemburg	1	0
	Netherlands	2	0
	Bot. Garden	11	0
Subgenus <i>Galeopsis</i>			
<i>G. bifida</i>	Belgium ^c	2	0
	Eastern Germany	3	0
	France	1	0
	Netherlands	8	0
	Sweden	2	0
	Switzerland	2	0
	U.S.S.R. ^c	1	0
	Western Germany	2	0
	Bot. Garden	3	0
	Netherlands	1 (19492)	+ and 0
	Switzerland	1 (22315)	+ and 0
<i>G. pubescens</i>	Eastern Germany	1	+
	Italy	1	+
	Netherlands	1	+
	Switzerland	14	+
	Bot. Garden	6	+
	France	1 (237)	0
	Italy	1 (234)	0
	Switzerland	1 (12394)	0
<i>G. speciosa</i>	Austria	1	0
	Eastern Germany	2	0
	Finland	1	0
	France	1	0
	Liechtenstein	1	0
	Netherlands	9	0
	Poland	2	0
	Western Germany ^c	1	0
	Bot. Garden	5	0
	France	3	+
	Great Britain ^c	1	+
	Italy	2	+
	Netherlands	45	+
	Norway	1	+
	Poland	1	+
	Switzerland	53	+
	Western Germany	1	+
	Netherlands	1 (18003)	+ and 0
		3 (14732, 14812, 23589)	0
	Switzerland	1 (20850)	+ and 0
	Switzerland	3 (4841, 8022, 8026)	0

^a Voucher numbers given only if results were aberrant.^b Scutellarin-test according to Molisch (1901) and Strecker (1909) performed with leaf fragments of herbarium specimens: 0 = no spherocrystals; + and 0 = some leaves positive and some negative; + = many spherocrystals in all fragments examined.^c Investigated specimens loaned from Rijksherbarium (L).

(preferably HCl, 1–5%) large, yellow spherocrystals slowly appear, mainly in the lower epidermis (fig. 1). On moistening with a solution of barium hydroxide their colour changes to rust-red, and subsequent treating with iodine turns them green.

By applying these reactions, Molisch could demonstrate the presence of scutellarin in 6 species of *Scutellaria*, in *Teucrium chamaedrys* L., and in *G. tetrahit* L. but he could not detect the compound in twelve other species of Labiatae. Strecker (1909) continued the search for scutellarin. He investigated 350 plant species, including many Labiatae, but found scutellarin only in representatives of four genera of Labiatae, *Galeopsis*, *Scutellaria*, *Teucrium*, and *Thymus*. Charaux and Rabaté (1940) isolated scutellarin from leaves of *Centaurea scabiosa* L. (Compositae). Accumulation of scutellarin by *G. tetrahit*, *Scutellaria*, and some species of Compositae was confirmed by Plouvier (1963); he described its isolation from four species of *Scutellaria*, *G. tetrahit*, and from the three composites *Erigeron canadensis* L. (= *Conyza canadensis* (L.) Cronq.), *Erigeron ramosus* Britton, Stern et Pogg. (= *E. annuus* (L.) Pers. subsp. *strigosus* (Mühl. ex Willd.) Wagenitz), and *Centaurea calcitrapa* L.

Strecker (1909) observed scutellarin in *G. ladanum* L., *G. versicolor* Curtis, and *G. tetrahit* L., but not in *G. bifida* Boenn., *G. pubescens* Besser, *G. walteriana* Schlecht., and *G. neglecta* Schultes. According to Briquet (1893), Porsch (1903), and Townsend (1972) *G. versicolor* Curtis (non Spenner) is a synonym of *G. speciosa* Mill., and *G. walteriana* Schlecht. (written *walterina* by Briquet and *walterana* in Flora Europaea, vol. 3 (1972), p. 351) a synonym of *G. pubescens* Bess. *G. neglecta* Schultes probably belongs to *G. tetrahit* var. *bifida* Lejeune et Courtois (= subsp. *bifida* Fries = *G. bifida* Boenn.) (Briquet 1893) or to *G. tetrahit* L. s.s. (Porsch 1903).

The results of Molisch and Goldschmidt and of Strecker induced me to investigate more closely the character “presence of scutellarin in leaves” for the whole genus *Galeopsis*. The results of this investigation are summarized in table 5.

With regard to *G. tetrahit* my results are in complete agreement with those of Molisch (Molisch and Goldschmidt 1901) and Strecker (1909). This species accumulates large amounts of scutellarin. I also agree with Strecker with regard to the absence of scutellarin from *G. bifida*. Presence or absence of scutellarin, as detected by the very simple hydrochloric acid test, discriminates nicely between these two closely related taxa. All my other observations, however, are at variance with Strecker's. The only other species of *Galeopsis* which accumulates scutellarin is *G. pubescens*. I did not find scutellarin in any leaf sample neither of subgenus *Ladanum*, nor of *G. speciosa*. Perhaps Strecker confused *G. pubescens* and *G. ladanum* on one side and *G. versicolor* Curtis and *G. versicolor* Spenner on the other side. *G. versicolor* Spenner (non Curtis) is synonymous with *G. tetrahit*.

Accumulation of scutellarin in the leaves of the allopolyploid species *G. tetrahit* must derive from *G. pubescens*. The presence of scutellarin in the latter species was confirmed by isolation and spectroscopic and chromatographic characterization (see Material and methods and tables 1 and 2). The absence of scutellarin from the foliage of *G. bifida* can be explained in several ways which are for the time being equally plausible.

The few aberrant observations included in table 5 need some comment. *G. bifida* 19492 grew at the bank of a brooklet, facing a population of *G. tetrahit*. *G. bifida* 22315 was collected in the transition-zone between a wet and swampy nature-reserve and the neighbouring agricultural fields. Under such circumstances some hybridization may well occur and be responsible for the varying amount of scutellarin in the foliage of *G. bifida*.

G. pubescens 234 and 237 were collected in the southwestern alps at the Italian and French side of the border respectively. The two specimens are very much alike. They are not only chemically but also morphologically atypical. The stems and branches are very thin, with the nodes hardly swollen. The general shape of the leaves is about normal but the leaves are very small (less than 3 cm long) and strikingly hairy. These plants from the southwestern edge of the range of the species possibly represent a distinct taxon.

G. tetrahit 8022, 8026, and 18003 were collected in habitats where some hybridization with *G. bifida* does not seem improbable (8022 and 8026 in Switzerland along the river Areuse; 18003 in the Netherlands, at the slope to a pool near the river Rhine).

For *G. pubescens* 12394 (Ticino, Switzerland), *G. tetrahit* 14732, 14812 (Netherlands), 4841 (Bern, Switzerland), and 20850 (Ticino, Switzerland) no explanation can be offered for the deviating behaviour because their habitats were not examined by me. All these plants are typical representatives of the respective species.




Fig. 3. Iridoid glucosides (= aucubinoids) of species of *Galeopsis* (compare table 4). I. $R = H$: antirrhinoside (1), $R = \text{glucosyl}$: 5-glucosyl antirrhinoside (2); II. $R = H$: galiridoside (3), $R = \text{glucosyl}$: 5-glucosyl galiridoside (4); III. $R = H$: harpagide (5), $R = \text{COCH}_3$: acetyl harpagide (6); IV. $R = H$: 6-deoxy harpagide (7), $R = \text{COCH}_3$: reptoside (8); V. $R_1 = \text{OH}$ and $R_2 = H$: ajugol (9), $R_1 = \text{OH}$ and $R_2 = \text{COCH}_3$: ajugoside (10) (Damtoft et al. 1981); $R_1 = R_2 = H$: glueroside (11) (= 6-deoxyajugol); VI. daunoside (12).

Table 6. Iridoids in *Galeopsis*

Taxon (number of leaf samples investigated by W-F ^a)	References	Iridoid compounds ^b										Plant part
		3	5	6	8	10	reptoside ^c	ajugoside ^d	other compounds			
Subgenus <i>Ladanum</i>												
<i>G. angustifolia</i> (7)	W-F ^a	—	+	+	+	+	—	—	—	—	—	leaves
<i>G. ladanum</i> (12)	W-F	?	+	+	+	+	(+)	—	—	—	—	leaves
<i>G. pyrenaica</i> (4)	W-F	+	+	+	—	—	traces	—	—	—	—	leaves
<i>G. reuteri</i> (1) ^e	W-F	—	+	+	—	—	—	—	—	—	—	leaves
<i>G. segetum</i> (5)	W-F	+	+	+	+	+	(+)	—	—	—	—	leaves
	Junod-Busch 1976 ^e	+	+	+	+	+	+	—	—	—	—	whole plant
Subgenus <i>Galeopsis</i> (= subgen. <i>Tetrahit</i>)												
<i>G. pubescens</i> (5)	W-F	++	+	++	++	++	—	—	—	—	—	leaves
	Sticher, Rogenmoser, Weisflog 1975 ^f	++	+	++	++	++	—	—	—	—	—	aerial parts
	Rogenmoser 1975 ^g	++	+	++	++	++	—	—	—	—	—	aerial parts
<i>G. speciosa</i> (5)	W-F	++	+	++	—	—	—	—	—	—	—	leaves
<i>G. bifida</i> (8)	W-F	++	+	++	—	—	—	—	—	—	—	leaves
	Junod-Busch 1976 ^k	++	+	++	—	—	—	—	—	—	—	aerial parts
<i>G. tetrahit</i> (11)	W-F	++	+	++	—	—	traces ⁱ	—	—	—	—	leaves
	Sticher 1970a, 1970b	++	+	++	—	—	—	—	—	—	—	leaves
	Sticher, Rogenmoser, Weisflog 1975 ^h	++	+	++	—	—	—	—	—	—	—	leaves
	Sticher, Weisflog 1975 ^m	++	+	++	—	—	—	—	—	—	—	leaves
	Weisflog 1975 ^m	++	+	++	—	—	—	—	—	—	—	leaves

^a Wieffering and Fikenscher 1974b. 5–10 g fresh or dried leaves were extracted; in the case of *G. reuteri* only a few leaves from an herbarium specimen were available.

^b + + = main constituent; + = present in appreciable amounts; (+) = detected in some samples only; — = not detected. Compounds with known structure indicated by numbers (1–12); Compounds with as yet unknown structure indicated by letters (H, O) (fig. 3; Junod-Busch 1976).

^c Identical with "Ester R_f 0.65–0.70" of W-F (Sticher and Weisflog 1975; Rogemoser 1975).

^d Identical with "Bifida-Stoff" of W-F (Sticher and Weisflog 1975; Junod-Busch 1976).

^e 2.4 Kg of fresh, flowering plants, including roots, were extracted.

^f In roots (3 samples): (+).

^g 18.16 Kg of fresh aerial parts of flowering plants were extracted.

^h In roots (1 sample): 6 + ; 10 + + .

ⁱ In roots (2 samples): 6 and 10 both + + .

^k 5.6 Kg of fresh aerial parts of flowering plants from a mixed population containing c. 60% *G. bifida* and 40% *G. tetrahit* were extracted. In roots the same pattern of iridoids was found.

^l In roots (2 samples): 6 + + : 10 + to + + : in seeds (1 sample): 6 + + : 8 + + .

Concluding it may be stated that the production of spherocrystalline masses of scutellarin by immersion of leaf fragments in cold hydrochloric acid is a useful feature to characterize *G. tetrahit* and *G. pubescens*. As far as is known at present, only the 7-glucuronide of 6-hydroxyapigenin does react precisely in the same way as was described by Molisch and by Strecker for scutellarin. Scutellarein glycosides (lacking the carboxylic group of glucuronic acid) and methyl ethers of scutellarein behave otherwise. *G. segetum* (= *G. ochroleuca*) contains scutellarein 7-glucoside and 6-hydroxyluteolin 7-glucoside (Trotin and Pinkas 1979) and *G. ladanum* contains ladanein (7,4'-dimethyl ether of scutellarein) and ladanetin (7-methyl ether of scutellarein) and several glycosides of ladanein (Gritsenko and Litvinenko 1969; Gritsenko, Litvinenko, and Kovalev 1969). Both taxa give a negative scutellarin test. 6-Hydroxyapigenin (= scutellarein) and 6-hydroxyluteolin are rather frequent flavonoids in families of Tubiflorae (Harborne and Williams 1971) but they rarely occur in the form of unmethylated 7-glucuronides. Though it was shown by me that baicalin and chrysin 7-glucuronide (see fig. 2 and Material and methods) do not give the scutellarin-reaction, it may well be that 6- and 8-hydroxyluteolin, when present in a plant as the 7-glucuronide, would produce the same microchemical reactions as does scutellarin.

Iridoid patterns of Galeopsis species

Sticher and his group performed very accurate phytochemical investigations with four species of *Galeopsis*: viz. *G. segetum*, *G. pubescens*, *G. tetrahit*, and *G. bifida*. Their *G. bifida* material was probably influenced by hybridization with *G. tetrahit*. Hybrid populations between these two species are by no means rare (Müntzing 1930a, this paper p...). The results of Sticher's group are compared with our results in table 6 and illustrated by fig. 3.

In most instances our comparative studies agree well with the iridoid glucosides as isolated from four species of *Galeopsis* by Sticher's group. Of course all minor components are not detectable by the analytical procedure applied by us (table 6, note a). Thus compounds **4**, **7**, **9**, **11**, and **12** (fig. 3) which were encountered as minor components in one or more species were not trace in our study.

There is one striking discrepancy, however, in the results of the two groups. It concerns acetylharpagide which is lacking in the leaves of *G. bifida* according to our observations and present in large amounts according to Junod-Busch (1976). This affects our former conclusion that acetylharpagide as a leaf constituent is typical of species of subgenus *Ladanum* and of *G. pubescens* but lacks in the other three species of subgenus *Galeopsis*. In this respect two facts should not be forgotten. Firstly there are appreciable differences between the different parts of a plant (Wieffering and Fikenscher 1974a). The material extracted by Junod-Busch (1976) did contain besides leaves also stems, and all parts of flowers. Secondly, hybridization may affect the iridoid patterns of a given population. Three of the *Galeopsis* iridoids are acetylated compounds, i.e. acetylharpagide (**6**), reptoside (**8**), and ajugoside (**10**). Compounds (**6**) and (**8**) are more characteristic of the species of subgenus *Ladanum* and of *G. pubescens* if only leaves are examined, and (**10**) occurs in roots and leaves of species of subgenus *Galeopsis*. It is not impossible that acetylation of glucosides is affected by hybridization. Junod-Busch assumed that the glucosides (**4**), (**6**), (**8**), (**9**), and (**10**) are typical for *G. bifida* because they were not encountered by Weisflog in *G. tetrahit*. On the other hand she could not trace gluroside (**11**) in her *G. bifida* material; the latter compound is therefore assumed to show intraspecific variation (this conclusion is based on the

presence of c. 40% *G. tetrahit* plants in the extracted sample of *G. bifida*). In this respect it is interesting to retain that Weisflog (1975) extracted leaves only in his study of the iridoid compounds of *G. tetrahit*. In agreement with us he did not find acetylharpagide in the leaves of this taxon. It is not impossible that most of the acetylharpagide isolated by Junod-Busch from *G. bifida* stemmed from stems and flowers not from leaves.

Another discrepancy which deserves to be mentioned is the fact that antirrhinoside (1) and glucosylantirrhinoside (2), which were isolated by Junod-Busch (1976) from *G. segetum* in amounts equalling those of galiridoside (3) and surpassing those of reptoside (8), were not detected by us. Most probably 1 was masked on our chromatograms by 6, and 2 was overlooked because it does scarcely react with Godin's reagent. Hitherto 1 and 2, two iridoids formerly known only from Scrophulariaceae, have been traced only in one species of subgenus *Ladanum*; possibly they represent a biochemical character of this subgenus.

I think it is safe to maintain that iridoid patterns represent characters worth of further study. One should realize, however, that the patterns may vary with plant parts and that they may be affected by hybridization. Moreover it is more than likely that there is some variation within each species, especially in the highly variable taxa belonging to subgenus *Galeopsis* (Briquet 1893; Porsch 1903; Henrard 1919). Therefore much more research has to be performed before these patterns can be safely used as taxonomic and biosystematic markers in *Galeopsis*.

Conclusions

The caryological and chemical characters treated in this paper contribute to the understanding of *Galeopsis* in an evolutionary sense.

Absence of infraspecific polyploidy suggests that there is only one way to polyploidy in *Galeopsis*; repeated hybridization and production of unreduced gametes by hybridogenic plants.

Presence of scutellarin in *G. tetrahit* and *G. pubescens* and its absence in all other species shows clearly that the character was introduced in *G. tetrahit* by *G. pubescens*.

It seems that a more detailed study of iridoid compounds of all species of *Galeopsis* would be rewarding in the context of efforts undertaken to disclose the phylogenetic history of the genus in all details.

Two essential questions remain still unanswered: (1) Did *G. tetrahit* have a monotypic or a polytopic origin? (2) Is *G. bifida* an ecodeme of *G. tetrahit* or does it have an independent origin from the two diploid species of subgenus *Galeopsis*?

Zusammenfassung

Chromosomenzahlen wurden für Herkünfte aller bekannten *Galeopsis*-Arten ermittelt (Tabelle 4); es wurden keine von den bereits bekannten (Tabelle 3) abweichende Zahlen gefunden.

Scutellarin wurde aus *G. pubescens* und *G. tetrahit* isoliert. Alle *Galeopsis*-Arten und viele Herkünfte wurden mit Hilfe eines Schnelltests auf Vorkommen dieses 7-Glucuronids geprüft (Fig. 1 und 2; Tabelle 5). Da auch andere 7-Glucuronide (Fig. 2; Tabellen 1 und 2) berücksichtigt wurden, durfte die mineralsäureinduzierte Auskristallisation in

den Blattzellen (Fig. 1) als ziemlich spezifisch für Scutellarin betrachtet werden. Scutellarin lässt sich zur Unterscheidung von *G. bifida* (fehlt) und *G. tetrahit* (vorhanden) heranziehen.

Die Literatur über Iridoiglycoside der Gattung und deren mögliche taxonomische Bedeutung werden kritisch besprochen.

Auf eine mögliche Bedeutung der chemischen Merkmale für das Verständnis der phylogenetischen Zusammenhänge in der Gattung *Galeopsis* wird hingewiesen.

The author is much indebted to Prof. Dr. J.B. Harborne, Prof. Dr. O. Sticher, Dr. L.H. Fikenscher, Dr. H.W.L. Ruygrok Prof. Dr. R. Hegnauer, Miss E.G.M. Schlatmann, the directors and staff members of many Botanical Gardens for sending seed samples.

References

Ball P.W. 1972. *Lamium* L. and *Lamiastrum* Heister ex Fabr.: T.G. Tutin et al. (ed.). *Flora Europaea* 3: 147–149. Cambridge Univ. Press, London and New York.

Bose S., S. Fröst 1967. An investigation on the variation of phenolic compounds in *Galeopsis* using thin layer chromatography. *Hereditas* 58: 145–164.

Brand C. van den, F.C.M. van Meel, J.H. Wieffering 1979: Å. Löve (ed.). Chromosome number reports LXIV. *Taxon* 28: 396.

Briquet J. 1893. Monographie du genre *Galeopsis*. F. Hayez, Imprimeur de l'Académie Royale, Bruxelles. 323 pp.

Charaux C., J. Rabaté 1940. Étude de *Centaurea scabiosa* L. *Jour. Pharm. Chim. Sér. 9 I*: 155–162.

Damtoft S., S.R. Jensen, B.J. Nielsen 1981. ^{13}C and ^1H NMR spectroscopy as a tool in the configurational analysis of iridoid glucosides. *Phytochemistry* 20: 2717–2732.

DeWet J.M.J. 1980. Origins of polyploidy: W.H. Lewis (ed.). *Polyploidy, biological relevance*, pp. 3–15. Plenum Press, New York and London.

Gadella T.W.J., E. Kliphuis 1963. Chromosome numbers of flowering plants in the Netherlands. *Acta Bot. Neerl.* 12: 195–230.

Goldschmidt G., E. Zerner 1910. Über das Scutellarin. *Monatsh. Chem.* 31: 439–491.

Gritsenko E.N., V.I. Litvinenko 1969. Flavonoids from *Galeopsis ladanum*. *Khim. Prir. Soedin.* 5: 55–56. (ex *Chem. Abstr.* 1969, 71: 10279).

Gritsenko E.N., V.I. Litvinenko, I.P. Kovalev 1969. Flavonoids from the red hemp nettle (*Galeopsis ladanum*). *Dokl. Akad. Nauk Azerb. SSR* 25: 55–58. (ex *Chem. Abstr.* 1970, 73: 63168).

Hagberg A. 1952. Heterosis in F_1 combinations in *Galeopsis*. I. *Hereditas* 38: 33–82.

Harborne J.B., C.A. Williams 1971. 6-Hydroxyluteolin and scutellarein as phyletic markers in higher plants. *Phytochemistry* 10: 367–378.

Harlan J.R., J.M.J. deWet 1975. On Ö. Winge and a prayer: the origins of polyploidy. *Bot. Rev.* 41: 361–390.

Henrard J. Th. 1919. *Galeopsis*. Een systematisch-floristische studie. *Ned. Kruidk. Arch.* 1918: 158–188.

Junod-Busch U. 1976. Isolierung, Charakterisierung und Strukturaufklärung der Iridoidglucoside von *Galeopsis segetum* Necker und *Galeopsis bifida* Bönninghausen. *Diss. ETH Zürich* 5747, 113 pp.

Kliphuis E., J.H. Wieffering 1972. Chromosome numbers of some angiosperms from the south of France. *Acta Bot. Neerl.* 21: 598–604.

Küpfer Ph. 1971. Liens génétiques entre les flores alpine et pyrénéenne. *Actes des colloques sur la flore et la végétation des Chaines Alpine et Jurassienne* pp. 167–185, Paris (ex R.J. Moore 1973).

Lewis W.H. 1980a. Polyploidy in species populations. In: W.H. Lewis (ed.). *Polyploidy, biological relevance* pp. 103–144. Plenum Press, New York and London.

— 1980b. Polyploidy in plant evolution: summary. In: W.H. Lewis (ed.). *Polyploidy, biological relevance* pp. 145–147. Plenum Press, New York and London.

Loon J.C. van, B. Kieft 1980: Á. Löve (ed.). Chromosome number reports LCVIII. *Taxon* 29: 540.

Löve Á., D. Löve 1956. Cytotaxonomical conspectus of the icelandic flora. *Acta Horti Gotob.* 20: 69–291.

Májovský J. et al. 1970. Index of chromosome numbers of the slovakian flora (part 1). *Acta Fac. Rerum Nat. Univ. Comenianae Bot.* 16: 1–26.

— 1974. id. (part 4). *Acta Fac. Rerum Nat. Univ. Comeinanae Bot.* 23: 1–23.

Marsh C.A. 1955. Glucuronide metabolism in plants 2. The isolation of flavone glucosiduronic acids from plants. *Biochem. Jour.* 59: 58–62.

Molisch H., G. Goldschmidt 1901. Über das Scutellarin, einen neuen Körper bei *Scutellaria* und anderen Labiaten. *Monatsh. Chemie* 22: 679–699.

Moore R.J. (ed.) 1973. Index to plant chromosome numbers 1967–1971. *Regnum Veget.* 90: Oosthoek, Utrecht, 539 pp.

Morton J.K. 1973. A cytological study of the british Labiatae (excluding *Mentha*). *Watsonia* 9: 239–246.

Mulligan G.A. 1959. Chromosome numbers of canadian weeds 2. *Can. Jour. Bot.* 37: 81–92.

Müntzing A. 1927. Chromosome number, nuclear volume and pollen grain size in *Galeopsis*. *Hereditas* 10: 241–260.

— 1929. Cases of partial sterility in crosses within a linnean species. *Hereditas* 12: 297–319.

— 1930a. Outlines to a genetic monograph of the genus *Galeopsis* with special reference to the nature and inheritance of partial sterility. *Hereditas* 13: 185–341.

— 1930b. Über Chromosomenvermehrung in *Galeopsis*-Kreuzungen und ihre phylogenetische Bedeutung. *Hereditas* 14: 153–172.

— 1931. Chromosomenvermehrung in *Galeopsis*-Kreuzungen und ihre phylogenetische Bedeutung. *Zeitschr. indukt. Abstamm. Vererb. Lehre* 57: 360–362.

— 1932a. Disturbed segregation ratios in *Galeopsis* caused by intraspecific sterility. *Hereditas* 16: 73–104.

— 1932b. Cyto-genetic investigations on synthetic *Galeopsis tetrahit*. *Hereditas* 16: 105–154.

— 1938. Sterility and chromosome pairing in intraspecific *Galeopsis* hybrids. *Hereditas* 24: 117–188.

— 1941. New material and cross combinations in *Galeopsis* after colchicine-induced chromosome doubling. *Hereditas* 27: 193–201.

— 1943. Fertility improvement by recombination in autotetraploid *Galeopsis pubescens*. *Hereditas* 29: 201–204.

Östergren G., W. Heneen 1962. A squash technique for chromosome morphological studies. *Hereditas* 48: 332–341.

Plouvier V. 1963. Sur la recherche des hétérosides flavoniques dans quelques groupes botaniques. *C.R. Acad. Sci. Paris* 257: 4061–4063.

Porsch O. 1903. Die österreichischen *Galeopsis*arten der Untergattung *Tetrahit* Reichb. Versuch eines natürlichen Systems auf neuer Grundlage. *Abhandl. k.-k. zool-botan. Gesellsch. Wien* 2 (2): 1–126.

Rogenmoser E. 1975. Isolierung, Charakterisierung und Strukturaufklärung der Iridoidglucosiden von *Galeopsis pubescens* Besser. Diss. ETH Zürich 5574, 93pp.

Sieber V.K., B.G. Murray 1980. Spontaneous polyploids in marginal populations of *Alopecurus bulbosus* Gouan (Poaceae). *Bot. Jour. Linn. Soc.* 81: 295–300.

Sokolovskaya A.P. 1963. Georgr. rasprostr. poliploidn. vidov rasten. (issled. flory poluostr. Kamchatki). *Vestn. Leningr. Univ., Ser. Biol.* 3 (15): 38–52.

Stace C.A. 1980. Plant taxonomy and biosystematics. Edward London, 279 pp.

Sticher O. 1970a. Galiridosid, ein neues Iridoidglucosid aus *Galeopsis tetrahit* L. (Labiatae). *Tetrahedron Lett.* 36: 3197–3200.

- 1970b. Galiridosid, ein Iridoidglucosid aus *Galeopsis tetrahit* L. (Labiatae). *Helv. Chim. Acta* 53: 2010–2020.
- , E. Rogenmoser, A. Weisflog 1975. Neue Iridoidglucoside aus *Galeopsis tetrahit* L. und *Galeopsis pubescens* Bess. (Labiatae). *Tetrahedron Lett.* 5: 291–294.
- , A. Weisflog 1975. Nachweis und Isolierung der Iridoidglucoside aus *Galeopsis tetrahit* L. (Labiatae) sowie Strukturaufklärung von Glurosid. *Pharm. Acta Helv.* 50: 394–403.
- Strecker E. 1909. Das Vorkommen von Scutellarin bei den Labiaten und seine Beziehungen zum Lichte. *Sitz. Ber. kais. Akad. Wissensch. Wien, math. naturw. Kl., Abt. I* 118: 1379–1402.
- Taylor H. L., G. A. Mulligan 1968. Flora of the Queen Charlotte Islands, Part 2. Cytological aspects of the vascular plants. Queen's Printer, Ottawa, 148 pp. (ex R. J. Moore 1973).
- Townsend C. C. 1972. *Galeopsis*: T. G. Tutin et al. (ed.). *Flora Europaea* 3: 145–147. Cambridge Univ. Press. London and New York.
- Trotin F., M. Pinkas 1979. Sur les polyphénols du *Galeopsis ochroleuca* Lam. (Labiées). *Plantes Méd. Phytothérapie* 13: 94–98.
- Weisflog A. 1975. Isolierung, Charakterisierung und Strukturaufklärung der Iridoidglucoside von *Galeopsis tetrahit* L. Diss. ETH Zürich 5503, 83 pp.
- Wieffering J. H., L. H. Fikenscher 1974a. Aucubinartige Glucoside als systematische Merkmale bei Labiaten – 1. *Lamiastrum*. *Biochem. Syst. Ecol.* 2: 31–37.
- 1974b. id. 2. *Galeopsis*. *Biochem. Syst. Ecol.* 2: 39–46.
- Zeilinga A. E., G. H. Kroon 1965. A method for making root-tip squashes permanent without removal of the cover slip. *Euphytica* 14: 36–38.