**Zeitschrift:** Botanica Helvetica

Herausgeber: Schweizerische Botanische Gesellschaft

**Band:** 98 (1988)

Heft: 2

**Artikel:** Principal component analysis of ecological indicator values of swiss

alpine flora

Autor: Raamsdonk, L.W.D. van

**DOI:** https://doi.org/10.5169/seals-68582

## Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

# Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF: 28.10.2025** 

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# Principal component analysis of ecological indicator values of swiss alpine flora

# L. W. D. van Raamsdonk

Institute for Horticultural Plant Breeding, P.O. Box 16, Wageningen, The Netherlands.

Manuscript accepted July 19, 1988

#### Abstract

L. W. D. van Raamsdonk 1988. Principal component analysis of ecological indicator values of Swiss alpine flora. Bot. Helv. 98: 195–205.

Indicator values for eight ecological parameters of 303 Swiss alpine and subalpine species were used to evaluate the preferences of the species and of main vegetation units, and to study the relation between the parameters. The species were divided into 15 different groups representing formal vegetation units which were indicated as snow fields, rocks and screes, meadows, swamps, arid vegetations, scrubs and forests. Most of these categories splitted according to calcareous or aciduous preferences. After principal component analysis the eight parameters were divided into three groups encompassing humus content, soil structure, humidity and continentality (group 1), light, temperature and nutrients (groups 2), and acidity (group 3). The humidity and acidity of the soil appeared to be the most important ecological parameters. The ecological preferences of species and vegetation units are discussed. Special attention was paid to species with identical indicator values and to the range in ecological preference of the species composition of six particular plots.

Key words: ecology, vegetation analysis, indicator values, Swiss alpine flora.

# Introduction

Organisms can only survive when adapted to the conditions of their own habitat. Especially plants are strongly influenced by climatic and edaphic factors. A close relationship exists between soil and vegetation especially in alpine regions. The acidity, humidity, temperature and structure of the soil on a particular site determine largely the floristic composition of typical vegetations.

Ecological indicator values of the Swiss flora were gathered by Landolt (1977). Each species included was characterized by the response to ten ecological parameters, i.e. humidity, acidity, nutrients, humus, soil structure (dispersion and aeration), light, temperature, continentality, salt tolerance and life form. The last two parameters mentioned dealing with the alpine flora were omitted by Landolt (1984). The growth and vigour of plant species depends on the ability to compete and on the relation between ecological parameters, and can hardly be indicated by a few figures used like weighted averages (ter

Braak and Barendrecht 1986), without indication of variation and differentiation. However, numerical representations make easy evaluation and comparison of ecological preferences of species possible. An indication of ecological circumstances of a particular location can be obtained by using as many species as possible (Landolt 1977).

The classification of vegetation units is based on the occurrence of indicator and differential species and, on the highest levels of the system, on physiognomy. The associations and other categories described in this manner can be characterized by a special preference for only one or two ecological parameters and tolerance for large fluctuations of others. The classes of alpine scree, snow field and rock vegetations are subdivided in groups of associations preferring calcareous and aciduous soils, respectively. Associations growing under wet and moist conditions, and those which are found in arid situations are placed into several not related classes (Ellenberg 1978).

The indicator values compiled by Landolt (1977) were used to analyse ecological preferences of species and of particular locations.

#### Material and methods

At first a selection of mainly alpine and some subalpine species with the vegetation units in which they occur was taken from Rothmaler (1976). Species occurring in several different vegetation types like snowfields, rocks and meadows were gathered in two main groups of general calcicole or general calcifuge soils, respectively. From this first selection a list of 303 species of the Swiss alpine and subalpine flora was compilated with indicator values taken from Landolt (1977) and modified according to Landolt (1984). Special attention was paid to the indicator and differential species mentioned by Ellenberg (1978) and Runge (1986). The eight ecological parameters are:

humidity of the soil
acidity of the soil
contents of nutrients in the soil
content of humus in the soil
soil structure (particle size and aeration)
average light emittance
average temperature during vegetation period
continentality (fluctuations of air temperature and humidity)

The 303 species were divided in 15 different groups which are comparable to formally described vegetation units as indicated by Rothmaler (1976). These groups are:

- 1 general calcicole species
- 2 general calcifuge species
- 3 snowfieldvegetation, calcareous

4 snowfieldvegetation, aciduous

- 5 rock and scree vegetation, calcareous
- 6 rock and scree vegetation, aciduous
- 7 alpine meadows, calcareous
- 8 alpine meadows, aciduous
- 9 nutrient-rich meadows
- 10 wet meadows and swamps
- 11 arid vegetations
- 12 serpentine rock vegetations

Arabidetalia caeruleae Salicetalia herbaceae

Potentilletalia caulescentis, Thlaspietalia rotundifolii

Androsacetalia vandelli, Androsacetalia alpinae

Seslerietalia variae

Caricetalia curvulae, Nardetalia

Arrhenatheretalia

Tofieldietalia, Caricetalia fuscae,

Montio-Cardaminetalia

Sedo-Scleranthetalia, Seslerietalia p.p.,

Festuca-Brometea p.p., a.o.

Drabetalia hoppeanae

13 subalpine forests (low light conditions)

Erico-Pinetea, Vaccinio-Piceetea

14 general meadow species without preference for groups 7 or 8

15 alpine and subalpine scrub vegetation

Adenostyletalia, Artemisietalia

The a-priori group division was used to prove the consistency of the present classification of vegetation units after an analysis of ecological indicator values.

The program package IRIS for multivariate analysis was used (van Raamsdonk 1986). Three steps were involved in the analysis. First, the Pearson correlation coefficients between the parameters were studied. Weighting factors were established for each parameter based on the correlation matrix and used to calculate the covariances. Secondly, principal component analysis was carried out according to the usual way based on the covariance matrix. The third step was the calculation of a weighted covariance matrix based on six parameters excluding humidity and acidity. Principal components were calculated in order to investigate the possibility of defining the two excluded parameters by means of the other six parameters used.

The species composition of six different localities were analysed by selecting the species in the principal component plots based on the covariance matrix of eight parameters. The localities were:

il

## Results

A number of species showed the same indicator values. These species are listed in tab. 1. Nine out of 48 species preferring meadows on calcareous soils (group 7) belong to the Papilionaceae against one out of 53 in meadows on silicate soils. Similar observations can be made concerning nutrient-rich meadows (group 9: 12 monocotyledons out of 21 species) and wet meadows and pastures (group 10: 11 Cyperaceae out of 26 species). In four genera two or three species showed identical indicator values. These are listed in tab. 1. Three groups of ecological parameters could be recognised based on Pearson correlation coefficients. Values higher than  $\pm 0.4$  were given in tab. 2. Humidity, amount of humus, soil structure and continentality appeared to be closely related. The highest correlation was found between humus and soil structure, and between humidity and continentality. The amount of nutrients, light and temperature formed a second group. Correlation coefficients between acidity and the other parameters were below  $\pm 0.3$ , thus forming a parameter of its own.

The weighting factors of soil structure, light, temperature and continentality were set to 0.5 prior to calculating the covariance matrix, leaving the other weighting factors to 1.0. This setting was made because of the calculated correlation coefficients and of relative importance of ecological circumstances in the environment.

The first principal component was mainly determined by acidity (-0.738), humus (0.687) and humidity (0.516) and accounted for 39.89% of total variation. Acidity and

Tab. 1. List of taxa with identical indicatorvalues. The scores on the first and second principal components of the points indicated in figure 2 are added.

Name	Group	In	dica	ator	val	lues		Compone	Component scores on			
	(IRIS code)									1st pc	2nd pc	
Hutchinsia alpina Soldanella minima	3	4	4	2	3	2	5	1	2	-0.021	0.051	
Arenaria biflora Primula integrifolia	4	4	2	2	4	4	4	1	2	0.096	0.002	
Cerastium latifolium Papaver rhaeticum Sedum atratum Saxifrage biflora Saxifraga moschata	5	3	4	2	2	2	5	1	3	-0.072	0.005	
Galium helveticum Chrysanthemum halleri Achillea atrata	5	3	5	2	2	2	5	1	3	-0.100	0.046	
Poa laxa Ranunculus glacialis	6	4	2	2	2	2	4	1	3	0.007	-0.044	
Cerastium uniflorum Eritrichium nanum Achillea nana	6	3	2	2	2	2	5	1	3	-0.016	-0.079	
Anemone narcissiflora Trifolium badium Trifolium thalii Hedysarum hedysaroides Astragalus frigidus	7	3	4	3	3	4	4	2	3	-0.004	0.066	
Polygala alpestris Scabiosa lucida	7	2	4	2	3	3	4	2	3	-0.046	0.007	
Bupleurum stellatum Armeria alpina Pedicularis kerneri	8	2	2	2	3	3	4	1	4	0.004	-0.090	
Carex curvula Gentiana alpina Phyteuma globularifolium	8	2	2	2	3	3	5	1	4	0.000	-0.097	
Senecio carniolicus Senecio incanus	8	2	2	2	4	3	5	1	4	0.027	-0.087	
Pulsatilla alba Antennaria dioica	8	2	2	2	4	4	4	2	3	0.052	-0.058	
Nardus stricta Geum montanum Potentilla aurea Gentiana kochiana	8	3	2	2	3	4	4	2	3	0.045	-0.040	
Androsace carnea Androsace obtusifolia Androsace brigantiaca Euphrasia minima	8	3	2	2	4	4	4	1	3	0.071	-0.034	

Tab. 1. (continued)

Name	Group	In	dica	ator	va	lues		Component scores on			
	(IRIS code)									1st pc	2nd pc
Coeloglossum viride Astrantia minor Plantago alpina	8	3	2	2	4	4	4	2	3	0.072	-0.030
Campanula barbata Arnica montana Hypochaeris uniflora Leontodon helveticus Hieracium aurantiacum											
Crocus albiflorus Crepis aurea	9	3	3	4	4	4	4	2	3		
Phleum alpinum Poa alpina	9	3	3	4	3	4	4	2	3		
Veratrum album Trollius europaeus	9	4	3	3	4	5	4	2	3		
Trichophorum alpinum Eriophorum angustifolium	10	5	2	2	5	5	5	2	3		
Trichophorum cespitosum Carex nigra	10	4	2	2	5	5	4	2	3		
Sedum annuum Sempervivum montanum	11	2	2	1	3	3	5	2	3		
Nigritella nigra Thesium alpinum	14	3	3	2	4	4	4	2	3		
Bulbocodium vernum Lilium bulbiferum	14	2	3	2	3	3	4	3	4		

Tab. 2. Pearson correlation coefficients (R) between seven ecological parameters. All values higher than  $\pm 0.4$  are shown, values below that limit only between parameters of the same group.

Humus content Soil structure Humidity Continentality	1.0 0.674 0.331 -0.219	1.0 0.410 -0.227	1.0 -0.625	1.0			
Light Temperature Nutrients		R < = plu	ısminus 0.3	38	$   \begin{array}{r}     1.0 \\     -0.456 \\     -0.443   \end{array} $	1.0 0.259	1.0
	humus	soil gr	humid.	cont.	light	temp. group 2	nut.




Fig. 1. Principal component diagram based on a weighted covariance matrix of ecological indicatorvalues. Species preferring general calcicole soils (upper open triangles), general calcifuge soils (lower open triangles), general meadows (rhombs), serpentine (upper solid triangles), wet places (plus-signs) and arid places (x) are indicated.




Fig. 2. Principal component diagram based on a weighted covariance matrix of ecological indicator values. Species preferring snowfields (calcareous: upper triangles, aciduous: plus-signs), meadows (calcareous: squares, acidous: rhombs), screes and rocks (calcareous: lower triangles, acidous: x-es). Only one general sign each and the species with identical indicatorvalues listed in table 1 are plotted.

Individual plots of other species are deleted.




Fig. 3. Principal component diagram based on a weighted covariance matrix of ecological indicator values. Species preferring nutrient-rich meadows (plus-signs), forests (rhombs) and scrubs (lower triangles) are plotted.




Fig. 4. Principal component diagram based on a weighted covariance matrix of six selected ecological indicatorvalues. Species originating from wet places (crosses), general meadows (rhombs), general calcicole species (upper triangles), general calcifuge species (lower triangles) and arid places (times-sign) are plotted.




Fig. 5. Principal component diagram based on a weighted covariance matrix of ecological indicator values. Species originating from meadows on Strelapass (Graubünden; upper triangles, scrub inhabitants: plus-signs) and Kühboden (Wallis; lower triangles, scrub inhabitants: x-es, Sempervivum species: solid lower triangles) are plotted.




Fig. 6. Principal component diagram based on a weighted covariance matrix of ecological indicator values. Species originating from silicate rock on Albulapass (Graubünden) are plotted. The preferences are: rocks (x-es), meadows (rhombs), snowfields (plus-signs), low light conditions (lower triangles) and wet places (upper triangles).

humidity showed also high factor loads on the second principal component (0.720 and 0.484 respectively with 26.03% of total variation). Soil structure showed a moderately high factor load (0.410) on the first component unless the weighting factor of 0.5.

The component scores of each species were plotted against the first and second principal component. Some general groups were shown in fig. 1. One diagonal is determined by humidity with moist preferring species in the upper right part and species of arid places in the lower left part. Silicate preference is located in the lower right part against limestone preference in the upper left part of the plot. Species predominantly found on serpentine act like inhabitants of calcareous soils. Scree and rock vegetations, snow field vegetations and meadows are shown in fig. 2. Snow fields appeared to be moist on the average, while rocky places have a low humid soil compared to meadows. Fig. 3 shows the regions covered by representatives of nutrient-rich meadows, forests and scrubs. These prefer more or less moist habitats such as deep soils with good water capacity.

An attempt was made to compose humidity and acidity axes based on the other ecological parameters by setting the weighting factors of the two parameters to zero. The first principal component (50.15% of total variation) was mainly based on soil structure (-1.029) and humus content (-0.711) which determined a cline from wet to dry in fig. 4. Acidity was not expressed. Representatives of meadows on limestone and silicate shared the same region. Limestone and silicate preferring species were even located on the same place (circle in fig. 4). Inhabitants of meadows without special preference took an intermediate position in fig. 4.

The position of the inhabitants of two meadows is presented in fig. 5. The variability in ecological amplitude found in the dolomite meadow (Strelapass) is greater than represented by the inhabitants of the silicate meadow (Kühboden). All four species showing ecological parameters identical to Nardus stricta as listed in tab. 1 were found in the silicate meadow together with for instance Arnica montana and Campanula barbata. Sempervivum montanum and S. tectorum were accompanying species on dry patches. Several species were found on both locations which are also representatives of forest vegetations. Species with very different ecological preferences were found on a location of about 30 square metres on a steep rock with some soil patches at the Albulapass (fig. 6). Representatives of rock vegetations, silicate meadows, snow fields, forests and marshes were abundant, representing a whole range from dry to wet conditions. Species adapted to low light conditions survive because of the northern exposition and shady places. Two different species-poor locations on scree together with a marsh were analysed, demonstrating contrasts between dry and wet conditions, and between calcareous and silicate rock. The inhabitants of the marsh show a large ecological amplitude.

## Discussion

The occurrence of related species in exactly the same habitat needs a close examination in relation to the possibility of sympatric evolution in plants (Grant 1981). The species pairs Saxifraga moschata and S. biflora, and Trifolium thalii and T. badium possess identical ecological preferences (tab. 1). However, the two species in both pairs belong to different species groups within the genus which indicates a parallel evolution (Hess et al. 1970). Senecio carniolicus and the related S. incanus are two geographically distinct species which are very closely related (Merxmüller 1952; Landolt 1984). These species can be regarded as being involved in a process of geographic evolution. The

allopatric species Androsace carnea, A. obtusifolia and A. brigantiaca belong to the same species group. The first two species are diploid inhabitants of silicate meadows and differ in some minor morphological characters. The morphologically intermediate A. brigantiaca occurs in a limited region in the southwestern Alps, is an aneuploid on the tetraploid level and shows a greater ecological amplitude compared to the other two species (Hess et al. 1970). A. brigantiaca can be considered to be an allotetraploid derivative of A. carnea and A. obtusifolia.

Several ecological circumstances vary greatly in relation with elevation (Ellenberg 1978): on higher levels in mountainous areas more light, especially ultraviolet, lower temperatures, greater temperature fluctuation (continentality) and a lower rate of production of soil (erosion) can be found compared to lower altitudes. When considered within the frame of the alpine flora these ecological parameters show less variation compared to factors like humidity, acidity and nutrient content of the soil (Ellenberg 1978). This relative importance is in agreement with the chosen weighting factors and the proposed groups of parameters.

The situation that acid soils are usually low in nutrient amount is not expressed in a high correlation. The nutrient content of the soil is associated with the occurrence of flowing ground water which offers an additional supply of nutrients. The floating water situation is not included in the numerical value of soil humidity, but in an additional letter (Landolt 1977). In an analysis of South African heath vegetations Low (1987) found no strong relation betwen high calcium content (acidity) and other high nutrient levels. Species with a low nutrient tolerance preferred acid soils with a low level of calcium. High nutrient preference was not related to calcareous soils (Low 1987). The range of acid preference in the inhabitants of wet places is far greater compared to drought-preferring species, which grow exclusively on silicate soils (fig. 1). This situation is due to the fact that large silicate mountain ranges are located predominantly in the dry inner Alps (Hess et al. 1967). Richard (1985) analysed 46 plots by using factor analysis based on the species composition. The first and second factor appeared to be determined by mainly the acidity and the mobility of the soil. A fairly large range of altitude was involved (Richard 1985).

A reverse relationship is indicated by Landolt (1977) between humidity and continentality which is illustrated by a strong negative correlation coefficient (tab. 2). Similarly a relation was found between the seed set and some climatic parameters in the *Ornithogalum umbellatum/angustifolium* group. Species originating from the coastal parts of western Europe (low continentality) showed a higher seed set in wetter seasons than in dryer years, while continental species showed the reverse situation (van Raamsdonk 1985).

A high humus content is strongly associated with low permeability, low ventilation and varying water content. The highest correlation coefficient was found in this group of parameters. For instance peat is mainly inhabited by indicators for oxygen deficiency (Landolt 1977).

Several microspecies were distinguished in the *Ranunculus montanus* group and in the *Cardamine pratensis* group which differ within each group with respect to humidity and acidity preference of the soil (Landolt 1971). These examples illustrate the important impact of both these ecological parameters on the vegetation structure. The correctness of classification of vegetation units according to acidity and humidity of the soil (Ellenberg 1978; Runge 1986) could not be proved for both parameters, since it was impossible to generate an acidity axis in the principal component plots based on the remaining parameters contrary to humidity which appeared to be related to several other parame-

ters (fig. 4 and 5). Indicator values for moisture obtained from Ellenberg (1979) were used to calculate average moisture values per plot based on the species composition (ter Braak and Gremmen 1987). Response curves were calculated for each species based on the average moisture values of the plots where each species was found. A reasonable consistency was found between the indicator values of Ellenberg (1979) and the recalculated averages from the response curves per species (Ter Braak and Gremmen 1987).

Landolt (1977) used the indicator values of species listed from several plots to discuss the ecological circumstances on these particular plots. The usefulness of multivariate methods to summarize the circumstances on selected locations is illustrated by the examples discussed.

Many thanks are due to Prof. Dr. E. Landolt (Zürich) for critical reading the manuscript and for his valuable remarks.

#### References

Braak C. J. F. ter, and Barendrecht L. G. 1986. Weighted averaging of species indicator values: its efficiency in environmental calibration. Math. Biosci. 78: 57-72.

Braak C. J. F. ter, and Gremmen N. J. M. 1987. Ecological amplitudes of plant species and the internal consitency of Ellenberg's indicator values for moisture. Vegetatio 69: 79–87.

Ellenberg H. 1978. Vegetation Mitteleuropas in den Alpen. Ulmer, Stuttgart.

Ellenberg H. 1979. Zeigerwerte der Gefäßpflanzen Mitteleuropas. 2<sup>nd</sup> ed., Scripta Geobotanica 9, Göttingen.

Grant V. 1981. Plant speciation, 2<sup>nd</sup> ed. Columbia Univ. Press, New York.

Hess H. E., Landolt E. and Hirzel R. 1967. Flora der Schweiz, Band 1. Birkhäuser, Basel, Stuttgart.

Hess H. E., Landolt E. and Hirzel R. 1970. Flora der Schweiz, Band 2. Birkhäuser, Basel, Stuttgart.

Landolt E. 1971. Ökologische Differenzierungsmuster bei Artengruppen im Gebiet der Schweizerflora. Boissiera 19: 129–148.

Landolt E. 1977. Ökologische Zeigerwerte zur Schweizerflora. Veröff. Geobot. Inst. ETH, Stiftung Rübek 64, 208 S.

Landolt E. 1984. Unsere Alpenflora. Verlag Schweizer Alpenclub, 318 S.

Low A. B. 1987. Nutrient patterns in Southern Hemisphere heathlands with particular reference to South Africa. XIV. Internat. Bot. Congr., Berlin.

Merxmüller H. 1952. Untersuchungen zur Sippengliederung und Arealbildung in den Alpen. Ver. Schutz Alpenpfl. u. -tiere, München, 105 S.

Raamsdonk L. W. D. van 1985. Pollen fertility and seed formation in the *Ornithogalum umbellatum/* angustifolium complex. Pl. Syst. Evol. 149: 287–297.

Raamsdonk L. W. D. van 1986. IRIS: program package for information regrouping to identify structures. Wageningen.

Richard J.-L. 1985. Pelouses xerophiles alpines des environs de Zermatt (Valais, Suisse). Bot. Helv. 95: 193-211.

Rothmaler W. 1976. Exkursionsflora, Kritischer Band. Volk und Wissen Volkseigener Verlag, Berlin.

Runge F. 1986. Die Pflanzengesellschaften Mitteleuropas. Aschendorff, Münster.