

A new subspecies of *Saccoloma elegans* : studies in the Flora of the Guianas, no 39

Autor(en): **Cremers, G. / Kramer, K.U.**

Objektyp: **Article**

Zeitschrift: **Botanica Helvetica**

Band (Jahr): **99 (1989)**

Heft 1

PDF erstellt am: **20.06.2024**

Persistenter Link: <https://doi.org/10.5169/seals-69128>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

A new subspecies of *Saccoloma elegans*. Studies in the Flora of the Guianas, no 39

G. Cremers¹ and K. U. Kramer²

¹ORSTOM, B.P. 165, F-97323 Cayenne-Cedex, France

²Institut für Systematische Botanik, Zollikerstr. 107, CH-8008 Zürich, Switzerland

Manuscript accepted January 31, 1989

Abstract

Cremers G., Kramer K. U. (1989). A new subspecies of *Saccoloma elegans*. Bot. Helv. 99: 45–48.

The fern *Saccoloma elegans* Kaulf. (Dennstaedtiaceae), widespread in tropical America, consists of two subspecies, ssp. *elegans* of S.E. Brazil and ssp. *chartaceum*, newly described, from Central America, the Greater Antilles, and N.W. South America.

The genus *Saccoloma* Kaulf. (Dennstaedtiaceae), including *Ithycaulon* Copel. and *Orthiopteris* Copel. (following, e.g., R. Tryon 1962, 1964, and Kramer 1978) comprises about ten species. It is pantropic in distribution but absent from continental Africa. When attempting to revise the genus, G. B. Nair, then working with the junior author, noted that the neotropical *S. elegans* Kaulf., as currently understood, consisted of two morphologically and geographically distinct entities. As his work was neither completed nor published, the newly discovered taxon remained undescribed. Within the framework of the recently begun "Flora of the Guianas", edited by Görts-van Rijn, Utrecht, it has become necessary to describe the new taxon formally. At the time, Nair regarded it as specifically distinct, but we prefer to treat it at subspecific rank, the two entities constituting what was *S. elegans* s. lat. being morphologically very close and geographically vicarious.

Saccoloma elegans Kaulf., Berl. Jahrb. Pharm. 21: 51, 1820.

Davallia saccoloma Sprengel, Syst. Veget. 4: 119, 1827 (non *D. elegans* Swartz 1801, nec Hedwig 1803), homotypic synonym.

Subsp. *elegans*

Type: not cited with the protologue; but in the same author's Enumeratio Filicum (1824, p. 224, pl. 1) Sello s.n. from Brazil is cited as the only collection, which must therefore be regarded as the type (Isotypes B! P!).

Rachis stramineous; most often 15–20(–29) pinnae to a side, these less than 3 cm wide, herbaceous; veins often forked, ca. (8–)11–17(–23) forked veins per 3 cm length of pinna.

Distribution: E-SE Brazil.




Fig. 1. – *Saccoloma elegans* Kaulf.: 1–2 subsp. *elegans*, 1, (right half of figure) lamina ($\times 0.4$); 2, part of a pinna with venation ($\times 1$). – 3–8 subsp. *chartaceum* Nair ex Cremers and Kramer: 3, rhizome with petiole bases ($\times 0.5$); 4, (left half of figure) lamina ($\times 0.4$); 5, pinna base ($\times 1.5$); 6, apex of a pinna ($\times 1$); 7, margin of a pinna with sori ($\times 3.5$); 8, part of a pinna with venation ($\times 1.3$) (drawn from Smith 2855, NY).

Other specimens examined: BRAZIL: Bahia: Blanchet 3976 (P); Rio Belmonte, Anon. 38 (BR); Ilheus, Luschnath 3936, comm. Blanchet 1857 (BR). – Minas Gerais: Pomba, Oliveira L. 1982 (P); road to São Miguel, Distr. Ilhem, Mexia 4964 (GH, NY, P, UC, US), 4639 (B, NY, P, U, US, Z); Gongo Soco, Bunbury s.n. 1863 (BR). – Rio de Janeiro: Faz. de Santa Ana, Glaziou 2376 (BR, P); Serra dos Orgãos, Gardner 159 (BM, P); *ibid.*, Pascoe s.n. (P); Glaziou 1746 (BR). – Santa Catarina: Arambuja-Brusque, Reitz 3454 (U). – Without loc., Miers s.n. 1878 (P); Riedel s.n. (P); Warre s.n. (P); Weddell 1006 (P); Langsdorff s.n. 1821 (BR).

Subsp. *chartaceum* G. B. Nair ex Cremers et Kramer, subsp. nova

Type: Guyana, Kanuku Mts, 3°08' N, 59°23' W, 200–400 m, Jansen-Jacobs et al. 338 (holotype U; isotypes B, BBS, BRG, CAY, NY, P, TRIN, US, Z).

Rhachis plus minusve profunde purpureo-castanea; pinnae pro latere 8–15(–17), ultra 3 cm latae; textura chartacea; venae saepius simplices, solum 1 ad 6 furcatae pro 3 cm longitudinis pinnae.

This subspecies diverges from subsp. *elegans* by the rachis being more or less deeply purplish-brown tinged; by the presence of only 8–15(–17) pairs of pinnae which are less than 3 cm wide; by the more or less distinctly chartaceous texture, and by the veins being mostly simple, (0–)1–6(–15) forked ones occurring over a length of pinna of 3 cm. We found no appreciable differences in the spores of the two subspecies; in both they are about 40 µm in diameter.

Distribution: Guatemala, Belize, Costa Rica, Panama, Cuba, Jamaica, Hispaniola, northern and western South America.

Paratypes: BELIZE: Big Creek, Schipp 90 (Z); – GUATEMALA: Alta Verapaz: von Türckheim 66 (BR, P); Quiche: Skutch 1807 (P). – COSTA RICA: Puntarenas: SE of Rincón de Osa, Evans 2812 (U). – PANAMA: Canal Zone: Maxon 4890 (P); Chagres, Isthmus of Panama, Fendler 411 (K); Panama: Juan Díaz, Killip 2331 (P); Finca Neptuna, NE of Lago Cerro Azul, road to Cerro Jefe, Nee 11531 (U); Darién: Cerro Sapo, Hammel, Franco, de Gaiza and Trainer 14879 (Z); without loc., Hooker 1870 (P). – CUBA: Without loc., Wright 1806 (P). – JAMAICA: Without loc., Ebbeke s.n. (P). – HISPANIOLA. REP. DOMINICANA: Barahona: Fuertes 995 (P, Z); Samaná Penins., Pitón de Azucar, Abbott 348 (P). – VENEZUELA: Mérida: Tovar, Fendler 376 (K); Amazonas: Ugueto, Croizat 798 (U). – GUYANA: Shodikar Cr., Essequibo R., Smith 2855 (GH, NY, US), Humirida Mts., Appun 1267 (GH); Sheenabowa, Potaro R., Jenman 1378 (BRG); Black Cr., Timoti R., Beckett s.n. 1906 (K); Potaro R., 1895, Jenman s.n. (BRG); Pakaraima Mts., Kopinang falls. Maguire et al. 46064A (NY, U, US); without loc., Appun s.n. (as Glaziou 12378) (P). – FRENCH GUIANA: Cayenne, Cascades R., Leprieur 260 (F, GH, P, US); Cayenne, Leprieur s.n. 1844 (B, P); Leprieur s.n. 1849 (P); without loc., Leprieur 267 (B); Desvaux s.n. (P); Richard s.n. (P). – COLOMBIA: Cundinamarca: Andes de Bogotá, 1857, Triana s.n. (P); Meta: Llano de San Martín, 1851/57, Triana s.n. (P); Triana 178 (K); Sierra de la Macarena, Caño Entada, Philipson and Idrobo 1750 (BM). – ECUADOR: Pichincha: Tocachi R., Sodiro s.n. (P); Verleysen 201 (P); Esmeraldas: Parr. de Concepcion, Playa Rica, Mexia 8429 (K, U). – PERU: Dept. unknown: Las Ganas, “Cordilleras peruv.”, Lechler s.n. 1854 (P). – BOLIVIA: La Paz: Larecaja, Copacabana, Krukoff 11134 (K, U); Mapiri: San Carlos, Buchtien 49 (Z), 1073 (P); Mapiri, San Antonio, Buchtien 1069 (P). – BRAZIL: Roraima: Botamatadi, Maita, Prance et al. 13611 (K, NY, U, US); Uaica airstrip, Rio Uracicoeira, Prance et al. 10821 (INPA, K, MG, U).




Fig. 2. Distribution of *Saccoloma elegans* Kaulf.: A, subsp. *elegans*; B, subsp. *chartaceum* Nair ex Cremers and Kramer.

Amazonas: São Gabriel da Cachoeira, Rio Negro, Spruce 2402 (P); Rio Purus, Rio Ituxí, Rio Curuquete, Prance et al. 14649 (INPA, K, MO, NY, U, US). Pará: Hwy. BR 22, Km 55, Prance and Silva 58897 (U); Hwy. BR 22, Km 96, Cachoeira, Prance and Pennington 1784 (U); Tanaii, Rio Acará near Belém, Spruce s.n. 1849 (P).

Literature

- Kramer K. U. 1978. The pteridophytes of Suriname. *Uitg. Natuurwet. Studiekr. Surin. Ned. Ant.* 93: 1–198.
- Stolze R. G. 1981. Ferns and fern allies of Guatemala. Part. III: Polypodiaceae. *Fieldiana Bot., New Series*, 6: 1–522.
- Tryon R. M. 1962. Taxonomic fern notes III. *Contr. Gray Herb.* 191: 100–106.
- Tryon R. M. 1964. The ferns of Peru. Polypodiaceae (Dennstaedtieae to Oleandreae). *Contr. Gray Herb.* 194:1–253.