

Zeitschrift:	Botanica Helvetica
Herausgeber:	Schweizerische Botanische Gesellschaft
Band:	100 (1990)
Heft:	1
Artikel:	Durio bukitrayaensis Kosterm. (Bombacaceae), a new species from Borneo
Autor:	Kostermans, A.J.G.H.
DOI:	https://doi.org/10.5169/seals-69709

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Durio bukitrayaensis Kosterm. (Bombacaceae), a new species from Borneo

A. J. G. H. Kostermans

Herbarium Bogoriense, Bogor, Indonesia

Manuscript accepted January 8, 1990

Abstract

Kostermans A. J. G. H. 1990. *Durio bukitrayaensis* Kosterm. (Bombacaceae), a new species from Borneo. Bot. Helv. 100: 29–31.

Durio bukitrayaensis is described as a new species from Borneo. It is related to *D. dulcis*, *D. grandiflorus*, and *D. testudinarum*.

Durio bukitrayaensis Kosterm., spec. nov.

Arbor mediocris, foliis alternantibus chartaceis oblongis (pro genere longis), longe cuspidatis basi rotundatis, supra subnitidis, sub lente minutissime reticulatis glabris, nervo mediano conspicue impresso, nervis filiformibus impressis, subtus nitidis adpresso lepidotis, squamulis non fimbriatis, nervo mediano prominente, nervis subpatentibus, epicalycis partibus ovato-acutis lepidotis, calycis partibus lepidotis squamulis majusculis, fructibus ellipsoideis, parvis, flavoviridibus spinis subcylindricis parvis acutis squamulis minutis fimbriatis obtectis.

Typus: Moga 4128 (BO).

Tree, 15 m tall, dbh. 15 cm. Bark dark brown, scurfy. Branchlets tightly crusty-scaly. Leaves scattered, chartaceous, oblong, 6 × 15 – 7 × 22 cm – 10 × 32 cm, acumen very slender, sharp, up to 4 cm long, base rounded; above green, under the lens minutely reticulate, nerves scabrous, no scales; below silvery brown or brownish-glaucous (fresh), glossy, covered with very tightly packed, flat, non-fimbriate scales of 0.5–0.75 mm diam., so tightly packed that their shape is hardly discernible, midrib strongly prominent, nerves rather patent, slender, prominent, c. 17 pairs with in between shorter ones. Petiole rather slender, densely lepidote, 10–15 mm long.

Flowers on the very base of the trunk, cauliflorous or on the thicker branches. Buds shortly ovoid, or conical. Epicalyx of broadly ovoid, acute, 2.5 cm long lobes, densely and tightly covered by non-fimbriate scales. Calyx lepidote, scales 1–1.2 mm diam.

Fruit ellipsoid, yellowish green, c. 7 cm long, 6 cm diam., rounded at both ends; spines rather slender, 1.5 cm long, densely, minutely fimbriate-scaly, at base c. 1 mm diam., stiff, sharp.

Distribution/Ecology. Lowland, 350 m, in wet, evergreen, tropical rainforest in the Bukit Raya Mts., Central Kalimantan.




Fig. 1. *Durio bukitrayaensis* Kosterm.

Note: The non-fimbriate, very tightly packed scales of the under surface of the leaves resemble those of *D. dulcis* and *D. grandiflorus* but the scales are much larger. Moreover, the leaves of *D. dulcis* and *D. grandiflorus* are smaller and differently shaped, the patent nerves are also different. Like in *D. testudinarum*, the flowers may occur on the very base of the bole, but the leaves are different, characteristic is the long draw-out slender acumen, and the scale are much smaller than those of *D. testudinarum*. The fruits are small, the spines look like those of *D. grandiflorus*.

Central Kalimantan, Bukit Raya in Upper Katingan (Mendawi) R. area, Upper Samba R., 60 km NNW of Tumbang Samba, 112°5' E, 0°60' S, alt. 350 m, Dec., fr. Mogea 4128 (BO); Upper Katingan R., NNW of Tumbau Samba, 112°30' E, 1°15' S, Dec., buds, Mogea 4417 (BO).