

**Zeitschrift:** Botanica Helvetica  
**Herausgeber:** Schweizerische Botanische Gesellschaft  
**Band:** 103 (1993)  
**Heft:** 1

**Artikel:** Cytotaxonomic studies in the Iberian taxa of the genus Lactuca (Compositae)  
**Autor:** Mejías, José A.  
**DOI:** <https://doi.org/10.5169/seals-71332>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 03.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Cytotaxonomic studies in the Iberian taxa of the genus *Lactuca* (Compositae)

José A. Mejías

Departamento de Biología Vegetal y Ecología, Facultad de Biología, Universidad de Sevilla, Apartado 1095, 41080 Sevilla, Spain.

Manuscript accepted January 19, 1993

### Abstract

Mejías, J. A. 1992. Cytotaxonomic studies in the Iberian taxa of the genus *Lactuca* (Compositae). Bot. Helv. 103: 113–130.

The 10 taxa representative of the genus *Lactuca* in the Iberian Peninsula have been studied, with the chromosome number for *L. viminea* subsp. *ramosissima* reported for the first time. All the taxa have been found to be diploid: nine of them with chromosome number  $2n = 18$ , and one with  $2n = 16$ . Some differences in chromosome morphology have been detected among diverse populations of some taxa; but in general, the separation of groups according to the karyological characters correlates with the present taxonomic classification of the group. The taxonomic status of *L. livida* is discussed.

*Key words:* *Lactuca*, chromosome morphology, Iberian Peninsula

### Introduction

The genus *Lactuca* is distributed world wide and can be considered the most evolved and successful from an evolutionary line comprising *Prenanthes*, *Cicerbita*, *Mycelis*, *Lactuca*, *Faberia* and *Cephalorhynchus* (Stebbins 1953). The genus *Cicerbita* is clearly implicated in the origin of *Lactuca*, and in turn may have arisen from *Prenanthes*, probably the most primitive genus of the group. This evolutionary process has involved the conversion of the columnar and beakless cypsela, characteristic of *Prenanthes*, into the compressed fruits of *Lactuca* which bear a beak, very short in their most primitive representatives.

The Mediterranean Region is an important centre of diversification of the genus *Lactuca*, where about 20 taxa can be found (Feráková 1977) from which a total of 10 taxa represent the genus in the Iberian Peninsula: *L. viminea* (L.) J. and C. Presl subsp. *viminea*, *L. viminea* subsp. *chondrilliflora* (Boreau) Bonnier, *L. viminea* subsp. *ramosissima* (All.) Bonnier, *L. serriola* L., *L. saligna* L., *L. virosa* L., *L. livida* Boissier and Reuter in Boissier, *L. perennis* L. subsp. *perennis*, *L. perennis* subsp. *granatensis* Fernández Casas and Charpin, and *L. tenerrima* Pourret; some of these (*L. livida*, *L. perennis* subsp. *granatensis* and *L. tenerrima*) are considered endemic in the Iberian Peninsula or the West

Mediterranean Basin. *L. sativa* L. has been excluded from this study because of its cultivation in the area.

The karyological studies in *Lactuca* are diverse because of the wide distribution of the genus and its economic interest; but they consist mainly of chromosome counts from some taxa without any particular interest. The most extensive studies are quite old (Babcock et al. 1937 sec. Feráková 1977, Thompson et al. 1941); two general publications on the tribe *Lactuceae* by Stebbins et al. (1953) and Tomb et al. (1978) also ought to be cited, because of some considerations on the taxonomic and chromosomic evolution of the tribe.

The general size and characteristics of the chromosomes in *Lactuca* enable the study of their morphology and the idiogrammatic formulas of the different taxa. Indications have sometimes been given (Stebbins et al. 1953, Feráková 1977) and, there is a very detailed study on *L. serriola* and *L. sativa* (Haque and Godward 1985a).

This paper reports cytological observations on the representatives of the genus in the Iberian Peninsula. The present investigations were carried out in conjunction with a study of the reproductive biology of *Lactuca* and some related genera with the aim of gaining a deeper biosystematic knowledge of the group.

## Material and methods

Mitotic and meiotic observations were made of material from 50 populations. Mitotic studies were done with root tips of plants obtained from seeds collected in the field, or with seedlings obtained by germinating either self-fecundating cypsela or wild seeds. The root tips were treated with 0.002 M 8-hydroxyquinoline for about 3.5 h at room temperature (Tjio and Levan 1950), fixed in Farmer's fluid – ethanol-acetic acid (3:1) (Löve and Löve 1975a) – and stored at 4°C in 70% ethanol. At first, material was stained in cold alcoholic-hydrochloric carmine solution; but staining with alcoholic orceine gave better results, so it was used afterwards. For meiotic studies, flower buds were fixed in the field in Carnoy's fluid: ethanol-chloroform-acetic acid (6:3:1) (Löve and Löve 1975a). They were kept in this for at least 24 h and preserved in 70% ethanol. In this case, staining was always in alcoholic hydrochloric acid-carmine solution.

Whenever possible, the idiogrammatic formula for the population has been indicated. The terminology of Levan et al. (1965) modified by Küpfer (1974) has been used for describing the morphology of chromosomes and the pairs established. These authors classified the chromosomes in six classes, depending on the rate  $r = \text{long arm length}/\text{short arm length}$ : metacentric (M) with the centromere in the median point ( $r = 1$ ); metacentric (m) with the centromere in the median region ( $1 < r < 1.7$ ); submetacentric (sm) with the centromere in the submedian region ( $1.7 \leq r < 3.0$ ); subtelocentric (st) with the centromere in the subterminal region ( $3.0 \leq r < 7.0$ ); telocentric (t) with the centromere in the terminal region ( $7.0 \leq r < 39$ ), and telocentric (T) with the centromere in the terminal point ( $39 \leq r \leq \infty$ ). The terminology of Stebbins (1938, 1971) has been used for apparent size of chromosomes and karyotype asymmetry. Coefficients  $A_1$  and  $A_2$  proposed by Romero (1986) have also been used for a more accurate estimation of karyotype asymmetry. If two chromosomes, assumed homologues, had to be included in different classes, an intermediate class for the pair was considered (e.g., m-sm for pairs comprising a metacentric chromosome and a submetacentric chromosome).

An average of three plants from each population was studied for meiosis and four for mitosis. Voucher specimens are conserved in the Herbarium of the Department of Botany, Faculty of Biology (see Appendix).

## Results

### *Chromosome numbers*

The chromosome numbers of the populations studied are given in Table 1. It can be seen that among the Iberian representatives the somatic number  $2n = 18$  is the most common, being reported in 9 of a total of 10 taxa: *Lactuca viminea* subsp. *viminea*, *L. viminea* subsp. *chondrilliflora*, *L. viminea* subsp. *ramosissima*, *L. serriola*, *L. saligna*, *L. virosa*, *L. livida*, *L. perennis* subsp. *perennis* and *L. perennis* subsp. *granatensis*. Only one other taxon, *L. tenerrima*, shows a different number,  $2n = 16$ . These chromosomes were always observed to go through regular meiosis, so the gametic numbers  $n = 9$  and  $n = 8$  have been reported, respectively.

### *Chromosome morphology*

In the present investigation, chromosome morphology was found to vary, in general between the classes metacentric (M and m) and telocentric (t), the class of submetacentric chromosomes (sm) being the most common overall and in every species, except for the representatives of the sect. *Lactuca* subsect. *Cyaniceae*: *L. perennis* and *L. tenerrima*. In these, metacentric chromosomes (M and m) prevailed, a class also frequent in the rest of the taxa. Some subtelocentric chromosomes (st) were also found, and a telocentric pair (t) was always observed in *L. virosa* and *L. livida*. This situation is reflected by the  $A_1$  coefficient that denotes *L. perennis* and *L. tenerrima* as the species with the morphologically most symmetrical chromosomes, whereas *L. serriola*, *L. saligna* and specially *L. virosa* and *L. livida* exhibit the most asymmetrical karyotypes. 1 or 2 satellite pairs of chromosomes are present in each karyotype.

The idiogrammatic formulas established from the study of the chromosome morphology are shown in Table 1. There is some degree of variation in arm ratios among the populations studied for a single taxa that is reflected in differences in the number of pairs included in each morphological class, and  $A_1$  coefficient values. Sometimes, these differences could have been increased by the antimitotic treatment, although special care was taken to homogenize the period of treatment. The most variable group in this sense is *L. viminea*, whose karyotype has been found to be composed mainly of metacentric and submetacentric chromosomes in variable ratios, regardless of the subspecies. Notable differences have been found in *L. perennis* and *L. tenerrima*, not attributable to antimitotic treatment. In the plants from Puerto de la Bonaigüa a regular karyotype comprising 9 homomorphic pairs of chromosomes has always been found, but among three plants studied from a population collected between Rosas and Cadaqués, two individuals with a heteromorphic pair of chromosomes were detected (Table 1, Figure 9), the remaining individual showing the same idiogrammatic formula indicated for Puerto de la Bonaigüa. It has not been possible to study the meiotic behaviour in the plants with the heteromorphic pair of chromosomes because of the lack of flower production in the experimental ground; in any case, they showed well-developed vegetative characters. Chromosome morphology in *L. tenerrima* was studied in material from two populations, those from Valle de Añisclo and Sierra de Cazorla (Table 1). Although the idiogrammatic formulas obtained are very similar, both karyotypes can be recognized by the position of the satellites, which are borne in the long arm of a pair of chromosomes in the plants from Sierra de Cazorla (Figure 12) and in a short arm in the plants from Valle de Añisclo (Figure 13), and in the remaining populations studied. The presence of conspicuous

Tab. 1. Chromosome number, idiogrammatic formula, karyotype asymmetry (A, A<sub>1</sub> and A<sub>2</sub> coefficients) and chromosome apparent length for the populations of *Lactuca* studied.

Taxon and locality	n	2n	Idiogrammatic formula	A	Coefficients		Chromosome length (µm)
					A <sub>1</sub>	A <sub>2</sub>	
<i>Sect. Phoenixopus</i> (Cass.) Bentham							
<i>L. viminea</i> (L.) J. and C. Presl							
subsp. <i>viminea</i>							
Jerte-Tornavacas	9II	18	4 m + 2 m-sm + 10 sm + 2 sm <sup>sat</sup>	2B	0.39	0.38	2.04–5.90
Escaro	9II						
Cazalla		18	8 m + 2 m-sm + 4 sm + 4 sm <sup>sat</sup>	2B	0.36	0.33	1.45–3.54
subsp. <i>chondrilliflora</i> (Boreau) Bonnier							
Rosas	9II	18	4 m + 10 sm + 4 sm <sup>sat</sup>	2B	0.41	0.32	1.07–3.19
La Molina-Alp	9II	18					
Viella-Bossost		18					
subsp. <i>ramosissima</i> (All.) Bonnier							
Tineo-Cangas		18					
Villaluenga		18	6 m + 6 sm + 4 sm <sup>sat</sup> + 2 st	2B	0.41	0.31	1.82–4.50
Puerto de la Ragua		18	8 m + 2 m-sm + 4 sm + 4 sm <sup>sat</sup>	2B	0.39	0.30	1.72–4.18
Sierra de Baza	9II						
Puerto de Alájar		18					
Viniegra		18	4 m + 2 m-sm + 8 sm + 4 sm <sup>sat</sup>	2B	0.43	0.32	2.79–6.22
Valdecastillo		18					
Algámitas		18	6 m + 2 m-sm + 6 sm + 4 sm <sup>sat</sup>	2B	0.38	0.34	1.72–4.72
Cazalla-Constantina		18					
<i>Sect. Lactuca</i> L.							
Subsect. <i>Lactuca</i>							
<i>L. serriola</i> L.							
Orrius-La Roca	9II	18	8 m + 6 sm + 2 sm <sup>sat</sup> + 2 st <sup>sat</sup>	2A	0.39	0.22	2.32–4.37
Lanavé		18	6 m + 8 sm + 2 sm <sup>sat</sup> + 2 st <sup>sat</sup>	3B	0.42	0.27	1.93–4.29
Villanueva-Beas		18					
Grajal		18	4 m + 2 m-sm + 10 sm + 2 st <sup>sat</sup>	3B	0.42	0.27	1.50–3.54
El Escorial		18					
Brenes	9II	18	6 m + 8 sm + 2 sm <sup>sat</sup> + 2 st <sup>sat</sup>	3B	0.44	0.25	2.14–4.39
Sevilla		18					
<i>L. saligna</i> L.							
Elciego	9II	18	6 m + 8 sm + 2 sm <sup>sat</sup> + 2 st <sup>sat</sup>	3A	0.38	0.24	2.04–4.07
Villanueva		18	6 m + 6 sm + 2 sm <sup>sat</sup> + 2 st + 2 st <sup>sat</sup>	3B	0.48	0.33	1.82–4.18
Grazalema		18	6 m + 8 sm + 2 sm <sup>sat</sup> + 2 st <sup>sat</sup>	3B	0.46	0.26	2.25–5.04
Otero de Sariegos		18	6 m + 8 sm + 2 sm <sup>sat</sup> + 2 st <sup>sat</sup>	3A	0.46	0.22	2.47–4.18
<i>L. virosa</i> L.							
Bienservida-							
Villaverde	9II						
La Espina	9II						
Puerto del Pico		18					
Orrius-La Roca		18	2 M + 8 sm + 2 sm <sup>sat</sup> + 4 st + 2 t	3A	0.55	0.23	2.14–4.07
Sierra de Baza		18	2 M + 8 sm + 4 st + 2 st <sup>sat</sup> + 2 t	3A	0.58	0.23	2.51–4.72
Los Marines-Aracena	9II						

Tab. 1. (continued)

Taxon and locality	n	2n	Idiogrammatic formula	A	Coefficients		Chromosome length ( $\mu\text{m}$ )
					A <sub>1</sub>	A <sub>2</sub>	
Fuenteheridos		18	2 M + 8 sm + 2 sm <sup>sat</sup> + 4 st + 2 t	3A-	0.56	0.23	3.11-6.33
Jaca		18		3B			
Puerto de Tíscar	9II						
Sierra de Cazorla		18					
Comillas		18	2 M + 8 sm + 2 sm <sup>sat</sup> + 4 st + 2 t	3B	0.54	0.27	2.57-5.57
S. Nicolás del Puerto		18	2 M + 8 sm + 2 sm <sup>sat</sup> + 4 st + 2 t	3A	0.58	0.23	2.89-5.68
<i>L. livida</i> Boiss. and Reuter							
S. Pablo de los Montes	9II	18	2 M + 8 sm + 2 sm <sup>sat</sup> + 4 st + 2 t	3A	0.59	0.22	2.36-4.32
subct. <i>Cyaniceae</i> DC.							
<i>L. perennis</i> L.							
subsp. <i>perennis</i>							
Rosas-Cadaqués	9II	18	2 M + 8 m + 2 m-sm + 2 sm + 2 sm <sup>sat</sup> + 2 sm-st <sup>sat</sup>	2A	0.34	0.17	2.89-4.93
Puerto de la Bonaigüa	9II	18	2 M + 8 m + 2 m-sm + 2 sm + 4 sm <sup>sat</sup>	2A	0.30	0.20	2.25-4.29
subsp. <i>granatensis</i>							
Charpin and Fdez. Casas							
Sierra de la Sagra		18	2 M + 10 m + 2 sm + 4 sm <sup>sat</sup>	2A	0.34	0.17	3.00-5.04
<i>L. tenerrima</i> Pourret							
Villajoyosa	8II						
Zahara-Puerto Palomas		16					
Bruguera		16					
Linares de la Sierra		16					
Valle de Añíscló		16	2 M + 4 m + 2 m <sup>sat</sup> + 2 m-sm + 6 sm	2A- 2B	0.33	0.21	1.18-2.57
Belerdas	8II						
Sierra de Cazorla		16	2 M + 4 m + 2 m <sup>sat</sup> + 8 sm	2A	0.34	0.16	1.40-2.47
Oseja de Sajambre		16					

secondary constrictions, especially in the larger pair of chromosomes, is notable in every case.

*L. livida* is an endemic Iberian species closely related to *L. virosa* for which there are no clear morphological boundaries (Feráková 1977, Linqvist 1960). Cytological observations have been made in material from the locality where *L. livida* was described by Boissier and Reuter: San Pablo de los Montes (see Appendix). In these plants the idiogrammatic formula was almost identical to those found in *L. virosa* (Table 1) which taxonomically reinforces the relationship of these taxa, and allows discussion on the validity of *L. livida* as species (see Discussion).




Fig. 1–7. 1. Karyogram of *L. viminea* subsp. *ramosissima* (SEV 126386),  $2n = 18$ . 2. Karyogram of *L. saligna* (SEV 126535),  $2n = 18$ . 3. Diakinesis of *L. viminea* subsp. *viminea* (SEV 126389),  $n = 9$ . 4. Somatic metaphase of *L. serriola* (SEV 126527),  $2n = 18$ . 5. Anaphase II (partial) of *L. serriola* (SEV 126524),  $n = 9$ . 6. Karyogram of *L. virosa* (SEV 124656),  $2n = 18$ . 7. Karyogram of *L. livida* (SEV 124612),  $2n = 18$ . Scale bar = 5  $\mu$ m.




Fig. 8–13. 8. Anaphase I of *L. perennis* subsp. *perennis* (SEV 126546),  $n = 9$ , 9. Karyogram of *L. perennis* subsp. *perennis* (SEV 126546),  $2n = 18$ . 10. Metaphase I of *L. tenerrima* (SEV 124653),  $n = 8$ . 11. Karyogram of *L. perennis* subsp. *granatensis* (SEV 126591),  $2n = 18$ . 12. Somatic metaphase of *L. tenerrima* (SEV 124646),  $2n = 16$ . 13. Karyogram of *L. tenerrima* (SEV 126571),  $2n = 16$ . Scale bar = 5  $\mu\text{m}$ .

Tab. 2. Previous chromosome counts for the Iberian taxa of *Lactuca*

Taxa	n	2n	Origin of material	Authors
<i>L. viminea</i> (L.) J. and C. Persl subsp. <i>viminea</i>		18	Slovenia	Feráková, 1970a: 14
		18	Portugal	Fernandes and Queirós, 1971: 83
		18	Portugal	Queirós, 1973: 311
		18	Bulgaria	Kuzmanov and Georgieva, 1980: 715
subsp. <i>chondrilliflora</i> (Boreau) Bonnier	9II		France	Whitaker and Jagger, 1939: 299
	9II		France	Thompson et al., 1941: 93
		18	Italy	Brullo and Pavone, 1978: 260
<i>L. serriola</i> L.	9II		Japan	Ishikawa, 1921: 156
	7II + 1IV		U.S.A.?	Cooper and Mahony, 1935: 846
	9II		U.S.A., U.S.S.R., Sweden	Whitaker and Jagger, 1939: 299
		18	U.S.A.	Heiser and Whitaker, 1948: 185
		18	Canada	Mulligan, 1957: 781 sub <i>L. scariola</i> L.
		18	cultivated	Chatterjee and Sharma, 1969: 580
		18	Afghanistan	Podlech and Dieterle, 1969: 230
		18	Slovenia	Feráková, 1970a: 13
		18	Portugal	Fernandes and Queirós, 1971: 84
		18	India	Koul and Gohil, 1973: 61
		18	Portugal	Queirós, 1973: 311
		18	Britain	Edmonds et al., 1974: 160
		18	Bulgaria	Feráková and Murin, 1976: 491
	9II		U.S.A.	Tomb et al., 1978: 719
		18	Italy	Brullo et al., 1979: 146
		18	Greece	Strid and Franzén, 1981: 842
		18	Britain	Haque and Godward, 1985a: 730
9II		Britain	Haque and Godward, 1985b: 841	
9II		India	Mehra et al., 1965: 44	
<i>L. saligna</i> L.	9II		U.S.A.	Thompson et al., 1941: 93
		18	U.S.A.	Heiser and Whitaker, 1948: 185
		18	Portugal	Fernandes and Queirós, 1971: 84
<i>L. virosa</i> L.	9II		Netherlands	Thompson et al., 1941: 93
		18	Portugal	Fernandes and Queirós, 1971: 84
		18	France	Kliphuis and Wieffering, 1972: 600
		18	Portugal	Queirós, 1973: 311
		18	Italy	Brullo et al., 1979: 146
	9II		Spain	Gallego, 1983: 132
		18	Spain	Blanca and Cueto, 1984: 185
		18	Spain	Fernández et al., 1985: 301
	18	Madeira	Dalgaard, 1986: 231	
<i>L. livida</i> Boiss. and Reuter		18	Spain	Blanca and Cueto, 1984: 185

Tab. 2. (continued)

Taxa	n	2n	Origin of material	Authors
<i>L. perennis</i> L. subsp. <i>perennis</i>	9II 9II		England cultivated	Whitaker and Jagger, 1939: 299 Thompson et al., 1941: 93
		18	Hungary	Pölya, 1949: 134
		18	Austria	Leute, 1974: 812
		18	Italy	Löve and Löve, 1982b: 584
subsp. <i>granatensis</i> Charpin and Fernández Casas		18	Spain	Charpin and Fernandez Casas, 1981: 237
<i>L. tenerrima</i> Pourret		16	Spain	Ubera and Ruiz de Clavijo, 1984: 296
		16+	Spain	Cueto and Blanca, 1987: 405
		1B, 2B		

### *Size of the chromosomes*

The apparent chromosome length of the Iberian representatives of *Lactuca* varies from about 1.07–6.33  $\mu\text{m}$  in the present study. Therefore, they are small, medium small and medium large; the class medium small being the most-represented. *L. tenerrima* shows the karyotypes with the smallest chromosomes, since in no case were they found to be longer than 2.57  $\mu\text{m}$ . In the remaining taxa only 1 or 2 pairs, if any, are small, the rest are basically medium small, and, in some cases, 1 or 2 medium large pairs are present. Considerable levels of variability were found within taxa, which in some cases can be attributed to different degrees of condensation, although special care was taken to homogenize the period of antimetabolic treatment.

The  $A_2$  index of interchromosomal variability ranges from 0.16 to 0.38. The most symmetrical species in this respect are *L. tenerrima* and *L. perennis*, representatives of the subsect. *Cyaniceae* of the sect. *Lactuca*. Higher indices were found in the representatives of the subsect. *Lactuca* from the same section: *L. serriola*, *L. saligna*, *L. virosa* and *L. livida*. All the subspecies of *L. viminea* (sect. *Phoenixopus*) show higher interchromosomal asymmetry levels, always exceeding 0.30. In this latter case, the constant presence of a pair of chromosomes longer than the others, and easily recognizable, was observed.

### Discussion

The chromosome numbers presented here for the genus *Lactuca* agree with previous counts made by various authors in all the taxa (Table 2) except *L. viminea* subsp. *ramosissima*, the chromosome number of which is reported for the first time (Table 3). Nevertheless, some specific points can be made: the formation of 9 bivalents has been observed in the meiotic studies of *L. serriola*, in disagreement with Cooper and Mahony (1935: 846) who indicated the formation of 7 bivalents and 1 tetravalent during meiosis

Tab. 3. Chromosome numbers and geographical distribution of *Lactuca* taxa. Distribution areas have been taken from "Index Kewensis" (Hooker and Jackson, 1946 and following) and modified according to Britton and Brown (1970), Feráková (1976; 1977) and Vuilleumier (1973). Authors: (1) Babcock et al., 1937 sec. Feráková, 1977: 85. (2) Blanca and Cueto, 1984: 185. (3) Charpin and Fernández Casas, 1981: 237. (4) Chatterjee and Sharma, 1969: 580. (5) Ching-I-Peng and Chien-Chang Hsu, 1977: 564. (6) Dvorák et al., 1981: 844. (7) Feráková, 1970a: 13–14. (8) Feráková, 1970b: 413. (9) Ghaffari, 1986: 901. (10) Gupta and Gill, 1980: 352. (11) Hanelt, 1973: 649. (12) Ishikawa, 1911 sec. Ishikawa, 1916: 438. (13) Ishikawa, 1916: 438. (14) Ishikawa, 1921: 156–157. (15) Keil and Pinkava, 1981: 705. (16) Kuzmanov et al., 1981: 702. (17) Laurenko et al., 1988: 606. (18) Laurenko et al., 1990: 1623. (19) Löve and Löve, 1975b: 674. (20) Löve and Löve, 1982a: 359–360. (21) Lungeanu, 1972: 682. (22) Malla et al., 1977: 446. (23) Martinoli, 1954: 79. (24) Mehra et al., 1965: 44. (25) Montmollin, 1986: 437. (26) Mulligan, 1957: 781. (27) Murín, 1971: 353. (28) Nazarova in Feráková. 1977: 85. (29) Nazarova, 1984: 972. (30) Ono, 1955: 270. (31) Ortega and Navarro, 1977: 76. (32) Packer, 1964: 483. (33) Parfitt, 1981: 516. (34) Provatova and Sokolovskaya, 1990: 1620. (35) Shetty, 1967: 570. (36) Sokolovskaya and Probatova, 1986: 1423. (37) Sokolovskaya et al., 1985: 126. (38) Stebbins et al., 1953: 427. (39) Strid and Franzén, 1981: 842. (40) Thompson et al., 1941: 93. (41) Tomb et al., 1978: 719. (42) Ubera and Ruiz de Clavijo, 1984: 296. (43) Whitaker and Jagger, 1939: 299. (44) Zakharyeva, 1985: 1699

Taxa	Geographical distribution	n	2n	Authors
<i>L. acanthifolia</i> (Willd.) Boiss.	Aegean Region		18	(25)
<i>L. altaica</i> Fisch. and Mey.	Altai Region	9		(40)
<i>L. aurea</i> (Schultz Bip. ex Panc.) Stebbins	Balkan Peninsula		16	(1)
<i>L. bourgeauii</i> (Boiss.) Irish and Taylor	Asia Minor	8		(40, 43)
<i>L. brunoniana</i> (Wall.) Clarke = <i>Prenanthes brunoniana</i> Wall.	Himalayan Region, Malaysia	8		(24, 35)
<i>L. canadensis</i> L.	Boreal America		16	(24, 38)
<i>L. cretica</i> Desf.	Asia Minor	8	34	(20, 40, 43)
<i>L. chelidonifolia</i> Makino	Japan	5		(40, 43)
<i>L. chinensis</i> Makino	Japan	16		(14)
<i>L. debilis</i> Benth. and Hook. f.	Japan	24		(14)
<i>L. dentata</i> Robinson				
<i>var. thumbergii</i>	Japan	12		(13, 14)
<i>var. albiflora</i>	Japan	12		(14)
<i>var. alpicola</i>	Japan	7		(14)
<i>L. denticulata</i> Maxim.	Japan	5		(13, 14)
<i>L. dissecta</i> D. Don	Himalayan Region	8		(24, 35)
<i>L. floridana</i> (L.) Gaertn. = <i>L. biennis</i> (Moench) Fern.	Boreal America	17		(33, 40)
<i>L. formosana</i> Maxim.	China, Formosa		34	(20, 26, 32)
<i>L. georgica</i> Grossh.	Transcaucasian Region		18	(5)
<i>L. gracilliflora</i> DC.	Himalayan Region, Malaysia	9		(29)
			18	(4, 24)
<i>L. graeca</i> Boiss.	Greece		18	(4)
<i>L. graminifolia</i> Michx.	Boreal America	17		(39)
<i>L. hastata</i> DC.	Eastern India	8		(15, 40, 41, 43)
			16	(24, 35, 43)
<i>L. indica</i> L.	Malaysia	9		(38)
				(40)

Tab. 3. (continued)

Taxa	Geographical distribution	n	2n	Authors
<i>L. indica</i> L.	Malaysia		18	(5)
<i>L. keiskeana</i> Makino	Japan	5		(14)
<i>L. lanceolata</i> Makino	Japan	5		(13, 14)
<i>L. lessertiana</i> Clarke	Himalayan Region		16	(38)
<i>L. livida</i> Boiss, and Reuter	Iberian Peninsula	9	18	(2)
<i>L. longidentata</i> Moris	Sardinia	8	16	(23)
<i>L. longifolia</i> DC.	Himalayan Region	8		(24, 35)
			16	(24, 38)
<i>L. ludoviciana</i> (Nutt.) Ridell.	Western Boreal America		34	(20)
<i>L. macrantha</i> Clarke	Himalayan Region		16	(38)
<i>L. macrorhiza</i> (Royle) Hook. f.	Himalayan Region	8		(24, 35)
		9		(24)
<i>L. matsumaruae</i> Makino	Japan	8		(14)
<i>L. orientalis</i> Boiss. = <i>Scariola orientalis</i> (Boiss.) Soják	Himalayan Region, Asia Minor		18	(9, 27, 29)
<i>L. palmensis</i> Bolle	Canary Isles	8		(31)
<i>L. perennis</i> L. subsp. <i>perennis</i>	Southern Europe	9	18	(43, see Table 2)
subsp. <i>grantensis</i> Charpin and Fernández Casas	Iberian Peninsula		18	(3)
<i>L. polycephala</i> Benth.	Eastern India, Burma	8		(10)
<i>L. pulchella</i> (Pursh) DC. = <i>L. tatarica</i> subsp. <i>pulchella</i> (Prush) Stebbins	Western Boreal America		34	(20, 38)
<i>L. quercina</i> L. subsp. <i>quercina</i>	Caucasus	9	18	(7, 8)
subsp. <i>wilhelmsiana</i> (Fisch. and Mey., ex DC.) Feráková)	Caucasus		18	(28)
<i>L. raddeana</i> Maxim.	Japan	9		(14, 40)
<i>L. rapunculoides</i> Clarke	Himalayan Region		16	(38)
<i>L. repens</i> Benth. and Hook. f.	?	8		(14)
<i>L. saligna</i> L.	Northern Hemisphere	9	18	(40, see Table 2)
<i>L. sagittarioides</i> Clarke	Himalayan Region	5		(22)
<i>L. sativa</i> L.	cultivated	9		(40, 43)
			18	(43)
<i>L. serriola</i> L.	Northern Hemisphere	9	18	(43, see Table 2)
<i>L. sibirica</i> (L.) Benth.	Siberia		18	(19)
			27	(17, 18, 36)
<i>L. sororia</i> Miq.	Japan, China		18	(5)
<i>L. spicata</i> (Lam.) Hitchc.	Northern America	17		(40)
<i>L. squarrosa</i> Miq.	Eastern Asia	9		(30)
			18	(30, 34)
<i>L. stolonifera</i> Benth, and Hook. f.	Japan	8		(14)
<i>L. tamagavensis</i>	?	8, 7		(14)
<i>L. tatarica</i> (L.) Mey. = <i>L. tatarica</i> (Mey.) Nijar	Western Asia, Eastern India	9		(40)
			18	(6, 11, 16, 21, 43)
<i>L. takhtadzhianii</i> Sosn.	?		18	(44)
<i>L. tenerrima</i> Pourret	Southern Europe, Morocco	8	16	(42, see Table 2)

Tab. 3. (continued)

Taxa	Geographical distribution	n	2n	Authors
<i>L. thumbergii</i> Maxim.	Japan	12, 11		(12)
<i>L. triangulata</i> Maxim.	Eastern Siberia	9		(14)
			18	(37)
<i>L. undulata</i> Ledeb.	Eastern an Boreal Asia	9		(9)
<i>L. villosa</i>	?	9		(13, 14)
<i>L. viminea</i> (L.) J. and C. Presl				
subsp. <i>viminea</i>	Northern Hemisphere	9	18	(7, see Table 2)
subsp. <i>alpestris</i> (Gand.) Feráková	Crete		18	(25)
subsp. <i>chondrilliflora</i> (Boreau) Bonnier	Mediterranean Region	9	18	(43, see Table 2)
subsp. <i>ramosissima</i> (All.) Bonnier	Mediterranean Region	9	18	(hic)
<i>L. virosa</i> L.	Northern Hemisphere	9	18	(40)

for this species. Also, no B chromosomes were detected in the populations of *L. tenerima* studied, while Cueto and Blanca (1987: 405) indicated the presence of B chromosomes in plants of this species from Sierra de Baza (Granada, Spain). Table 4 gives the chromosome numbers of the Iberian taxa of *Lactuca* and many other representatives of the genus from around the world, including their geographic distribution. A high diversity can be seen in numbers including  $2n = 10, 14, 16, 18, 22, 24, 32,$  and  $34$ , which display an interesting geographic pattern. The representatives from the Mediterranean Region, that encompasses the highest number of representatives of Europe and North Africa, have mainly the number  $2n = 18$ , but two taxa, *L. aurea* and *L. tenerima*, have  $2n = 16$ . No incidence of polyploidy has been found in this area. The number  $2n = 16$  is the most common number found among the representatives from the Himalaya, India and Malaysia. In North America, a group of species occurs sharing the number  $2n = 34$ . It has been proposed to be composed of amphiploids originating from species with  $2n = 18$ , and  $2n = 16$  (Whitaker and Thompson 1941) from Eurasia that colonized North America via the Bering Strait (Whitaker 1944, Vuilleumier 1973). It seems that all of them have a common origin since a high interfertility level was found (Thompson et al. 1941, Whitaker and Jagger 1939, Whitaker and Thompson 1941). Some additional taxa showing the number  $2n = 18$  are also distributed in this region, but they are presumed to have been introduced from the Old World (Vuilleumier 1973). The representatives from Japan form the most diverse group from the karyological point of view since all the numbers are represented. Some of these correspond to endemic taxa studied by Ishikawa (1916, 1921) but not later complete confirmation of his observations has been reported. According to Tomb (1977) these results are questionable.

Each of these regions is related to different centres of speciation of the genus. The number  $n = 9$ , considered the basic one of the tribe (Stebbins et al. 1953, Tomb et al. 1978), although Turner et al. (1961) postulated  $x = 4$  or  $x = 5$  as the ancestral basic number in the Lactuceae, prevails only in the Mediterranean Region, that has been considered as centre of origin of *Lactuca* by Feráková (1977). In this area 4 sections of the genus *Lactuca* can be found: *Mulgedium* (Cass.) Clarke, *Lactucopsis* (Schultz-Bip ex Vis. et Panc.) Rouy, *Phoenixopus* (Cass.) Benth. and *Lactuca* L.; all four are represented in the Eastern part of the area, while in the West only the last two sections can be found.

The Eastern Mediterranean Region is probably a centre of origin of the genus. Babcock et al. (1937 sec. Stebbins et al. 1953) considered both  $x=8$ , and  $x=9$ , as basic chromosome numbers in *Lactuca* and proposed a polyphyletic origin for the genus, but Stebbins et al. (1953) considered  $x=9$  as the only primitive one, arguing that the number  $n=8$  had not been found in *Cicerbita*. In fact, although some species of *Cicerbita* showing  $n=8$  have been reported (e.g. Bolkhoskikh et al. 1969), they could hardly be considered as ancestors of *Lactuca*. In the area of the Himalayas, India and Malaysia, where a high incidence of the number  $n=8$  has been found, the reduction of the chromosome number should take place during an early evolutionary stage. Nevertheless, the high diversity of the genus and the geographical pattern of distribution allows a polyphyletic origin to think about, that is illustrated by the taxonomic study of a small group of scandent species of *Lactuca* from Africa, not related to any other taxa of the genus, which probably originated in this area from an isolated section of *Prenanthes* (Stebbins 1937). Unfortunately no karyological studies are on record.

As indicated before (see Introduction) chromosomes of *Lactuca* are quite appropriate for studies of chromosome morphology but few references can be found on this subject, except on *L. serriola*, the chromosome set of which had been studied before by Haque and Godward (1985a). These authors proposed the idiogrammatic formula  $8m + 2m - sm + 6sm^{sat} + 2st^{sat}$  for British plants, a formula very similar to our results. The asymmetry type found by these authors was 3A, and the asymmetry coefficients calculated in the work presented here from the arm length ratios are  $A_1 = 0.44$  and  $A_2 = 0.21$ . They also indicated that *L. serriola* and *L. sativa* were karyologically very similar. Certain other authors (Chatterjee and Sharma 1989, Whitaker and Jagger 1939) had made observations on the chromosome morphology of these taxa but nomenclature differences preclude a comparison. In addition, it has been possible to study the idiogrammatic formula of *L. virosa* and *L. livida* from the detailed photographs published by Blanca and Cueto (1984). In both cases the formula deduced is  $2M + 8sm + 2sm^{sat} + 4st + 2t$  (Table 1), which coincides with our observations.

It is interesting that the study of the karyotypes in the Iberian representatives reflects their taxonomic relationships (Table 1) and could also be an useful tool in certain taxonomic considerations. Three groups can be separated: 1) *L. viminea*, represented by three subspecies with very similar idiogrammatic formula and karyotype asymmetry; 2) *L. serriola*, *L. saligna*, *L. virosa*, and *L. livida* which are widespread weed representatives of the sect. *Lactuca* subsect. *Lactuca*, an evolved group in the genus. They present the most asymmetrical karyotypes of the Iberian taxa as could be expected to occur in colonizer weeds (Stebbins 1971: 90); 3) *L. perennis* and *L. tenerrima*, included in the subsect. *Cyanicae* sect. *Lactuca*, are very closely related morphologically. Superficially, their chromosomes seem quite different because of their lengths and some secondary constrictions found in *L. tenerrima* (Fig. 12 and 13), but in both cases, the karyotype asymmetry is 2A, and  $A_1$  and  $A_2$  coefficients are very similar (Table 1).

*L. viminea* presents pronounced morphological diversity comprising four subspecies treated as species by many authors (e.g., Willkomm 1861, Coste 1903) and considered well-marked subspecies by Feráková (1977). Three of them are represented in the Iberian Peninsula, but not clear morphological and taxonomic delimitations have been found in the material studied. Some characters currently used may be fixed, e.g. the length of the beak in the cypsela but some others, such as the mode of ramification, may depend on climatic features and sun exposure. Based on the karyological data no differentiation is possible since the diversity found is as similar between subspecies as it is within subspecies (Table 1).

Various authors have dealt with the taxonomic status of *L. livida* Boissier and Reuter. Blanca and Cueto (1984) indicated that *L. livida* is a species clearly differentiated from *L. virosa*. In contrast, Linqvist (1960) and Feráková (1977) considered them as closely related species without clear differential characters, and Velasco (1981) proposed the new combination *L. virosa* subsp. *livida* (Boissier and Reuter) Ladero and Velasco. In fact, the number of florets per flower head, the shape of the leaves, and the anthocyanin purple colouring, considered as the most important differential characters in the taxonomic diagnosis, have been found to show a continuous variation in the area, and sometimes to be different in wild populations, and in the material from these populations growing in the experimental fields. Also, no important karyological differences have been found among the material studied as can be seen in Table 1. It is reasonable to assume that a thorough taxonomic revision is necessary for clarifying definitively the taxonomic boundaries and status of these taxa. Plants with some typical characters of *L. livida* can be found in several separate localities of the Iberian Peninsula, which indicates that their distribution area is wider than usually presumed (Feráková 1976: 330, 1977: 75). In fact, although Feráková indicated that the distribution of *L. livida* is restricted to a small area in the centre of Spain, Velasco (1981) estimated that it has a wider distribution.

The author is grateful to R. Churchill, who corrected the English of the manuscript.

## References

- Blanca G. and M. Cueto 1984. Números cromosómicos de plantas occidentales, 290–296. *Anales Inst. Bot. Cavanilles* 41: 185–189.
- Boissier E. 1839–1845. *Voyage botanique dans le midi de l'Espagne*, 2. Guide et Cie., Paris. 757 pp.
- Bolkhoskikh Z., Grif V., Matvejeva T. and Zarharyeva O. 1969. Chromosome numbers of flowering plants. Leningrad. 926 pp.
- Britton N. L. and H. A. Brown 1970. *An illustrated flora of the Northern United States and Canada*. Dover Publ. New York. 637 pp.
- Brullo S. and P. Pavone 1978. Numeri cromosomici per la flora italiana: 464–483. *Inform. Bot. Ital.* 18: 248–266.
- Brullo S., P. Pavone and A. Zizza 1979. Numeri cromosomici per la flora italiana: 602–616. *Inform. Bot. Ital.* 11: 139–147.
- Charpin A. and J. Fernández Casas 1981. *Iter secundum austrohispanicum*. *Candollea* 36: 223–241.
- Chatterjee T. and A. K. Sharma 1969. Cytotaxonomy of Cichorieae. *Genetica* 40: 577–590.
- Ching-I-Peng and Chien-Chang Hsu 1977. In Á. Löve (ed): IOPB chromosome number reports, LVIII. *Taxon* 26: 562–564.
- Cooper D. C. and K. L. Mahony 1935. Cytological observations on certain Compositae. *Amer. J. Bot.* 22: 843–848.
- Coste H. 1903. *Flore descriptive e illustrée de la France*, vol. 2. Paul Klincksieck, Paris. 627 pp.
- Cueto M. and G. Blanca 1987. Números cromosómicos de plantas occidentales, 392–402. *Anales Jard. Bot. Madrid* 43: 403–409.
- Dalgaard V. 1986. Chromosome numbers in flowering plants from Madeira. *Willdenowia* 16: 221–240.
- Dvorák F., F. Grüll, K. Sutory and B. Dadáková 1981. In Á. Löve (ed.): IOPB chromosome number reports, LXXIII. *Taxon* 30: 844.
- Edmonds J. M., P. D. Sell and S. M. Walters 1974. Some British chromosome counts in the Compositae sub-family Cichorioideae. *Watsonia* 10: 159–161.
- Feráková V. 1970 a. In J. Májovský et al. (eds.): Index of chromosome numbers of Slovakian flora. Part 1. *Acta F. R. N. Univ. Conmen. – Botanica* 16: 1–26.
- Feráková V. 1970 b. A biosystematic study of critical species of *Lactuca* sect. *Lactucopsis*. *Folia Geobot. Phytotax.* 5: 401–427.

- Feráková V. 1976. Genus *Lactuca* L. In G. Tutin et al. (eds.): Flora Europaea, vol. 4: 328–331. Cambridge Univ. Press, Cambridge.
- Feráková V. 1977. The genus *Lactuca* L. in Europe. Univerzita Komenského, 122 pp.
- Feráková V. and A. Murin 1976: IOPB chromosome number reports, LIII. Taxon 25: 490–491.
- Fernandes A. and M. Queirós 1971. Contribution à la connaissance cytotaxonomique des Spermatophyta du Portugal. II: Compositae. Bol. Soc. Brot., Sér. 2, 4: 5–121.
- Fernandez I., M. J. Díez and J. Pastor 1985. Números cromosómicos para la flora española, 373–381. Lagasalia 13: 299–302.
- Gallego M. J. 1983. Números cromosómicos para la flora española, 290–294: Lagasalia 12: 131–132.
- Ghaffari S. M. 1986. In Á. Löve (ed.): IOPB chromosome number reports, XCIII. Taxon 35: 900–901.
- Gupta R. C. and B. S. Gill 1980. In Á. Löve (ed.): IOPB chromosome number reports, LXVII. Taxon 29: 351–352.
- Hanelt P. 1973. In Á. Löve (ed.): IOPB chromosome number reports, XLII. Taxon 22: 648–649.
- Haque M. Z. and M. B. E. Godward 1985 a. Comparison between two genera, species and cultivars in Lactuceae. I. Karyotype analysis. Cytologia (Tokyo) 50: 725–738.
- Haque M. Z. and M. B. E. Godward 1985 b. Comparison between two genera, species and cultivars in Lactuceae. II. Effects of seed-irradiation on chiasma frequency. Cytologia (Tokyo) 50: 841–853.
- Heiser C. B. and T. W. Whitaker 1948. Chromosome numbers, polyploidy and growth habit in California weeds. Amer. J. Bot. 35: 179–186.
- Hooker J. D. and B. D. Jackson 1946 sqq. Index Kewensis, 2 vols. and 15 supplements. Oxford Univ. Press. Oxford.
- Ishikawa M. 1916. A list of the number of chromosomes. Bot. Mag., Tokyo 20: 404–448.
- Ishikawa M. 1921. On the chromosomes of *Lactuca*. Bot. Mag., Tokyo 35: 153–158.
- Keil D. J. and D. J. Pinkava 1981. In Á. Löve (ed.): IOPB chromosome number reports, LXXII. Taxon 30: 705–706.
- Kliphuis E. and J. H. Wieffering 1972. Chromosome numbers of some angiosperms from the south of France. Acta Bot. Neerl. 21: 598–604.
- Koul A. K. and R. N. Gohil 1973. Cytotaxonomical conspectus of the flora of Kashmir (1). Chromosome numbers of some common plants. Phytion (Horn) 15: 57–66.
- Küpfer P. 1974. Recherches sur les liens de parenté entre la flore orophile des Alpes et celle des Pyrénées. Boissiera 23. 321 pp.
- Kuzmanov B. and S. Georgieva 1980. In Á. Löve. (ed.): IOPB chromosome number reports, LXIX. Taxon 29: 715.
- Kuzmanov B., S. Georgieva, V. Nikolova and I. Penceva 1981. In Á. Löve (ed.): IOPB chromosome number reports, LXXII. Taxon 30: 701–702.
- Laurenko A. N., N. P. Serditov and Z. G. Ulle 1988. Chromosome numbers in some species of Asparagaceae, Asteraceae and Ranunculaceae from the North Urals (Kuomi ASSR) Bot. Žur., 73: 605–607.
- Laurenko A. N., N. P. Serditov and Z. G. Ulle 1990. Chromosome numbers in some species of flowering plants of the Urals (Kuomi ASSR). Bot. Žur., 75: 1622–1624.
- Leute G.-H. 1974. In Á. Löve (ed.): IOPB chromosome number reports, XLVI. Taxon 23: 811–812.
- Levan A., K. Fredga and A. A. Sandberg 1965. Nomenclature for centromeric position on chromosomes. Hereditas 52: 201–220.
- Lindqvist K. 1960. On the origin of cultivated lettuce. Hereditas 46: 319–350.
- Löve Á and D. Löve 1975 a. Plant chromosomes. J. Cramer. Vaduz. 125 pp.
- Löve Á. and D. Löve 1975 b. In Á. Löve (ed.): IOPB chromosome number reports, L. Taxon 24: 673–675.
- Löve Á. and D. Löve 1982 a. In Á. Löve (ed.): IOPB chromosome number reports, LXXV. Taxon 31: 344–360.
- Löve Á. and D. Löve 1982 b. In Á. Löve (ed.): IOPB chromosome number reports, LXXVI. Taxon 31: 583–587.
- Lungeanu I. 1972. In Á. Löve (ed.): IOPB chromosome number reports, XXXVIII. Taxon 21: 682.

- Malla S. B., S. Bhattarai M. and Gorkhali 1977. In Á. Löve (ed.): IOPB chromosome number reports, LVII. *Taxon* 26: 444–446.
- Martinoli G. 1954. La cytotaxonomie expérimentale appliquée aux espèces végétales de la Sardègne et en particulier aux endémiques. VIII Congr. Intern. Bot., Compt. Rend. Sci. et Rapp. et Comm. parvenus avant le Congrès, séct. 8 et 16: 78–79.
- Mehra P. N., B. S. Gill, J. K. Mehta and S. S. Sidhu 1965. Cytological investigations on the Indian Compositae. I. North-Indian taxa. *Caryologia* 18: 35–68.
- Montmollin B. 1986. Etude cytotaxonomique de la flore de la Crète. III. Nombres chromosomiques. *Candollea* 41: 431–439.
- Mulligan G. A. 1957. Chromosome numbers of Canadian weeds. I. *Canad. J. Bot.* 35: 779–789.
- Murin A. 1971. In Á. Löve (ed.): IOPB chromosome number reports, XXXII. *Taxon* 20: 353.
- Nazarova E. A. 1984. Chromosome numbers in the Caucasian representatives of the families Asteraceae, Brassicaceae, Fabaceae, Limoniaceae. *Bot. Žur.* 69: 972–975.
- Ono H. 1955. Intergeneric hybridization in Cichorieae. *Jap. J. Genet.* 30: 269–274.
- Ortega J. and B. Navarro 1977. Estudios en la flora de Macaronesia: algunos números de cromosomas. III. *Bot. Macaronésica* 3: 73–80.
- Packer J. G. 1964. Chromosome numbers and taxonomic notes on western canadian and arctic plants. *Canad. J. Bot.* 42: 473–494.
- Parfitt B. D. 1981. In Á. Löve (ed.): IOPB chromosome number reports, LXXI. *Taxon* 30: 515–516.
- Podlech D. and A. Dieterle 1969. Chromosomenstudien an afghanischen Pflanzen. *Candollea* 24: 185–243.
- Pólya L. 1949. Chromosome numbers of some hungarian plants. *Acta Geobot. Hung.* 6: 124–137.
- Probatova N. S. and A. P. Sokolovskaya 1990. Chromosome numbers in some representatives of the families Asclepiadaceae, Asteraceae, Boraginaceae, Chenopodiaceae, Lamiaceae, Onagraceae, Scrophulariaceae, Solanaceae, Urticaceae from the Soviet Far East. *Bot. Žur.*, 75: 1619–1622.
- Queirós M. 1973. Contribuição para o conhecimento citotaxonomico das Spermatophyta de Portugal. II. Compositae. Supl. 1. *Bol. Soc. Brot.*, Sér 2, 47: 299–314.
- Romero C. 1986. A new method for stimating karyotype asymmetry. *Taxon* 35: 526–530.
- Shetty B. V. 1967. In Á. Löve (ed.): IOPB chromosome number reports, XIV. *Taxon* 16: 552–571.
- Sokolovskaya A. P. and N. S. Probatova 1986. Chromosome numbers in some representatives of the Asteraceae, Iridaceae, Poaceae, Primulaceae, Violaceae families from the Far East of the USSR. *Bot. Žur.* 71: 1423–1425.
- Sokolovskaya A. P., N. S. Probatova and E. G. Rudyka 1985. Chromosome numbers in the species of the families Asteraceae, Poaceae, Rosaceae from the Primoyre region, Kamchatka and Sakhalin. *Bot. Žur.* 70: 126–128.
- Stebbins G. L. 1937. The scandent species of *Prenanthes* and *Lactuca* in Africa. *Bull. J. B. Brux.* 14: 333–352.
- Stebbins G. L. 1938. Cytological characteristics associated with the different growth habits in the dicotyledons. *Amer. J. Bot.* 25: 189–198.
- Stebbins G. L. 1953. A new classification of the tribe Cichorieae, family Compositae. *Madroño* 12: 65–81.
- Stebbins, G. L. 1971. Chromosomal evolution in higher plants. Edward Arnold Ltd. London. 216 pp.
- Stebbins G. L., J. A. Jenkins and M. S. Walters 1953. Chromosomes and phylogeny in the Compositae, tribe Cichorieae. *Univ. Calif. Publ. Bot.* 26: 401–430.
- Strid A. and R. Franzén 1981. In Á. Löve (ed.): Chromosome number reports, LXXIII. *Taxon* 30: 829–842.
- Thompson R. C., T. W. Whitaker and W. F. Kosar 1941. Interspecific genetic relationships in *Lactuca*. *J. Agric. Res.* 63: 91–107.
- Tjio J. H. and A. Levàn 1950. The use of oxyquinoline in chromosome analysis. *Anal. Est. Exper. Aula Dei* 2: 21–64.
- Tomb A. S. 1977. Lactuceae-systematic review. In Heywood V. H. et al. (ed.): The biology and chemistry of the Compositae 2: 1067–1079. Academic Press, London.

- Tomb A. S., K. L. Chambers, D. W. Kyhos, A. M. Powel and P. H. Raven 1978. Chromosome numbers in the Compositae: 14. Lactuceae. Amer. J. Bot. 65: 717–721.
- Turner B. L., Ellison W. L. and R. M. King 1961. Chromosome numbers in the Compositae. IV. North American species with phyletic interpretations. Amer. J. Bot. 48: 216–223.
- Ubera J. L. and E. Ruiz de Clavijo 1984. Números cromosómicos para la flora española, 342–348. Lagasalia 12: 294–296.
- Velasco A. 1981. Sobre *Lactuca livida* Boiss and Reuter. Anales Jard. Bot. Madrid 38: 316.
- Vuilleumier B. S. 1973. The genera of Lactuceae (Compositae) in the southeastern United States. J. Arnold Arbor. 54: 42–93.
- Whitaker T. W. 1944. The inheritance of certain characters in a cross of two american species of *Lactuca*. Bull. Torrey Bot. Club 71: 347–355.
- Whitaker T. W. and J. C. Jagger 1939. Cytogenetic observations in *Lactuca*. J. Agr. Research, Washington 58: 297–306.
- Whitaker T. W. and R. C. Thompson 1941. Cytological studies in *Lactuca*. Bull. Torrey Bot. Club 68: 388–394.
- Wickbom T. 1949. The time factor of chromosome spiralization. Hereditas 25: 245–248.
- Willkomm M. 1870. Compositae L. In M. Willkomm and J. Lange (eds.): Prodrum Florae Hispanicae 2: 24–272. Sumtibus E. Schweizerbart, Stuttgartiae.
- Zakharyeva O. I. 1985. Chromosome numbers of some flowering plants from the Caucasus and Middle Asia. Bot. Žur., 1699–1701.

#### Appendix. – Origin of the material studied (and voucher specimens)

*Lactuca viminea* subsp. *viminea*: CACERES: Entre Jerte y Tornavacas; 15. IX. 1985; García y Mejías (SEV 126388). LEON: Escaro; 22. IX. 1985; García y Mejías (SEV 126389). SEVILLA: Cazalla de la Sierra; 10. VIII. 1986; Herrera, Mejías y Moreno (SEV 126384).

*Lactuca viminea* subsp. *chondrilliflora*: GERONA: Rosas, Playa de La Pelosa; 7. VIII. 1985; Mejías, Polo y Romero (SEV 126381). GERONA: Entre La Molina y Alp; 10. VIII. 1985; Mejías, Polo y Romero (SEV 126380). Entre Viella y Bossost; 11. VIII. 1985; Mejías, Polo y Romero (SEV 126382).

*Lactuca viminea* subsp. *ramosissima*: ASTURIAS: Entre Tineo y Cangas de Narcea; 14. VIII. 1987; Barrera y Mejías (SEV 126385). CADIZ: Villaluenga del Rosario; 7. IX. 1985; Herrera, Mejías y Muñoz (SEV 126383). GRANADA: Puerto de la Ragua; VIII. 1985; Muñoz (SEV 126386). Sierra de Baza, Calar de Rapa; 29. VII. 1987; Mejías y Talavera (SEV 126579). HUELVA: Puerto de Alájar; 1. IX. 1985; Mejías (SEV 126379). LOGROÑO: Entre Viniegra de Arriba y Viniegra de Abajo; 18. IX. 1985; García y Mejías (SEV 126378). LEON: Valdecastillo; 22. IX. 1985; Gracia y Mejías (SEV 126387). SEVILLA: Peñón de Algámitas; 25. VIII. 1986; Mejías y Muñoz (SEV 126377). Entre Cazalla de la Sierra y Constantina; 10. VIII. 1986; Herrera y Mejías (SEV 126384).

*Lactuca serriola*: BARCELONA: Entre Orrius y La Roca; 6. VIII. 1985; Mejías, Polo y Romero (SEV 126524). HUESCA: Lanavé; 13. VIII. 1985; Mejías, Polo y Romero (SEV 126527). JAEN: Entre Villanueva del Arzobispo y Beas de Segura; 2. VIII. 1985; Mejías, Polo y Romero (SEV 126528). LEON: Grajal; 23. IX. 1985; García y Mejías (SEV 126529). SEVILLA: Brenes; 12. VIII. 1989; Mejías (SEV 126584). Sevilla; 12. VIII. 1989; Mejías (SEV 126585).

*Lactuca saligna*: ALAVA: Elciego; 19. IX. 1985; García y Mejías (SEV 126538). ALBACETE: Villapalacios; 2. VIII. 1985; Mejías, Polo y Romero (SEV 126533). CADIZ: Grazalema, Puerto del Boyar; 23. IX. 1984; Herrera y Mejías (SEV 126535). VALLADOLID: Otero de Sariegos; 23. IX. 1985; García y Mejías (SEV 126537).

*Lactuca virosa*: ALBACETE: Entre Bienservida y Villaverde de Guadalimar; 18. VII. 1985; Arroyo y Mejías (SEV 124631). ASTURIAS: La Espina; 14. VIII. 1987; Barrera y Mejías (SEV 126567). AVILA: Puerto del Pico; 17. IX. 1985; García y Mejías (SEV 126564). BARCELONA: Entre Orrius y La Roca; 6. VIII. 1985; Mejías, Polo y Romero (SEV 126594). GRANADA: Sierra de Baza; 27. VII. 1984; Mejías (SEV 124656); HUELVA: Entre Los Marines y Aracena; 12. VII. 1984; Mejías (SEV 124667). Fuenteheridos; 1. IX. 1985; Mejías (SEV 126555). HUESCA: Jaca; 13.

VIII. 1985; Mejías, Polo y Romero (SEV 125558). JAEN: Puerto de Tíscar; 18. VIII. 1984; Herrera y Mejías. Sierra de Cazorla; Torcal Llano; 20. VII. 1984; Herrera y Mejías (SEV 124664). SANTANDER: Comillas; 21. IX. 1985; García y Mejías (SEV 126566). SEVILLA: San Nicolás del Puerto; 18. VIII. 1985; Mejías (SEV 126558).

*Lactuca livida*: TOLEDO: San Pablo de los Montes, Baños del Sagrario; 20. VII. 1985; Arroyo y Mejías (SEV 124612).

*Lactuca perennis* subsp. *perennis*: GERONA: Entre Rosas y Cadaqués; 20. V. 1987; Mejías (SEV 126546). Idem; 20. X. 1988; Mejías (SEV 126547); LERIDA: Puerto de la Bonaigua; 10. VIII. 1985; Mejías, Polo y Romero (SEV 126542). Idem; 1. VII. 1987; Díaz, López y del Pino (SEV 126544).

*Lactuca perennis* subsp. *granatensis*: GRANADA: Sierra de la Sagra; 31. VII. 1988; Mejías (SEV 126591).

*Lactuca tenerrima* Pourret: ALICANTE: Villajoyosa; 5. VI. 1985; García y Mejías (SEV 124616). CADIZ: entre Zahara de la Sierra y el Puerto de las Palomas; 25. VI. 1986; Fernández, García, Mejías y Pastor (SEV 126572). GERONA: Bruguera; 8. VIII. 1985; Mejías, Polo y Romero (SEV 126574). HUELVA: Linares de la Sierra; 1. IX. 1985; Mejías (SEV 126575). Huesca: Valle de Añisclo; 9. IX. 1986; Mejías y Muñoz (SEV 126571). JAEN: Belerdas; 18. VII. 1984; Herrera y Mejías (SEV 124653). Sierra de Cazorla, Torcal Llano; 18. VII. 1984; Herrera y Mejías (SEV 124646). LEON: Oseja de Sajambre; 22. IX. 1985; García y Mejías (SEV 124663).