Zeitschrift:	Botanica Helvetica
Herausgeber:	Schweizerische Botanische Gesellschaft
Band:	103 (1993)
Heft:	2
Artikel:	Reproductive biology of Petrocoptis crassifolia Rouy (Caryophyllaceae), a chasmophilous endemic plant of the Central Pyrenees
Autor:	García, María B. / Antor, Ramón J. / Villar, Luis
DOI:	https://doi.org/10.5169/seals-71333

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

# Download PDF: 22.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# Reproductive biology of *Petrocoptis crassifolia* Rouy (Caryophyllaceae), a chasmophilous endemic plant of the Central Pyrenees

# María B. García, Ramón J. Antor and Luis Villar

Instituto Pirenaico de Ecología (C.S.I.C.), Apdo. 64, E-22700 Jaca, Huesca, Spain

Manuscript accepted April 26, 1993

# Abstract

García, M. B., Antor, R. J., and Villar, L. 1993 Reproductive biology of *Petrocoptis* crassifolia Rouy (Caryophyllaceae), a chasmophilous endemic plant of the Central Pyrenees. Bot. Helv. 103: 133–140.

The reproductice biology of *Petrocoptis crassifolia* Rouy (Caryophyllaceae), an endemic plant of the Central Pyrenees (Spain), was studied. Its nectariferous flowers show an ultraviolet pattern in the corolla and an incipient zygomorphism in the arrangement of petals and stamens. *P. crassifolia* may produce seeds by autogamy but in smaller numbers than when it is pollinated by insects (mainly Hymenoptera). The fruit-set ranges between 43-80%, depending on the years and populations studied. Fruits develop towards the rock face, and seeds exhibit myxospermy; both these traits are adaptations that facilitate the deposition of seeds in rocky substrata. Survival of the seedlings in rocky fissures at the end of the first year is 10%. Reproductive traits suggest that this species is well adapted to its habitat, and that it is not endangered. Its rarity seems to be associated mainly with a lack of dispersal mechanisms for colonizing suitable but remote habitats.

Key words: Reproductive biology, Petrocoptis crassifolia, endemic plant, Pyrenees.

# Introduction

The Pyrenees, like other mountains in the Iberian Peninsula, are one of the areas with the largest diversity of vegetation and endemism in Europe (Favarger 1972, Gómez-Campo and Malato-Beliz 1985, Villar and García 1989). Until now, studies of endemic plants in the Pyrenees have focused on taxonomy and biogeography (see, for example, Montserrat and Villar 1972, Küpfer 1974, Delay and Vivant 1978, Vivant and Delay 1980, Baudière and Cauwet-Marc 1987, Villar 1988), and only recently have they begun to focus on biology and ecology (Dajoz 1991, García and Antor 1992, García et al. 1992), even though this kind of knowledge is very important in deciding their status and their possibilities of survival (Gómez-Campo et al. 1987).

Petrocoptis crassifolia Rouy (Caryophyllaceae) is a perennial herb belonging to a genus endemic to the Iberian Peninsula (NE Spain; see Montserrat and Fernández Casas

1989; Saule 1991). It occupies a narrow geographic range (the central Pyrenees) and has a high habitat specificity (crevices of limestone cliffs and overhangs), allowing us to catalogue this species as "rare" (*sensu* Rabinowitz 1981). Since several studies have demonstrated that the rarity of some species is associated with reproductive problems (see, for example, Ross 1936, Davies 1960, Greig-Smith and Sagar 1981, Herrera 1987, 1989), this paper deals with the reproductive biology of this singular endemic plant, and discusses the causes of its rarity.

We have investigated the most important populations of *P. crassifolia*, located throughout the Añisclo Valley (Ordesa-Monte Perdido National Park, Spain) and "Las Devotas" (Lafortunada, Cinca Valley), where we have recorded floral traits, pollinators, dependence on insect visits for reproductive success, fruit set, seed set, and the survival of seedlings.

# **Floral traits**

The flowers of *P. crassifolia* are composed of a tubular calyx and five free petals each of which bear two coronal scales. Although the corolla has a more or less uniform pinkish-purple colour, UV photographs show "nectar-guides" at the mouth of the floral tube within which nectar is found (Fig. 1). This UV pattern, invisible to the human eye but visible to many insects, is due to the absorption of UV wavelengths by coronal scales,



Fig. 1. Visible light (left) and ultraviolet (right) pattern in *P. crassifolia* flower. The right-hand photograph was taken using a filter B+W 403 and a film TMAX 400 ASA (Kodak).



Fig. 2. Example of frequent floral symmetry in *P. crassifolia*: the petals are aggregated in the lower part of the corolla and the anthers in the upper part of the floral tube.

contrasting with the rest of the corolla that reflect them. Detailed UV patterns, composed of reflecting and absorbing regions, can be found in many plant species (Horovitz and Cohen 1972, Jones and Buchmann 1974, Frohlich 1976, Kay 1984), helping to direct the pollinator's behaviour in a manner that assures pollen transfer (Silberglied 1979).

Because this plant grows on vertical and overhanging walls, its stems are usually pendulous, and the petal plates of the corollas are usually vertically oriented (average  $\pm$  standard deviation:  $90^{\circ}39' \pm 8^{\circ}2'$ ; range:  $80^{\circ} - 109^{\circ}$ ; N = 60; calculated following Batschelet 1981). Floral symmetry is not strictly radial, as has usually been assumed, but slightly zygomorphic (Fig. 2). This bilateral symmetry comprises two characteristics: two or three petals are grouped in the lower part of the corolla, and all anthers are located in the upper part of the floral tube. Both stamens and petals are radially arranged at the bottom of the floral tube, but change their orientation within it. This incipient zygomorphy seems to be associated with the vertical position of the petal surfaces and also with the behaviour of the pollinators, because it has been found in other species of the genus that resemble *P. crassifolia* and with similar pollinators (García et al. 1992).

The main reward that flowers of *P. crassifolia* offer to pollinators is nectar, accumulated at the bottom of the floral tube. Nectar measurements were made in cut inflorescences kept in plastic bags for 24 hours at ambient temperature (nectar measured in this way is usually less abundant and more diluted than in the field; see Cruden et al. 1983 for a review of this subject). Each flower produces about two microlitres of nectar per day (average  $\pm$  standard deviation:  $2.3 \pm 1.0 \mu$ ; N=36; measured in calibrated micro-

pipets), and the concentration of sugars is about 16% ( $15.7 \pm 2.0\%$ ; measured with a Bellingham and Stanley pocket refractometer).

# Reproduction

The main pollinators of *P. crassifolia* are Hymenoptera (68% of visits to flowers), principally *Anthophora plumipes* and *Bombus pratorum*, followed by Lepidoptera (27%; principally the hawkmoth *Macroglossum stellatarum*) and Diptera (5%; *Bombylius venosus*). These pollinators are very similar to those visiting *P. grandiflora* (Guitián et al. 1992). All of these were recorded as diurnal visitors during 20 censuses of 15' in 1990 and 1991, but we do not know if *P. crassifolia*, like some *Silene* species, is also pollinated by nocturnal insects (see, for example, Pettersson 1991). These insects hang from the coronal scales (as the petal surface is vertical) in order to suck the nectar from the floral tube.

The fruit-set in *P. crassifolia* ranges from 43% to 80% (ratio fruits/flowers), depending on population and year (Table 1). Each fruit usually contains between 5 and 8 seeds, which means that more than half of the ovules develop into seeds (ratio seeds/ovules: 59.2% during 1991 in Añisclo, N=131 fruits examined). Nevertheless, a given population may have a considerable interannual variability in the fruit-set (Las Devotas II:  $\chi^2 = 98.8$ ; d.f. = 1; p < 0.001), while no statistical difference was found between populations for the same year (1990:  $\chi^2 = 1.39$ ; d.f. = 1; n.s.).

Some fruits can be formed by self-pollination (22% of the 82 flowers that were enclosed in a fine nylon mesh produced capsules); however, in this way *P. crassifolia* produces significantly fewer fruits than when it is pollinated by insects ( $\chi^2 = 558$ ; d.f. = 1; p<0.001). Although autogamy is possible, 53% of self-pollinated flowers did not produce mature fruits although the ovary began to inflate, which indicates that self-compatibility is not totally functional in *P. crassifolia*. In addition, the seed-set was also lower in enclosed flowers ( $3 \pm 2.4$  seeds per fruit, N=14) than in free flowers (see Table 1).

The dispersal of seeds takes place near the rocky substrate, because capsules develop towards the wall, where they open (Dendaletche 1970). The seeds each have a tuft of hairs that become mucilaginous in contact with water (myxospermy), which allows them to stick to humid surfaces. We have occasionally observed that seeds may be dispersed by ants. On the other hand, anemochory seems unlikely because the seeds lack adaptations for this and are relatively heavy (seed mass: 1.5 mgr.; see Westoby et al. 1992).

Year	Population	Fruit-set			Seed-set	
11		N	Fruits	% fruits/flowers	N	$\overline{x} \pm s.d.$ seeds/fruit
1990			and the second second second		-	
	Añisclo I	66	46	69.7	41	$7.8 \pm 2.5$
	Las Devotas I	274	201	73.4	133	$4.7 \pm 2.5$
	Las Devotas II	357	285	79,8	40	$5,1 \pm 2,0$
1991						
1000	Añisclo II	297	165	55,5	131	$7,5 \pm 3,2$
1992	I as Devetes II	220	1.45	10.0		
	Las Devotas II	338	145	42,9		

 

 Table 1. Fruit-set and average number of seeds per fruit of P. crassifolia, in four different populations in three years.

Botanica Helvetica 103/2, 1993



Fig. 3. Percentage of seedling survival of *P. crassifolia* in three different microhabitats during the first year of life.

In some populations throughout the Añisclo Valley, plants located at the bottom of the cliffs (near shrubs) suffered heavy floral predation (up to 100% of the flowers), which entirely prevents seed production. We observed the caterpillar of *Trigonophora flammea* (Lepidoptera, Noctuidae) feeding on flower buds.

## Survival of seedlings

The germination of the seeds takes place early in spring. Seedlings can be seen on the ground, on ledges with some soil, and in rocky crevices. In April of 1991 we marked 242 young seedlings in Añisclo and Las Devotas populations and recorded their microhabitat. During their first year of life they were checked monthly in order to estimate their survival rate.

Spring and summer were the periods with the highest seedling mortality, and survival curves became stable from September onwards (Fig. 3). On the ground all seedlings died in the first two months (N=32), and only 10% of seedlings in fissures and on ledges survived the first year (N=210).

### Discussion

*P. crassifolia* has some reproductive traits indicating a good adaptation to the vertical and overhanging rocks where it exclusively lives. The incipient pattern of floral zygomorphism of petals and coronal scales provides a small platform that could facilitate visits by pollinators that land on flowers, such as bees (see Stebbins 1970). On the other hand, the movements of the fruits and the myxospermy of the seeds are frequent features in

decreasing dispersal ability, which has been claimed advantegeous in habitats where seeds have few suitable places to germinate (Grubert 1974, Ellner and Shmida 1981). These latter traits suggest that founding of new populations must be quite infrequent, because seeds tend to stay near the mother plant.

In addition to the difficulty in placing seeds in favourable microhabitats for germination, most seedlings die during their first year of life, although the estimated 10% of survival during this time is probably a good result, since the year of monitoring was unusually dry. The main problem of *P. crassifolia* seems to be the scarcity of good microhabitats for the development of the species. Floral predation in some small populations also affects reproduction, reducing the number of flowers and therefore the production of seeds. All these characteristics affect the recruitment in established populations negatively.

In spite of its rarity, the reproductive success of *Petrocoptis crassifolia* in terms of fruiting can be regarded as normal since it is similar to or slighty higher than that which plants with hermaphrodite flowers usually have (Sutherland and Delph 1984). On the other hand, by assuming that *P. crassifolia* is a very old taxon (tertiary and subtropical, Montserrat 1979), it seems clear that its survival up to the present is already a success. The most important factor in its survival is probably its ecological specialization, which has allowed it to avoid competition from other, more widespread species which cannot survive in this habitat. In fact, the rock habitats in the Mediterranean region shelter numerous relict taxa, many of which are endemic (Davis 1951, Baudière and Cauwet-Marc 1986). Living in these habitats with weak interspecific competition, *P. crassifolia* shows some limitations associated with its reproduction and dispersal of seeds, problems that it reduces by means of the floral zygomorphism, the moving of fruits toward the substrate, and the myxospermy of the seeds.

At the moment, our results do not indicate that *P. crassifolia* is endangered: its reproduction is good and many of its inaccessible populations escape human and herbivore influences; moreover, most of them are fortunately protected in the Ordesa-Monte Perdido National Park. The limited distribution of the species seems to be fundamentally associated with a great habitat specificity, which usually leads to decreasing competitivity in other habitats (Drury 1974, Smith 1976), and with a lack of dispersal mechanisms for colonizing suitable but remote habitats (Stebbins 1980).

We would like to thank J. Buesa for providing the UV filter and J. Muñoz for pollinator identification. This research was supported by a grant (B CB-13/88) of the Diputación General de Aragón (Spain) and by the ICONA-CSIC project on "Cartografía de la vegetación del Parque Nacional de Ordesa y Monte Perdido". A part of this text was presented as a poster to the "III Colloque International de Botanique pyrénéo-cantabrique" (Biarritz, France, September 1992).

# References

Batschelet E. 1981. Circular statistics in Biology. Academic Press, London.

- Baudière A. and Cauwet-Marc A.-M. 1987. Les endémiques pyrénéennes, specialisation écologique et signification phytogéographique. In: Amigo J.J, Baudière A. and Muscat A. (eds.), Actes I Colloque International de Botanique Pyrénéenne. Groupement scientifique Isard, Toulouse, pp. 269–286.
- Cruden R. W., Herman S. M. and Peterson S. 1983. Patterns of nectar production and plant-pollinator coevolution. In: Bentley B and Elias T. S. (eds.), The biology of nectaries. Columbia University Press, New York, pp. 80-125.

- Dajoz I. 1991. Cycles de vie, variabilité morphologique et polymorphisme pollinique chez quelques endémiques pyrénéennes. Thèse, Université de Paris-Sud, Paris, 132 pp.
- Davies P. A. 1960. Pollination and seed production in Shortia galacifolia. Castanea 25: 89-96
- Davis P. H. 1951. Cliff vegetation in the Eastern Mediterranean. J. Ecol. 39: 63-93.
- Delay J. and Vivant J. 1978. Sur quelques endémiques pyrénéennes. Cytotaxonomie (1° partie). Bull. Soc. bot. Fr. 125: 485-492.
- Dendaletche C. 1970. Sur les *Petrocoptis* pyrénéens. Notes éthologiques. Bull. Soc. Hist. Nat. Toulouse 106: 306-311.
- Drury W. H. 1974. Rare species. Biol. Conserv. 6: 162-169.
- Ellner S. and Shmida A. 1981. Why are adaptations for long-range seed dispersal rare in desert plants? Oecologia 51: 133-144
- Favarger C. 1972. Endemism in the montane floras of Europe. In: Valentine D. H. (ed.), Taxonomy, Phytogeography and Evolution. Academic Press, London and New York, pp 191-204.
- Frohlich M. W. 1976. Appearance of vegetation in ultraviolet light: absorbing flowers, reflecting backgrounds. Science 194: 839-841.
- García M. B. and Antor R. J. 1992. Variabilidad fenotípica de un endemismo localizado en islas ecológicas: *Pinguicula longifolia* subsp. *longifolia* (Lentibulariaceae). Pirineos 139: 97-104.
- García M. B., Antor R. J. and Villar L. 1992. Son actinomorfas las flores de *Petrocoptis*? Influencia de la posición floral y los polinizadores en su simetría. Anales Jard. Bot. Madrid 50: 116–117.
- Gómez-Campo C. et al. 1987. Libro rojo de especies vegetales amenazadas de España peninsular e islas Baleares. Ministerio de Agricultura, Pesca y Alimentación, Madrid.
- Gómez-Campo C. and Malato-Beliz J. 1985. The Iberian Peninsula. In: Gómez-Campo C. (ed.), Plant conservation in the Mediterranean area. Dr W. Junk Publishers, Dordrecht, pp. 47-70
- Greig-Smith J. and Sagar G. R. 1981. Biological causes of local rarity in *Carlina vulgaris*. In: Synge H (eds.), The biological aspects of rare plant conservation. J. Wiley & Sons, Chichester, pp. 389-400.
- Grubert M. 1974. Studies on the distribution of myxospermy among seeds and fruits of angiospermae and its ecological importance. Acta Biol. Venez. 8: 315-551.
- Guitián J., Fernández J. M. and Rodríguez M. 1992. Flowering phenology and fruit-set in *Petrocop*tis grandiflora (Caryophyllaceae). Int. J. Pl. Sc. 153: 409-412.
- Herrera C. M. 1987. Distribución, ecología y conservación de *Atropa baetica* Willk. (Solanaceae) en la Sierra de Cazorla. Anales Jard. Bot. Madrid 43: 387-398.
- Herrera, C. M. 1989. Biología y ecología de Viola cazorlensis. II: Uso de sustratos, reprodución y consumo por los herbívoros. Anales Jard. Bot. Madrid 47: 125-138.
- Horovitz A. and Cohen Y. 1972. Ultraviolet reflectance characteristics in flowers of crucifers. Am. J. Bot. 59: 706-713.
- Jones C. E. and Buchmann S. L. 1974. Ultraviolet floral patterns as functional orientation cues in hymenopterous pollination systems. Anim. Behav. 22: 481-485.
- Kay Q. O. N. 1984. Variation, polymorphism and gene-flow within species. In: Heywood V. H. and Moore D. M. (eds.), Current concepts in plant taxonomy. Academic Press, London, pp. 181-199.
- Küpfer Ph. 1974. Recherches sur les liens de parenté entre la flore orophile des Alpes et celle des Pyrénées. Boissiera 23:1-322.
- Montserrat P. 1979. Biogéographie de la graine des Petrocoptis. Webbia 34: 523-527.
- Montserrat P. and Fernández Casas J. 1990. *Petrocoptis*. In: Castroviejo S. et al. (eds.), Flora iberica II; Real Jardín Botánico, Madrid, pp. 304-312.
- Montserrat P. and Villar L. 1972. El endemismo ibérico. Aspectos ecológicos y fitotopográficos. Bol. Soc. Broteriana 46: 503-527.
- Pettersson M. W. 1991. Pollination by a guild of fluctuating moth populations: option for unspecialization in *Silene vulgaris*. J. Ecol. 79: 591-604.
- Rabinowitz D. 1981. Seven forms of rarity. In: Singe H. (ed.), The biological aspects of rare plant conservation, New York, pp. 205-217.
- Ross M. N. 1936. Seed production of Shortia galacifolia. J. N. Y. Bot. Garden 37: 208-211.
- Saule M. 1991. La grande flore illustrée des Pyrenées. Ed. Milan, Toulouse, 765 pp.

Silberglied R. E. 1979. Communication in the ultraviolet. Ann. Rev. Ecol. Syst. 10: 373-398.

Smith R. L. 1976. Ecological genesis of endangered species: the philosophy of preservation. Ann. Rev. Ecol. Syst. 7: 33-55.

Stebbins G. L. 1970. Adaptative radiation of reproductive characteristics in Angiosperms, I: Pollination mechanisms. Ann. Rev. Ecol. Syst. 1: 307–327.

Stebbins G. L. 1980: Rarity of plant species: a synthetic viewpoint. Rhodora 82: 77-86.

Sutherland S. and Delph L. F. 1984. On the importance of male fitness: patterns of fruitset. Ecology 65: 1093–1104.

Villar L. 1988. El elemento endémico en la flora del Pirineo occidental español. Monograf. Inst. Piren. Ecol. 4: 371-381.

Villar L. and García B. 1989. Vers une banque de données des plantes vasculaires endémiques des Pyrenées. Acta biol. mont. 9: 261–274.

Vivant J. and Delay J. 1980. Sur quelques endémiques pyrénéennes. Cytotaxonomie (2e. partie). Bull. Soc. bot. Fr. 127: 493-505.

Westoby M., Jurado E. and Leishman M. 1992. Comparative evolutionary ecology of seed size. Trends Ecol. Evol. 7: 368-372.