

Zeitschrift: Boissiera : mémoires de botanique systématique
Herausgeber: Conservatoire et Jardin Botaniques de la Ville de Genève
Band: 8 (1949)

Artikel: The birth and spread of plants
Autor: Willis, J.C.
Register: Index General
DOI: <https://doi.org/10.5169/seals-895607>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 23.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Index General

Abundance 12-14; of community members (§) 413
Acanthacea Ch. IX, 195-221, 378, 423, 435; dispersal, numerical and geographical 198-9; leaders by size 206; new genera 32; pollen chars. 197, 207-8; taxonomy 205 seq.; in W. I. 203
Accident, introduction by 408
Accumulation at heads of lists 23-5
Aceraceae 307
Achariaceae 307
Achene and follicle 29, 368
Achillea (Comp.) 181
Acquirement of new chars. 211; from above by law 263; selection possible, but rare 299
Acrotrema (Dillen.) 145-8, 150, 303
Actinidiaceae 307
Action of law 299, 263; of vital factors 51, 54-5 &c.
Adaptability 9, 12, 32, 38, 43, 99, 387, 399, 400; born with sp. 12; needed as well as local adaptation 399
Adaptation 9-21, 26-9, 32-5, 37-45, 50-1, 60, 62-3, 85, 88, 95-105, 108-9, 114, 179, 184, 187, 190, 276-9, 314, (§) 317-22, 387, (§) 393-9, 405, 413-4, 417, 433-4, 467, 483; advantage, selection (§) 317, (§) 393, 299; and adaptability go together 12, 32; at birth 11-12, 32, 42, 190, 387, 394-5, 393-9, 405-6; automatic (law 9), 483, 418-21; becomes independent of structure 276; boom in, 316;

born with sp., see Adn. at birth; divergence rarely of adaptl. value 423; does not operate to separate by incompatible divergences 27-8, 362; gradual adn. inherent in nat. sel. 98; local 387; little evidence for necessary improvement 276, 398; must take up new position 16, 276; of coast plants in *Rubiaceae* &c. 434; of little influence 320, 179; of new spp. to new conditions 417 seq., 426, 433-4; of small fams. like *Juncaceae* 346-7, 426; out of court 26; progress under adaptn. on old theory (§) 393-7, 415; question simplified 417; structural adaptation worked too hard 414-5; supposed operation of 9-21, 27-9, &c.; to climate &c. gradual as plants travel 394, 418-21; to drier climate 415; takes a different place 276.
Adoxaceae 341
Advantage (§) 317, 398; for endc. genus 420
Aestivation imbr/conv in *Acanthacea* and others 195
Affinities 322
Africa 198, 200, 202-3, 268, 274-5, 347, 414-5, 421, 442-3
Agave (oideae) 348-9
Age and Area (§) 16-19, 22-6, 32-5, 38, 49, 64-9, 81-5, 96-7, 167-70, 185, 187-8, 226, 248-50, (§) 303, 315, 336, 360, 386-9, 404-5, 416, 448-9, 451-2, 458, 460, 481 (law 2); age ignored, but explains better 313, 318,

384-8, 397, 404-6; of char. is what really matters most 170, 219, 336; of supreme importance 321, 360, 386-8, 448; size and area also very important 193, 226, 386

Age and Size 19-25, 167, 303, 313, 332, 388, 481 (law 4)

Agreement, Ceylon/Madras 74, Spain/Balkans 69/74, *Cruciferae/Umbelliferae* 457, leading trop. fams. 466; Scillies/Shetlands 466.

Akaniaceae 341

Alangiaceae 307

Alismaceae (origin) 375

All As do not defeat all Bs 28, 89, 399; potentialities of fam. should be in leader 367

Allied groups of species represent more or less stable positions of equilibrium in cell division 97

Alpine plants 458, 460-3

Alston, A. H. G. 114, 151

Amarantaceae 75, 90, 92, 94, 174, 400, in Galapagos 475

Amaryllidaceae by size 348, 174

America 198, 200-1, 203, 233-4, 268, 273, 285, 288, 290-1, 347, 349, 353

Ammineae (Umb.) by size 186

Anacardiaceae 154, 173

Analysis of Ceylon flora 109-10; of Brit. communities 410-2

Ancestral forms should usually be simpler 103; fossil should be in leading tribe, if any 344-5

Andes flora 459, 463-4; watershed 455

Anemone(eae) 10, 19, 29-30, place 33-5, 49-50, 56, 101, 113-6, 174-5, 312, 321, 368, 387, 436; not superior to *Helleboreae* 321

Angiosperms, leading 45 fams. 329-30; 30 gen. 333-4

Animals 210, 322, 431

Annonaceae 135, 173, 295

Anodynes, verbal 12, 118

Anthemideae (Comp.) 176, 181, 438

Any char. may turn up almost anywhere, or be altered by mutation at any time 133, 142-5, 156, 175, 196-8, 210, 424

Any genus may carry a char. found elsewhere 102-3, 196-7

Any member of a fam. probably carries all or most of its potentialities 133-6

Appearance of duplicates in size in fam. lists 219-20; same char. in different places 196-7

Application of inductive methods 23

Approach to generic centres 50

Aquifoliaceae 307

Araceae Ch. XI, 248; by sizes 267-72; discontinuity 273, 275, 281; division by Pacific 273; endosperm 436-7; gen. 269 seq.; polyphyly 273 seq., 287; suggested origin 287; taxonomy 279-8; in Africa 274-6; in W. I. 466-7

Arber, Mrs. 255

Arctic-alpine grassland 26, 412

Arctotideae (Comp.) 442

Area occupied 35, and cf. AA; by endes. 65-7, 92, 117, 137, 139, 140, 198-202, 242-7, 386-7, 389, 398-9, 451; contour lines 57-8; of child usu. within or close to that of parent 399; of disp. 24

Argentine mountains 463

Arithmetical results and figures many, from 10-58, 81, 153-228 seq., 300-368, 422-78; basis for evolution 265; order begins to show in new places 259-66

Aroideae (Arac.) 271

Arrival in new place (rarely in order of age) 9, 11, 22, 37-8, 89-93, 192-3, 426, 437-57; of Brit. flora from France 447

Artificial system of classification 225, 227, 235

ASA laws of, 23, and rest of book; most important in dispersal 50, 85; override vital factors 45

Asclepiadaceae 173

Asia 198-202, and see Ceylon, India

Associations, see Communities

Assumptions of selection theory 109, 152, 418; *Evol.* 167

Atavism 103, 118

Auckland Is. (N. Z.) 17-19

Australian *Compositae* 443-7; *Inuleae* (Comp.) 443-5

Automatic isolation 310, (§) 316

Available total of chars seems increasing 313

Average diminution in size towards centre of gen. 50; dispersal Brit/France 50-4; of *Acanthaceae* 214-5; size of young genus much the same in different continents 200

B repeats behaviour of A 265, 313

Bahamas 471-2

Baillon, H. 212

Balkan flora 44, 46-7, 61, 69-80, 82-3, 86-7

Barriers 32-49, 66, 94, 387-8, 393, 397, 399, 400, 404-5, 431, 443, 447-8, 453-5, 458-61, 465 seq., and see *AA*, *Evol.*

Basellaceae 252

Basis of dispersion largely mechanical 139, 392

Beginning of genus &c. (and see these) 190, 297, 312-3, 405

Behaviour of endemics a miniature of that of "wides" 390; genera (§) 342-50, 29, 49-58; top genus in a family 243

Bentham and Hooker 131, 175, 218, 239-40, 322-3, 341

Berberidaceae 89, 432

Berthelot's experiment 373

Beta, dispersal 58

Betulaceae 84, 339

Big genera mainly composed of spp. of small dispersal 21

Bignoniaceae 314

Biology acquiring more arithmetical basis, and beginning to look like the result of a great thought translated into action 259-66, 392

Birth, adaptation at, see Adn.; arrival anywhere in order of birth improbable 425; birth of sp. 2 from sp. 1 191

Blue flower, production of 103

Boissier, E. 57, 59, 445

Bonnier, G. 49, 447

Boom in adaptation, relictum, &c. 316

Boraginaceae 87, 173

Bower, F. O. 370, 126, 150, 238

Bracts in *Cruciferae* 104-5

Branching of head into sub-heads (see Leaders) 250

Breaking off of islands 465; up structurally (cf. Taxonomy) tends to destroy regularity of curves 188, 91, 108, and geographical community 140, 159, 169 seq., 193-214, 246, 258; usually gives proof of former land connection 465

Britain, British 112s 23-5, 167-8; "ones" 20-1; dispersal goes largely with local size, and size in neighbouring countries 20-1, 166; ecology 7-9, 11-12, 26-8, 36, 46, 393-9, 404 seq.; families, esp. the large *Comp.* 175-88, 439, *Crucif.* 51, 185, *Gram.* 39-41, *Legum.* 172-3, *Ranunc.* 9-14, 28-38, 49-50, *Umbel.* 52, 172-3, 186; fams. not showing leader in Brit. usually from warm countries 193; flora reduced in outlying islands 48, 466; mostly from France 447-8; shows many sub-fams., tribes, and sub-gen. 108; gen. of one sp. in Brit. 21, of two spp. 312; gen. mostly larger than non-Brit. of same fam. 48; usually with wide diver-

gent in early spp. 164-5, 312; has all gen. 1-11 in Cichorieae, 1-6 in Anthemideae 438-9; leaders and sub-leaders 170-87, and see Leaders; local dispersal 83; mountain flora 461; outlying islands 48, 466; pliocene flora 36, 167; relationship to Spanish flora 453; southern origin 46-9, 82-7, 166, 185, 402, 418, 447-8, 453; "unsuccessful" fams. 11-12; varieties in Brit. 186; and see Communities, Ecology, Leaders, &c.
Bromeliaceae by size 348-9, 174
Brownlowieae (Tili.) 366
Buchenau, Fr. 347, 427
Buffon, G. L. L., 386
 Bulk of spp. in few largest gen. of fam. 307, 359
Burseraceae 135
Buxaceae 158-9

Calloideae 272
Calyceraceae 359-61
Carex 21, 26, 42, 312, 333, 400, 462-3, isl. 468, 475
 Carriage by water 48-9
Caryophyllaceae 10, 25, 42, 160
 Casual appearance of single chars. 133-5, 324
Cataclysms 219, 57
Cauliflory 355-6
Celastraceae 174
 Cell-division probably electrically controlled 130
Centaurea (Comp.) 21, 36, 44, 78, 80, 102, 176, 333, 445
 Centre of creation or evolution 63, 488; of generic area usually shows most spp. 243-4
Cercidiphyllaceae 254
 Ceylon flora 13, 17, 33 (curve *Dipterocarp.*), 67, 74, 89, 106-14, 121-6, 145-58, 201-2, 228-30, 234-5, 240-2, 244, 318, 356, 402, 406-10, 449, 455, 465-6; analysis of flora, 109-13; botanically two islands 465; Ceylon-Madras 74, 107, 402; chars. WE and

WEE gen. 111-3, 121-6; *Dilleniaceae* 145-51; *Dipterocarp.* 33, 153-6; endemics 106-8, 145-57, 390-2; *Hortonia* 202, 228, 230, 234-5, 240-2; Indian Ocean as barrier 244; introduced weeds 406-10; *Julostylis* 156-7; *Monoporandra* 155; mountains 459; *Pityranthe* 157; proportion of endemics increases with local size of gen. 157, 390; *Rhamnus* 107, 143, 353; *Ritigala Coleus* 318, 356; *Rubiaceae* by size 468-473; *Schumacheria* 145-8; *Trichadenia* 151-3; *Vernonia* 149; wet and dry regions (watershed) 401. Chalk grassland 26, 37, 410-4
Chanaz, J. P. St. A. de 374
 Chance, or happening 124, 154-5, 157, 284, 299-300, 324, 359-66, 373-80, 393, 397-9, 408-9, 418, 424, 470; enters largely into the matter 399; of change in important (old) chars. lessens as no. of chars. increases 450
Chandler, Miss 345
 Change in conceptions of classification 251; not necessarily adaptive 378; of climate acts as barrier 399-400; of composition of community (§) 414; of conditions 9, 66, 105, 185; of species sudden 276
Chaparral 420
 Characters. Most of the book is more or less concerned with this subject; some of the more important items in it are listed here; and see 500; and relationships 231 seq.; acquired by compulsion, destruction, or encouragement 302; any char. may turn up anywhere 156, 210; any member of a fam. probably carries potentialities of most 131-2, 150, 153, 155, 294; at A may also show far off at B, but not between 437; based

on divergence do not necessarily follow lines of genetic descent 264; "chance" largely shown in incidence 155, 293-4, 365-8; and *cf.* Chance; close relationship, with structural divergences 185, and see Leaders; combinations of chars. 105, 128, 236, 362, 367, 374-5, 381-2; complexes 211; contrasting chars. in small gen. 132; curves 391-2; discontinuity 90, 147, 154, 232-4, 239-41; due usually to sudden mutations 96, 105, 470; fuse in ancestry, however divergent 111; genetically acquired from above 263, 367 or handed down (dom. or recess.) 190-1, 211, 294, 367, 450; importance due to age 185, 217-20, 226, 236-8, 336, 425; incidence of chars. 143 seq., (§) 355, 359, 417; lack of transitions (§) 367-9; may be received without being shown by parent 114, 116, 122, 392 or appear in more than one place 436-7; more chars. should be used 237; no standard for 144; not casually acquired, due to inheritance from above, whether shown by parent or not 164, 263; not visible in parent 104, 106; now no longer certain that descend from something closely similar 164; of *Algae* &c. 384; of fams. perhaps a "chance" lot 299; of E1 or E2 may come from same parent 121-6; of new spp. 417; old chars. 450; perhaps less numerous than spp. 189, 375; probably all carried as a mixed lot of genes, present as potentialities in the leader 135; rank rises with number of forms showing the char. 252; seized upon for keys 380; that do not show in parent 122;

usually shown in perfect condition 60; value in one fam. or place no clue to value in another 189
Chatham Is. (New Zealand) 18, 158, 465-6
 Chemical analogies 372; chemistry and economic botany 510
 Chenevard P. 462
Chenopodiaceae 92, 174, 468
 Chief fams. of island endes. 469; ultimate factors in dispersal 449
 Child usually begins within or near range of parent 399
Chilean Compositae 440-2
Chrysosplenium 278
Cichorieae (1-11 in Brit.) 439
Cistaceae 456
 Classification 14, 104, 116, 129-30, 139-49, 170-94, 207, 214, 217, 222-3, 225-7, 241-7, 250-1, 255, 264, 273, 294, (§) 316-7, (§) 322, 341, 344; a more or less mechanical result of DDM 250-1; at present almost only by structure 175, 322; by doubling law 141; destruction of transitions largely obsolete 246-7, 250-3, 258; early gen. trace out lines of classification 222 &c.; for practical purposes 143-4, 217, 227; natural 142-3, 214, 225-7, 246-7, 250-1, 264, 316, (§) 322; a good one cannot yet be made 144; no standard for valuation of chars. 144, 170; often necessarily artificial 175, 225-7
Cleisostoma (Orch.) WEE 126
Clematis 10, 49-50, 56, 100, 113, 363
 Climate (§) 59-62, (§) 399, 277-8, 401, 454-60, and see *AA*, *Evol*, and Zones of Vegetation below; a barrier when changing 400-1; and weather 278
 Climax vegetation 408

Climbing plants 422
 Close structural relationship not a certain proof of genetic 143
 Closest possible genetic, least structural, relationship, towards head of fam. 108, 188
Cneoraceae 456
 Coalescence of chars must occur in going back 123
 Coast plants 434, 470
Cochlospermaceae 371
 Cockayne, L. 32
Coelachne (Gram.) WEE 113
Coelogyne (Orch.) WEE 125
 Cohabitation of *Podostemaceae* of different spp. 96
Coleus, 67, 318, 356
 Column monadelphous sta. 146
 Combination of chars. often apparently casual 99, 128-9, 147-8, 349-50, 360, 367, 374-6.
 Commencement of growth and dispersal of spp. 314, 434-5
 Common parent of divergence probably carried both chars. 93
 Communities 7, 8, 25-8, 37-9, 279, (§) 410-15; abundance of members 413-4; analysis of 410-12; arctic-alpine grassland 412; chalk grassland 26, 37, 410-11; change of composition of 414-5; distribution in 7-9, 413, dominant spp. in 411-3; formation and grouping 409-13; little taxonomic relation between members 7; *Molinietum coeruleae* 412; *Nardetum strictae* 412; of waste ground in Ceylon 409; pastures on highland ploughlands 412; *Quercetum roburis* 410-11; usually not very large 410
 Comparison Brit., France, Spain, Balkans 46-87; Brit.-Seychelles 448; Ceylon-Madras 74
 Compensation, principle of 263
 Competition 7, 8, 15, 19, 27-9, 32, 63, 80, 98, 114-5, 209, 244-5, (§) 253, 259, 276, 282, 301-6, 396; dependent upon whole outfit of competitor 396; lessens for spp., not for individuals 259; most severe among near allies 209; rarely specific 259, (§) 304-5; under natural selection 114-5
 Complexes (§) 211, 302, 362, (§) 369,
Compositae 10, 25, 42, 70-7, 84, 86, 101, 159-60, (§) 173-88, 263, 331, 354, 359-81, (§) 437-47, 456, 467; and *Umbelliferae* obey DDM and ASA well 187; dispersal of 437 seq.; early genera 176-88, 359; first 30 gen. have 9025 spp. other 1149 have 9004, 389; in Austr. 440-6; in Brit. 173-88, 439; in Medit. region 445; in New Zealand 439; leaders 175-8
 Composition of flora 109, of fam. 198-9, 228-30, 266-87, 302
 Compound interest (Yule) 298
 Confusion by splitting 214
Connaraceae by size 141-4, 363
 Continental floras Ch. II, p. 44
 Continuous leadership of original leader 425
 Contour lines and maps 58-9
 Contradiction of Darwinism 19, 87, 95, 116, &c.
 Contrasting chars. in WE and WEE Chs. V, VI, 95 seq.
 Control by ASA and DDM almost complete up to size of 15-20 spp. 388
 Cook's Strait, behaviour of W and E there 319
 Correlation of chars. 133 seq., 236-40
Corynocarpaceae 154, 158
 Course of Evolution 320
Coussareae (Rubi.) 468
 Crepis, endc. sp. in N. Zealand 438
 Cretan endemics 80
 Crombie 305
 Cross mutation 195, 205, 239
Cruciferae 50-1, 79, 81, 84, 104, 173, 365; bracts 365;

France and Brit. compared 50-1; parallelism with *Umbelliferae* 457
Cunoniaceae 342
Curcuma WEE 126
 Curve of genera lengthens both ends 33-4, 192
 Cushion plants 347
Cyclanthaceae for divergent variation, *Evol.* p. 76
Cymopterus (Umb.) 432
Cynareae (Comp.) 176, 179-80, 421, 445-6, 456
Cynocrambaceae 456
Cynomoriaceae 456

Darwin, C. 38, 151, 248, 396, 466; Sir Francis 96
 Darwinism and its contradiction 11, 14, 19, 21, 28-9, 34, 38, 59, 64, 69, 80, 87, 92, 95, 98, 115, 124, 168, 242, 252, 259, 304, 345, 364-5, 376
 Death of parent not necessary 259, 298, 395
 Decrease in emphasis of mutation 315
 Deductions, see Predictions; probable truth of their premises 319-20; how made 264-5
 Definition of a fam. 264-5
 Destruction of intermediates and transitions 19, 27-9, 32-9, 41-2, 66-70, 88-89, 108-9, 114-5, 118, 122, 143-7, 231-2, 251, (§) 253-9, 275-6, 286, 301, 304, 314, 339-40, 361-72, 396-7, 494; inherent in selection 98
 Development and dispersal of a flora and its parts, most pp. to 350, and after that esp. 357-65, (§) 367-72, 374-80, 382, 385-488, 491, 505; and see Laws, Spread, &c.; abroad and at home 17, 22, 168; as amenable to inductive reasoning as other branches of botany 45; begins to take form 99, and cf. Arithmetical; current theo-

ries helpless 45; discontinuity Ch. IV, p. 88; does not seriously depend upon adaptation; dynamic, not static (law 1) 481; geographically and numerically treated 198; goes with size of genus and world dispersal 20, 168; in Brit. 7-43, 167-8, and see Brit.; in Brit., France, Spain and Balkans 44 seq.; in Brit., France, &c goes with dispersal in world 54; in Brit. of genera of various sizes 20; in Balkans 44 seq.; in Ceylon 13, 109-12, and see Ceylon; in France 51, 447; in Spain 69 seq.; in islands 464 seq.; in mountains 458 seq.; in world 165; into most suitable situations largely the work of vital factors, partly overridden by ASA and DDM 45; largely dependent upon world and local size 25-6, 167-8; laws of, see Laws; made more discontinuous by splitting 108, 214; more or less mechanical 11-14, 17, 44-49, 184, 242-5, 258; (§) 430; of endemics a special case, a miniature of larger 64-9, 95; of dominants by vice-counties 410-3; of family 433; of fams. downward 453; of family names (Swiss) like that of plants 448; of flora (§) 447; of genus (§) 430; of members of a community 410-14; of small spp. or sub-spp. 21; of spp. and gen. (§) 242, 233, 430, 433; of spp. into vice-countries 9-10; of xerophily in dry climates 415 seq.; selective destruction no longer needed 117-8; study of problems in distribution 416, 424-9, 437-46

Diapensiaceae 66, 89

Dichotomous divergent mutation Ch. VIII, 164; 218, 220-1, 255, (§) 309, 314-5,

essential features (including survival of parent, accidentally omitted) 99-100

Dicots 82; Di/Mon differences 10, 24, 26, 42, 47, 52, 70-3, 80-3, 85, 168

Dictionary (sizes of gen.) 190-1 (reprint of 6th. ed. just out, 1948)

Differences between neighbouring floras mainly in small genera 449 seq.; betw. spp. 1 and 2 in 45 largest fams. 335-6; in floras at Ceylon watershed 455, 459, 465; in rate of travel 405; in uppermost tribes in size 187-8, 327, 447

Different combinations of chars. 367

Differentiation 186, 188-9, 192

Difficult to place a fam. in its order 332, 349, 382

Difficulties of nat. selection 42, 98

Dilleniaceae 96, 145-51, 192, 303

Diminution in average size towards centre of gen. 50; in flora upwards in mountains 459-60; in gaps towards foot of hollow curve 22

Dioscoreaceae 11, 288, 334

Dipterocarpaceae 33, by size 153-5, 331, 366

Direct mutation avoids long-standing difficulties 111

Discontinuous distribn. Ch. IV 88-94, real 89, structural 90; 108-9, 128, 140, 151, 250-3 (§) 253, 268-77, 281-4, 285-7; of char. incidence 116; of variation, one sp. to next 297

Discrepancy (structural) almost a law 353; of *Rhamnus*, and of *Siparuna* 353; betw. taxonomy and distr. 139-41, 144-5, 148-9, 152, 185, 193-4, 227-30, 242-7, 272-5, 279, 285, (§) 352-4; cf. 498-500

Dispersal see Development; of communities usually above the average 411; of *Compositae* 437 seq.

Distribution, see Development; basis of 385; problems, their treatment 443-7, 484-5

Divergence, most pp. to 375; Ch. VIII 164-94, 377, 449-50, 491; a great difficulty to Darwin 252; a principal mark of evolution 259, 339, 392; and evolution go together 249-50; any char. may be thus altered 424; automatic, probably electrical 425; as shown in Ceylon, WE gen. 111-13, WEE 121 seq.; as shown in Madras WE gen. 75 &c; between endemics in WEE gen. 125 seq.; between parent and child 170; chars. must come from higher up 123, seq.; divergent chars. 127 &c; each new char. may bring others in its train 258; early mutations tend to marked divergence 423-4; endemic often a descendant of wide close by 106; even one divergent child may form a new sp. 258; first feature to show in a fam. 263; formation of new sp. often insured 107; fusion of divergent chars. in ancestry 111; gives clue to wide taxon. separation of old groups 169; impressed at birth 251; increases upwards 209, 250-1, in fruit 368-9; in most small gen. 131; in mutation 105-7, 392; in taxonomy 449; necessarily affects more of important chars. as one goes back 383; no adaptational value as a rule 423; no longer due to selective destruction 246-7 of char. 131, 147, 332-6, (§) 355; of variation one of Darwin's difficulties 252; on whole diminishes as fam. grows 168, 209, 263; shows

early 392; subgeneric in *Rhamnus* &c. 107; taxonomic subdivisions dependent on it 451; usually neutral or indifferent 303; within the genus 181, 426-8

Divergent mutation, see Dicotomous

Dominants of communities 410-3; widely distributed 412-3

Doubt thrown on current opinions (Arch. Incompl.) 324

"Downward" evolution 120, 171-2, 188, 222, 250, 314, 344, 357, 364, 386, 396; much simpler 450-3, 396-7

Dropping out of smaller gen. in going north 402

Duplicates by size 178, (usu. scattered) 191, 219

Dying out 18, 34-65

Each genus in turn behaves like its predecessor 343; new one may require a new group for itself 158

Earliest descendants of early genera tend to pass automatically into other tribes 176, 423-4; mutations probably "larger" 310, 384; tend to head divisions 310

Early bird gets the worm 8, 13, 39 &c.; Brit. or other spp. most widely dispersed, and most divergent 167; comers get the best chances 8, 29, 39, 488; development of fam. 333; divergences tend to be larger 356; fams. widely scattered taxonomically 330; formation of sub-groups 39-40, 141-4, 170-2, 305-10, 322, 381; gen. probably largely direct descendants of sub-leaders 250; gen. tend to belong to different tribes 170-2, 330; gen. trace out chief taxon. lines of classn. 310, 322; giving off of sub-leaders by leader 173-86; growth and dispersal probably slow

431; mutations tend to head sub-leaders 171-3, 250, 284, 310, 424-5; stages of a fam. 219-21

East Asia, East N. Am., distribution to 89, 287

East Medit. region 445

Echinophoreae (Umb.) 457

Ecological changes need relatively short periods 96-7

Ecology 7, 8, 26, 28, 36, 41, 44-6, 54, 59-61, 279, (§) 415

Economic and chemical possibilities opened out 121, 372-4, and *Evol.* 8, 89, 177

Effects of age hitherto ignored 313; of climate 59; of migration 36-7

Efficiency 299

Eichhornia (introd.) 408

Electrical control of cell-division 130, 425

Element of artificiality in classification 108; chance 399

Emphasis of mutations 315

Endemism, endemics. A few chief headings are given here; and see §, p. 388, and law 12, p. 484; a direct contradiction of "Darwinism" 87, 116; a godsend to selectionists 63-4; a miniature of distribution in general 390-2; and invasion 393; a normal feature of a flora of reasonable size 87; areas 67, 137; as a special case of early stages in dispersal 67; as local adaptations 63-4, 95; as relics 63-4, 87, 95, 114-7, 446; as young beginners 55, 63-5, 67, 95, 313; at first a case of localisation and specialisation 393, 398; based on some mathematical formula 294, 392, 452-3; behave just like newly arrived spp. 390-1, 409-10, 486; by countries, see countries; by DDM (essential features p. 99 should include survival of parent, accidentally omitted) 310,

&c; *Evol.* 110; by fams. see fams.; by law 67-8, 76, 87; by size of genus 390; by sudden mutation, not selection 68, 453; chars. may be received without being shown by parent 115, 392; definition 64, 188; determined by law 67-8, 87; first formation 106, 190; genera 85, 140-1, 145, *AA Ch.* XVI, p. 169, espec. 175-83; usually small 139 and *Ch.* VII, 137-63; in large fams. 140; in Ceylon 142-57; of other countries, see countries; illustrate early stages of dispersal 390-1, 484 (law 12); in broken and/or mountainous country common 63, 140; in *Cynareae* 445; in *Inuleae* (Australia) 443; isolation 140, (§) 316; lack of definition 55; limits of endemism 67; mainly in large and "successful" families 69; mark invasions 140, 202, 393; miniature of dispersal in general 139, 390-2; new spp. tend to evolve to suit new conditions better 189; no immediate adaptational structural basis 109; no line can be drawn between E and W 64, 139-40; northern barriers in old world 66; of islands 464; of mountains 458; of one country shows definite relation to others close by 69-80; perfection of chars. 335, *Evol.* 114; permutations and combinations of chars. 362; phenomena parallel to those of non-endemics 139; proportion increases with local size of genus 390-1; relationship between endm. of S. Eur. and composition of Brit. flora 82-3; selection cannot control 14-100; swamping 478; small genera usually endemic 129-30; unsuccessful genera 129; usually descendants of

W or E close by 106 seq.; WE and WEE &c. 106 seq.; young beginners 67, 95, 319; youth of endemics shown at Cook's Strait 319
Engler, A. 266, 336, 339
Engler-Diels 238, 326, 337, 483
Epacridaceae 342, 368
Epigaea, disc. 89
Equilibrium of nuclear formation 362, 372
Erica vagans 412
Ericaceae by size 368-9, 342, 364; *Eric.* and *Epacr.* 342
Erythroxylaceae 195
Essential features of evolution by DDM 99 (survival of parent accidentally omitted) 276, 310; of taxonomic difference is divergence 449
Euphorbiaceae by size 338
European flora, Chs. I, II, III
Even one divergent child may form a new sp. or gen. 258
Every new gen. may need a new group for itself 158
Evidence of fossils, see Fossils; for dying out, once a reasonable area covered, very small 35; of inferiority in dispersal due to structural divergences little 11, 23
Evolution is largely dealt with throughout, but a few features require special marking; a necessary consequence of life; and distribution go together 8; any member of fam. probably carries most or all of its potentialities 133-4, 209-10; appearance of some chars. not a cause of evolution, but a by-product of it 131; as set forth here is a reversal of "Darwinism" 168; automatic, probably electrical 425; back to front 328, 331, 344, 355, 416; based on some mathematical conception 123-4, 215, 259, 294, 300, 318, 322, 328, 391, 416, 452-3; before a fam. is of any size,

its chief taxonomic divisions are marked out 250, 322 and see Leaders; biology begins to develop other arithmetical bases 259; and seems to be the result of a great thought or principle which is working itself out by law 493, see Super-Mendelism; by law 67-8, 87, 124, 300 by sudden mutation 453; char. of more value in a fam., the number that show it (i.e. the older it is) 128, 217; classification (taxon.) a more or less incidental result of DDM, 246; close genetic relationship may exist with wide structural divergence 188; complexes 211; cf. *Evol.* p. 139, and p. 128 above; *Compositae* in the light of our work 172-81, 437-47; DDM and ASA well obeyed in *Comp.*, *Umb.* &c.; destruction of connecting links 41-2, 99, 108, 146, 158, 227, 286; dispersal by ASA and DDM 452; divergence a mark of 210, 259; divergent mutation 113, *Evol.* 76 and Test case IX, p. 110; dominant plan of evolution 452; downwards 14, 16, 99, 103, 168, 173, 175, 184, 188, 222, 233, 238, 242, 251, 297-8, 315, 331, 344, 348, 392, 450-3, 482 (law 5), 489, 495, 497 (evidence almost overwhelming 168-9); early gen. tend to go into other tribes 246; essential features of evolution (incl. survival of parent accidentally omitted) 99, 310; extension of the laws of Mendelism now required 263; genera artificial divisions 210; genetics and taxonomy often at variance 142; geogr. distrn. still largely based on selection 361 (and cf. Brit. flora, esp. 427-8); geogr. localisa-

tion of struct. features goes towards species 14, 147, 331, and see downwards; handing down of chars. 187-94; has hitherto been taken backwards 331; hollow curves see h. c.; incidence of char. needs much study 141 seq., 207, 286; incomplete evolution (teratology) 100-04; independent of nat. selection 14, 16-7, 40, 60-8, 76, 81, 93-5; inheritance of chars. that are not shown in immediate ancestors 206-7; intermediates 16, 169; isolation Ch. VII, p. 140; kaleidoscopic mutation (§) 376, 294, 300-1; laws of 97; little to do with adaptation 99, 109; mainly controlled by DDM and ASA 452; mechanism of 242; must apparently go on 302; mutations at top on whole larger 171; new spp. to suit new conditions 189; no immediate adaptational structural basis 109, 392; no longer a matter of chance, but of law 23, 124, 171-2; no standard for chars. 144; now coming into line with physical sciences 293; of a fam. as seen from above downwards 346-50; of spp. downwards 242, and cf. gen. and fam. under Leaders; Pandora's box 94, 133-4, 147-8; perfection of chars. T. C. X, *Evol.* 114, 45, 127-8; permutations and combinations of chars. 147-8, 152, 189-90, 362; polyphyly 150-2, 291, 293; proofs of downward theory cf. Islands, Leaders, Cynareae, Inuleae; rank of mutations 207 (of gen. the higher the larger 226); relictism 253-9; seems the result of a great thought or principle, which is working itself out by law 493 (no. 34); selection can hardly control

DDM 14-100; picks out individuals 258; (unsatisfactory as explanation 300); simplified 293, 297, 450-2; size and space 19, 201; small gen. usually endemic, or behave as such 129, 131 (usually satellitic, TC XVIII *Evol.*, p. 128); super-Mendelism 103, 263; swamping (endcs. only in non-endc. gen) 202, 478; test-cases, see *Evol.*; transitions to other fams or gen. 169, 314; under steady influence in one direction 276-9; relative sizes of gen. TC III, *Evol.*, p. 95; variety of char. in uniform conditions TC XXIX, *Evol.* 156; young beginners 64, and see Endcs.; Yule and Willis, see Y and W. Explanation by selection very unsatisfactory 300

Extension of laws of Mendelism wanted 263, 294

External conditions probably much to do with evol. of char. 104-5, 210

Extinction 69, 92-3, 167

Fagaceae 339

Failures 19

Family(ies); a more or less connected combination of chars. 349; any member probably carries most or all chars. in some genic admixture 131-6; becomes genetically confused by structural subdivision 108, 214; begins as one specialised and localised endemic 190; development and dispersal 432-5; of *Compositae* 437 seq.; early produces sub-leaders, cf. Leaders; first 45 by size 329-35 (to 50 gen. 173-5); first and second gen., relative size 191, 335; follows rule of early divergence *Evol.* 199; grouping artificial 142; happens to show in common a

group of well-marked chars. 362; in Brit. 192; reaching 112 there 84-5; in order of size 173-5; and see Leaders; local size usually depends upon local age 390; more or less concentrated about place of origin 245; more tribes, and proportionately more endemics, in large fams. 140-1, 252; names in Switzd., distribution of 448 (*Evol.* 35-8); not showing leader in Brit., usually of warm countries 193; or genus soon shows taxonomic divisions 141, 176, 182-3, &c.; or tribes, with Medit. centres 456-7; pairs of 342; potentialities carried by first genus 241; relative size of first and second genus of 191, 335; small 341; subdivisions, see Leaders; to size 15-20, 388; with headquarters in Medit. region 456

Few actual generations from leader to foot, even in large fam. 356

First 45 fams. by size 329-335; 30 genera by size 333-4; 30 genera of *Compositae* by size have more spp. than the other 1149, 359; come, first served 13, 412, 488 &c.; dozen or so of gen. trace out taxon. division 322; genus, in 5 subfams. of *Araceae*, divided from second by Pacific Ocean 273; genus of fam. begins as specialised, localised, endc. 190; genus, on average, double the second 191, 335, 359, *Evol.* 101; no good evidence that early mutations really larger 251; tribes often smaller than later in very old fam. 327

Fischer, C. E. C., 74-5

Fleeming Jenkin 304, 318

Flacourtiaceae 151-3

Flora, development and dispersal (§) 447-54; of any one

country tends to show great divergences 172

Floral regions (§) 454-7; imply char. fams. 455-7

Floras, Alps 461, Andes 463, Bahamas 471-2, Balkans 69 seq., Brit. 7-61 (adjacent islands 464-78, Brit. Fr. Sp. Balkans 46 seq.), Ceylon 107 seq., Crete 82, France 44 seq., Galapagos 472-3, Italy 74, Malay Penin. 127, Medit. Is. 80 seq., Medit. region 418, 447-54, Pliocene of Brit. 36-7, Seychelles islands 464 seq., of mountains 458 seq.; Inuleae in Australia 444, Cynareae in world 445

Fluctuating differences 379

Formation of community 410; of duplicates in size as lines of descent separate 219-20; of flora 406, 447-54; of new sp. involves divergence 105-7, 431

Fossils 35-8, 57, 99, 169, 219, 254-7, 343-6 (§) 479, 494; may be sidelines of evolution 219

France, flora of 44-87, 185, 447; invasion of Brit. 185, 447-8; *Crucif.* in 51 seq., *Ranunc.* in 53-5, *Umbell.* in 52

Fries, R. E. 463

Fruit chars. 368-9; divergence in *Eric.* 369; *Calycerac.* 359-61

Functional adaptation 61-3

Fungi 302

Further back in time, mutations tend to be larger 166-9; proofs for downward evolution 188

Fusion of divergent chars. in ancestry 111

Gaertnera (Rubi.) 189-90, 236-7, 361

Galapagos Is. 75, 472-3, 475

Galium (Rubi.) 80, (leaders) 182, size and place 468

Galopina (Rubi.) 132-3

Galton's Law 226, 232, 239, 294, 417

Gaps between larger gen. in fam. in size 22, 34; *Senecio* and followers in *Senecioneae* 177, 423-4; times of arrival of spp. 22

Garcinia (Guttif.) WE 114

Gardenieae ahead of Psycho-trieae, Madagascar &c. 471

Gardiner, J. S. 49

Genera Plantarum 135

General development and spread (§) 403, and see Developmt; distribution of plants Ch. XIII, p. 385; review in *Araceae* 248

Generic (see genus) and fam. differences 98, 185; leaders and sub-leaders 181-4, 427-8

Genetics 96-7, 108, 133-4, 142, 175, 188, 207-8, 234, 263-6, 285-6, 293-4, 300, 312, 317, 416; allies need not be structurally close in all chars. 93, 322; and taxonomy at times at variance 139, 273; groups 236-7; grouping does not lend itself well to classification 139-40, 236-8; lines cross taxonomic in places 108; relationship tends to be closest at top of fam. 188

Genic changes 97, 300

Gentianaceae 218, 300, 422

Genus, genera, generic, &c.; an artificial division 214; and fam. differences more difficult to explain by selection 95-9; and spp. do not usually arrive in order of birth, but note *Compositae* and esp. *Cichorieae* in Brit. and Islands 178; beginning of 190, 277, 312-3, 431-3; behaviour of 51-3, 342-50; chars. produced by a kind of shuffling 189-90; closest relationship at top of a fam. 147 seq., 188; development (§) 430; distribution in Brit. 20-27; endemic Ch. VII, p.

137; in S. Eur. 63-90, and *cf.* countries; each new one liable to need a new group for itself 158; first gen. of fam. begins as localised specialised endc. 190; follow same rules as spp. 137; great proportions on mountains and islands 140, 201, groups of 235-7; have most spp. local 69, 184; in a flora chosen by reactions to laws and conditions 187; in both wet and dry zones of a country 401, and see Ceylon; larger Mediterranean mostly in Brit. 83 &c.; larger than 15-20 spp. not so closely governed by ASA and DDM 140, 221, 388; largest in Brit. have most 112s 25, 42, 84-5; largest 30 genera in world 333-4; leaders and sub-leaders 141, 173-94, and see Leaders; nearer centre of fam. smaller in aver. size 49-50; not known in Brit. soon appear on continent 44; numbers of small gen. of small area very large 44 seq. 127; of Ceylon, endemic 145; of other countries see countries; of dispersal 112 on whole the largest in Brit. 25, 42, 84-5; of *Ranunculaceae* in Brit. by size 49-50; of size 15-20 come within range of great climatic and other changes 140, 388; outlying the largest 27; produced by a kind of kaleidoscopic shuffling 189-93; rank higher the larger in the fam. 222; relationship not to be found with certainty by taxonomic methods only 279-80; selection little to do with chars. 189; sizes, see p. 350, and Leaders and other lists; small, discontinuity in 109; usually will not arrive in exact order of birth 418; with one or more

spp. in one vice-county only
21

Geographical distribution; see Distribution, Development; as open as rest of botany to inductive reasoning 43; mainly determined by ASA and DDM, at least in earlier stages to size of 15-20 spp. 449

Geography 8, 88-99, 117, 130-2, 137-63, 198-204, 222-38, 242-7, 266-79, 352, 385-495; and taxonomy do not always agree 90-9, 143, 148-50, 209, 227, 230, 274-5, 285-6; bringing order into distribution (with DDM) 193; cannot be neglected 143; classification (Schellenberg) 142; continuity often lacking 89-94, 229-31; destruction need not be invoked so much as hitherto 118; discontinuity 266, 276; of fams. very important 198, 202, 230; relationships as important as structural 198, 205; separation 139-44, 129; taxonomic discontinuity 273

Gesneraceae 45, 75, 82-3, 173

Globulariaceae 82-3, 456

Gorgonidium (Arac.) disc. 272

Glands at base of stamen 240

Gradual acquirement of adaptation under selection 60; adaptation of fam. to new conditions 426; evolution 276, 300; transition inherent in nat. selection 98

Gramineae 39-41, 86, 92-3, 173; in Brit. 39-42

Graphic curves *Ranun.* *Diptero.* 31-3; 15 leading fams. 260

Gregory, Mrs. 428

Great concentration of early fams. and gen. in Seychelles 448; divergence shown by early fams. 322-4

Grisebach on age 386

Group of allied spp. represents so many more or less stable positions of equilibrium in cell division 97

Growing through 126, and see Bower

Growth by mutational division 331; of fam. 218, of gen. 241-4, of spp. 243; of few endemics into flora 444-5

Grumilea (Rubi.) complex 212

Guettarda (Rubi.) 470

Guppy, H. B. 48, 303, 313, 315, 328, 331, 448, 467, 481-2

Gynura 122, 125

Gypsophila (Caryo.) 44

Gyrostemoneae (Phytolacc.) unisex. fls. 134

Hairiness 149

Handing down of chars. 191-2

Handling problems of distribution 176-194, 424-5; examples Inuleae of Australia, Cynareae, 443-54, &c.

Happening, see Chance, and Prediction

Harland S. 300, 317, 393, 479

Harmonisation, geogr. and taxon. 143-4, 205, 207-8, 231-8, 272, 275-6, 281, 339-40; between taxon. and genetics 205, 210

Hawaiian flora and endemics 24-5, 38, 67, 75, 116, 145, 159-63, 415-6, 467, 473, 477

Hayward, W. R. 186

Heads of all divisions usually early formed 187; of fams. carry all potentialities 170-2; of tribes chiefly in large gen. 140-1, 312; usually first to reach another country and get largest dispersal there 312

Heights reached by trees in Brit. 460

Helenieae (Comp.) 441-2, 447

Heliantheae (Comp.) 159-60, 181

Helichrysum (Comp.) 440-4

Helleboreae (Ran.) 10, 19, 29, 31, 33-5, 56-7, 321, 387

Herbarium 227

Hibiscus (Malv.) 49, 156, 244, 292, 429-30

Hieracium (Comp.) 21-3, 47, 78, 101-2, 107, 116, 176-9, 182-4, 263, 333; and Rubus 183, 314

Highland pasture land 412

Hillebrand 159

Himalaya 463-4

History of plants largely one of invasion 402

Hitchcock 309, 311

Hollow curves 16-23, 29-34, 51-2, 65, 67, 91-2, 170, 192, 218, 220, 258-62, 298-309, 388, 390-431, 452, 504; and see *Evol*; automatically produced 219-20; curves of 15 largest fams. 260, mixed curves 261, log. curves 262; lengthens at both ends 33; necessary result of DDM with survival of parent 305; poor in *Monimiaceae* 234

Holmes, O. Wendell, 209

Hooker, J. D. 42, 124, 140, 298, 328, 383, 467-8

Hortonia (Monim.) 228-35, 241-2

How long does first genus continue to lead? 192-3, 425

Hurst, C. C. 105, 108

Hutchinson, J. 12, 37, 239, 337-9

Huxley, T. H. 12, 257

Hydrocharitaceae by size 349, 357, 426

Hypothesis, working 49, 96, 99, 192, 388

Icacinaceae 174

Imbricatae (Acan.) 200; imbr./conv. aestivation 195

Importance of chars. 217, 225-6, 236-7, 250, 450, &c.; goes with age 236-7, 250, 336, 368

Incidence of chars. (needs much study) 141 seq., 417, &c.

Incidence of mutations probably obeys some law 226-7

Incidental showing of bracts in *Cruciferae* 105

Incompletae 238, 323-4, (§) 336-9, *Evol.* 104; incomplete evoln. 104

Incongruity, geogr. distr. and taxon. 88-9, 91, 228, 232, 234-7, 241, 269-72, 275, 279-285, 323-4, 330, (§) 336-9, 344-6, (§) 352-4

Increase in number of spp. with approach to generic centre 44; in size of continental flora in going south mainly due to increase in Brit. genera 47-8; in size of tribes downwards in large fams. 327; of divergence upwards 209; of hollow curves at both ends 192

Index Kewensis 32, 117, 190, 307

India 113-4; and see Madras, Ceylon, Indian Ocean

Individuals, not species, chosen by nat. selection 99

Induction and inductive methods 23, 76, 193, 231, 248-9, 344-5, 386-8

Infiltration of plants 403

Influence of outside conditions 276-7

Inheritance, laws of 190; like Mendelism 127; of acquired chars. 416; often does not follow genetic lines 436-7; probably obeys some law like Galton's 232; probably carried as some assortment of genes 133

Injurious insects 28

Intermediates 168-9, 258, 292, 298, 336-9

Introduced weeds 406-9

Introductions do not owe their success to any adaptation for the purpose 408

Inuleae (Comp.) by size 181; of Australia 440, 443-5, 486

Invasion (§) 401-2, of New Zealand 403, (maps) 407-8, 465-6, 477-8; effect of barriers 447-8; of Brit. from south 68 (§) 401-3

Investigation of problem in distribution facilitated by laws of ASA and DDM 176-194, 424-5, 443-54

Iridaceae (leaders) 174

Irregularity in small fams. 307

Islands 17, 80-3, 161, 138-40, 201, 204, (§) 464-78

Isodendrion (Viol.) e.g. 160-1

Isolation 140, (§) 316, 415-6, 446, 460 seq., 493

Italy flora 74

Janchen 172

Jenkin Fl. 304, 312, 318, 396

Juan Fernandez 468, 475

Julostylis (Malv.) e.g. 156

Juncaceae by size 346-8, 426-7, 434

Juncus sub-leaders 346-8, 426-7

Justicia (Acan.) Ch. IX, p. 195, complex 211; and *Ruellia* 423-4

Kaleidoscopic mutation 32-4, 104-5, 128, 282, 300, 362, (§) 374-80, 430, 449-50, (law 6) 483

Kerner A. 48

Keys of monographs 245-6

Killing out 28, 38, 66, 108, 435

Komarov 404

Labiatae 46, 293

Lack of transitions (chars.) (§) 367, and see Chars

Lamarck J. B., 360, 386

Land connection 48-9, 202, 487

Lantana (Verb.) 406, 410

Large fams. or gen. 10, 25, 42, 48-52, 69, 75, 80, 81, 140, 188-90, 231-9, 320, 327-8, 314-5, 335, 339; carry all or most potentialities in fam. 211; contain bulk of endemics 75, 113-4, 137-9, 159, 389-90; groups usually headed by large gen. 32-3, 140-1; cf. Leaders; leader usually followed by many followers 425-6; pass barriers better 47-9

Larger a fam., the greater proportion of exceptions and complexes 311; the larger the area occupied 389; the more tribes in most cases

357-8 &c.; a genus in Medit. region, the more likely to occur in Brit. and to have many southern endemics 83; a genus, the greater its range of chars. 364; a genus, the greater the proportion of 112s 25; a genus, the more likely to reach a given country 48; a genus, the more taxonomically important 232; a genus, the older on the whole 328, 25; fams. and gen. show greater range of chars. 358, 362-4; fams. of Angiosp. and location by different authors 329-30; fams. usually headed by larger gen.; gen. need most subdivisions on whole 151; groups show more "relics" 314; structural differences usually inherited from far back, 170, 172; the fam. or gen. the more important in later history 231, 359; the number of possible parents in a fam. the more small gen. 52, *Evol* p. 91

Largest 45 fams. 329-30; 30 gen. 333-4; fams. and gen. on whole the oldest 327-8; and the most widely dispersed 25 seq.; fams. show good hollow curves 306, fig. 260; genera at top of fam. most closely related 207, 318; contain most of the endemics ("relics") 75, 137, 159, 320; genera go furthest out in dispersal 27, 48-9; genus in fam. usually about twice the second 329-30, 335; species vary most a consequence of DDM 345

Lauraceae 115, 232, 234

Laurelieae (Monim.) 229-30, 234 5; closely obeyed by small fams. and gen. 140

Laws of ASA (§) 22-3, 45, 62, 84, 185, 187, 205, 257, 297,

303, 315, 321, 388, 413; much more to do with dispersal than has adaptation 49, 50, 140

Laws of Distribution 8, 13, 17, 27, 64, 480-88; (1) dispersal dynamic, (2) age and area, (3) size and space, (4) age and size, p. 481; (5) evolution downwards 482; (6) kaleidoscopic mutation 483; (7) survival of parent, (8) dichotomous divergent mutation, (9) adaptation automatic, 483; (10) rate of progress greater with age 484; (11) leaders and sub-leaders 484; (12) endemism the earlier stages 484; (13) divergence the more marked the nearer to leader 484; (14) influence of barriers 297, 307, 321, 485; (15) to him that hath 495

Laying-out of a problem in distribution should be done primarily by the laws of ASA and DDM 424-5, examples 444-6

Leaders and sub-leaders 30-1, 141, 153, 156, 176-86, 206, 230, 267, 269-72, 301, 325-6, 338, 346-9, 368, 422-9, 437, 444-6, 496; do not always lead 191-3, 425; early descendants tend to go into other groups 423-4; of all geographically possible groups usually appear in any country 423; overtaking by younger genera 429-30

Leaders of orders 428-9

Leaders families: *Acanth.* 206; *Amaryll.* 348; *Araceae* 267; *Bromel.* 348; *Compositae* 176; *Connar.* 141; *Dipterocarp.* 153-4; *Ericaceae* 368; *Euphorb.* 338; *Hydrochar.* 349; *Juncac.* 346; *Legum.* 325; *Malvaceae* 156; *Monim.* 230; *Orchid.* 326; *Palmac.* 474; *Penaeac.* 301; *Ranunc.* 30, 31; *Rubiaceae* 468;

Rubi. in islands 469; *Triurid.* 437; *Umbellif.* 186

Leaders sub-families, or tribes : Ammineae (Umb.) 186; Anthemideae (Comp.) 181; Aroideae (Arac.) 271; Astereae (Comp.) 180; Cichorieae (Comp.) 178; Colocasioideae (Arac.) 270; Cynareae (Comp.) 179; and in Medit, and world 445-6; Inuleae (Comp.) 181; and in Australia 444; Lasioideae (Arac.) 270; Mendonioideae (Acan.) 216; Monstroideae (Arac.) 271; Nelsonioideae (Acan.) 216; Pholidoroideae (Arac.) 269; Pothoideae (Arac.) 269; Seneioneae (Comp.) 177; Thunbergioideae (Acan.) 216

Leaders of genera and subgenera : Desmodium (Legum.) 428; Galium (Rubi.) 182; Hieracium (Comp.) 182; Juncus (Junc.) 347-8, 426-7; Lamium (Labi.) 427; Luzula (Junc.) 348, 427; Rhamnus (Rham.) 107; Rubus (Ros.) 183; Saxifraga (Sax.) 428; Scirpus (Cyper.) 427; Tiliandsia (Bromel.) 349; Viola (Viol.) 428

Leadership of first genus continuous or not 191

Leaf chars. of small gen. 131-2

Lecythidaceae 146

Leguminosae 135, 172, 325-6, 332, 334-5, 418-9

Lemnaceae 10

Lepidaceae (Crucif.) 92

Less marked features of *Monimiaceae* 239

Liabinae (Comp.) 443

Like causes, like results on similar material 277-9, 289, 301, 424-5

Liliaceae, chars. 135

Limits of area 243-4, of distr. of endemics 66

Lindau, G. 197-9, 205-9

Linnean system 155, 227, 264

List of tribes of leading genera of fams. 173

Little endemism north of Alps 44; variation in rate of spread between allied spp. 169

Local adaptation 63-4, 95-7; species chiefly in larger genera 45-7, 63-4, 69, 184-5

Localisation and specialisation 393, 398

Locally larger gen. overlap smaller above and below 21

Logarithm curve 18-19, 211, 218, 431, diagram 262, 306, 319

London Catalogue 9, 29

Long tail with small head 441, 471

Loss of improvement by crossing 396

Luxemburgia (Ochn.) 146

Luzula (Junc.) 348, subgen. 427

Lysimachia (Prim.) 28

Madagascar 139, 230-1, 234, 471, 476

Madras 74-5, 107, 113, 143; and Ceylon 74-5, 402

Magnoliaceae 327-8, 337-9

Malaya 106, 116, 127, 273, 283, 476

Maldives 49

Malopeae (Malv.) 157

Malvaceae by size 156, 292, 429-30

Many difficulties cleared up by taking evolution downwards 297; tribes in Brit. 172-5

Maqui 420

Marantaceae 371-2

Mathematical theory of evolution 300

Maxims, taxonomic 351

Mechanical distribution 17, 22

Mediterranean region 55, 63, 68, 76, 79, 82-5, 455-7, 418-20; a centre of creation or evolution 63

Members of a flora determined by reactions to laws, especially mechanical 187

Mendelism 122, 184, 190, 268, 291-3, 311, 417

Menispermaceae map of distribution 137-8

Mesophytes 278-9

Methods of study of distribution problems 424-5

Migration 15, 36, 66 (§) 401-3

Mikania, intr. weed 408

Mivart, G. S. 331

Mixture of crops 409

Molinietum coeruleae 412

Monadelphous stamens 146

Monimiaceae Ch. X, p. 222; by size 230, lack of geographical connection 229-30; taxonomically grouped 228, geograph. 229, of Madagascar 234; probably not a simple genetic fam. 238

Monocots 26, 39-42, 47, 85, 123-4, 157, 266, 375; reaching 112 in Brit. 42; Alps 462

Monographs, keys to 245-6

Monoporandra (Dipt.) 153, 155, 432

Monotypes 350; and *cf.* 158

More taxonomic division, more geographical 108-9, 149-51; we break up a fam. the more artificial and discontinuous the divisions look 108, 193, 198

Moringaceae 456

Most small gen. closely obey ASA 256; most widely dispersed in Brit. show most southern endemics 83

Mountains (§) 458-64

Mutation, throughout the book, espec. 14, 34-5, 44-5, 58, 96-100, 103, 105-7, 111-14, 128-9, 150-1, 166, 187-94, 223, 228, 246, 252, 277, (§) 309, 324, 352, 369-70, (§) 376; cross 202; divergence 163, 298; earlier tend to be larger 171; first usually takes offspring into another tribe 186, 190; incidentally involves structural change 388; of specific chars. (Hurst) 105; probably elec- trically controlled 162; producing gen. probably employs more chars. than sp. 186; rarely changes one char. only 162; results more perfect than by selection 103; single mutn. can cover any difference parent/child; small in series 15, 126; usually adaptationally indifferent 100

Mutisieae (Comp.) 441

Myosotidium (Borag.) 156

Myosurus (Ran.) 15, 101

Myrsinaceae/Primulaceae 342

Nardetum strictae 412

Nasturtium terat. 102

Natural classification 142-5, 197, 214-5, 217-8, 225-8, 242, (§) 322; at present impossible 143, 217, 226-7

Natural selection, see Selection; system 217

Nearer centre of fam., the more small gen. 50, 59

Nelsonioideae (Acan.) 195, 216

New chars. acquired by law, not casually 263

Newcomer usually suited to very few communities 413-4

New directives in the study of distribution Ch. VIII, p. 164

New fam. or gen. 305-16; genera in France 49-50; genera usually satellite 32; probably of very few individuals 490

New spp. adapted to new conditions 190, 391; formation involves divergence 106-9; not produced by trial and error 191-2; production of 398; usually within or close to area of parent 201

New Zealand flora 17-18, 58-9, 65, 158, 204, 304, 318-9; effects of Cook's Strait 319; endemics 158; maps of invasions 407-8; invasions 403, 407-8, 477-8; *Comp.* 439-40; *Rubi* 473; zoning 319

No char. can be used alone

with confidence 107; no evidence for serious dying out 35-6; no line possible between endemic and wide 64, 139; no longer needful to find intermediates, living or fossil 298; no necessary destruction of transitions 118; no serious difference between allied spp. in rate of spread 169; no standard for chars. 144, 170; no use value in divergences 301

Non-Brit. genera 44-7

Nothing to show superiority of one sp. over another 304

Nuclear reactions and readjustment 96-7, 388

Numbers of sub-divisions of fams. and gen. in Brit. 11, and see Leaders; of small genera very great 129

Numerical differences, upper and lower tribes of *Comp.* 187-8

Nymphaeaceae 434

Ochlandra (Gram., sta. difft) 365

Often, frequently, usually, &c. continually used in taxonomic descriptions 128, 355, 363, 424, 483

Oldenlandieae (Rubi.) 468-9, 471-3

Old position now becomes untenable 233

Older a sp. the more rapid its spread 398, 404-5, *AA* 34; spp. find country more open to colonisation, and may easily become pioneers 405

Oldest genus usually first arrival in country (in fam.) 192-3, 449, 453; tribes in large fams tend to be smaller 187-8

Onagraceae by size 174; in Spain and Brit. 453

One char. may produce a second; possible increase 211-12

One divergent child may produce a whole new sp. 258;

genus through another 150 and see Bower; new sp. required every 15-30 years, in one place only 16

"Ones" 38%, 488

Orchidaceae by size 326-7, 10, 157

Order begins to show in new places 121, 259, 315 (and see Arithmetical)

Orders or cohorts 323, 330-334, 357, 382, 428-9; to which the first 45 fams. belong 330; first 30 gen. 334

Orders by leaders 428-9; steps above fam. hard to follow 382; show well at top 334 &c.

Origin and dispersal 168, 187-8, 190; of *Alismaceae* 375

Orkney Is. 23-4

Orthotropous ovule 239-40

Osyrideae (Santal.) 90

Other factors begin to show effects when genus reaches 15-20, 306

Othonninae (*Comp.*) 442-3

Outlying genera the largest 27, 49, 245; islands of Brit. 466 and see Islands for others

Outside conditions an important guiding force in evol. 276-7 seq., 302

Overlap of chars. tends to increase with age 210; of genera at foot of list 198

Overriding genus 131, 146-8, 151, 232, 283, 291-2, 301, 363, 437

Overtaking of leader by followers 429

Owen, Sir Richard 331

Oxalidaceae 195

Paeonieae (Ran.) 11, 33-4, 50, 57, 362

Pagamea (Rubi.) 189

Pairs of fams (§) 342

Palmaceae by size 473-4; in islands, espec. Seychelles 473-474

Palmgren 48, 465

Pandora's box 94, 133-4, 147-8, 212, 282

Paniceae (Gram.) 39-41

Pantrop. gen. very old 202

Papilionatae (Legum.) 325, 335

Parallelism of endemism Balkans/Spain 69 seq.; Ceylon/Madras 74; distribn. in *Crucif./Umbell.* 457

Parasites 422

Parent, and child 93, 96, 170, 313; carries both chars. 93, 116; carries possibilities of unexpected chars. 435; survives at a mutation 67, 395

Passage of fams., one into another 92-4

Past history of plants largely a record of invasions 402

Penaeaceae by size 301-2; 279, 295, 301-2, 348, 354, 434

Penzig, O. (teratology) 100-05, 365-6, 371, 436

Perfection of chars. 60

Permutations and combinations 92, 105, 147-8, 189-90, 208, 232, 299, 301, 354, 362-3, 368, (§) 374-6

Peumus (Monim.) 228-9, 234, 241

Philodendroideae (Arac.) 268-9

Phrymaceae 240

Phyllanthoideae (Euph.) 91, 338

Phytolaccaceae 90, 134, 354

Pioneers 405, 410

Piperaceae 288-9, 292-3, 308

Pistia (Arac.) 266

Pityranthe (Tili., e.g.) 157

Plenty of room at the top 98

Pliocene flora (Brit.) 36-7, 62, 167, 183

Podostemaceae 96-7, 99, 240, 276-7, 289-90, 302, 317, 345-6, 357, 360, 363, 366, 481

Pollen patterns (Acan.) 197, 207-10

Polyphyly 15, 96-7, 99, 148-9, 151, 240, 276-7, 289-90, 302, 317, 351-2, (§) 369-71, (§) 435-7

Polyploidy 11

Polyporus 302

Potentialities exist in head in some kind of recessive condition 170-1, 208, 237, 241, 367

Pothoideae by size 269; 267-9

Predictions 17-18, 23, 26-7, 50-1, 64, 76, 79, 82-3, 85, 97, 141, 144, 172-4, 186, 191, 193, 203-4, 243-5, (how done) 248-9, 256, 319-20, 313-4, 453

Primitive types 37

Primulaceae 378

Principle of compensation 251

Prionium (Junc.) 347

Probably any member of a fam. carries all its chars. 133

Problems in distribn., treatment 231, 424-5, 443-7, & Cynar. Inul. and cf. Handling

Production of chars. 211

Progress of evol. 420

Proofs of downward evol. 157-8, 175, 222, 328, 470

Proportion of spp. and gen. to size of leader 187-8; and distrn. Span. and Balk. endcs. 86

Proportional differences upper and lower divisions of fams. 187-8

Proviso of single mutation 96

Psychotria (Rubi.) 189, 212, 311-12, 430; in Hawaiians 162, in Seychelles 469, in Bahamas 471, in Ceylon 472

Puccinia 302

Punicaceae 456-7

Put sub-gen. together, ground better covered 108

Quercetum Roburis dispersal 411

Range of dispersal, in Brit. 83-5, 88; of *Ranunculus* 117; of *Symplocos* 117; of tribal size in *Acanthaceae* 215

Rank and range 307, (Guppy) 313, 225, 307-12, 353; of chars. 188; of mutations 209; of newcomer 97, 353, 398, 425; of fam. 425, of gen. 225; owed to age 313, 425

Ranunculaceae (us) 9-14, 29-39, by size 49-50, 53-7, 65, 70-1, 78, 100-1, 113-7, 167, 251, 254, 319-21, 344, 357, 362-3, 367-8, 371, 375, 377, 387, 389-90, 436, 461, 463, 514

Rate of spp. production, cf. Yule; of spread much the same in groups of allies 169-70, 405

Real discontinuity 89, 202

Recent acquisition of water habit 417

Recessiveness of one char. in divergence 106 seq., 234, 370

Red and black seeds (crab's eyes) 355

Reed swamp community 28

Reflexed calyx 371

Regularity of curves destroyed by splitting 214

Reid, Mrs Clement 36-7, 167, 183, 255-7

Relationship closer as one goes up 225, 234-7; genetic r. crosses taxonomic at times 108; goes with position in dichotomous system, whatever actual structural features may be 207

Relic, relicdom, 21, 32, 34-5, 43-5, 54, 62-3, 68-9, 95, 106, 113-6, 127, 129, 136, 155, 247, 250, 252, 254-9, 304, 310, 312-3, 318-20, 337, 510, and see *AA* and *Evol.*

Representation goes largely with size 176, 179, 449, 487; and cf. 44

Resedaceae, a Medit. fam. 75, 456-7

Retention of char. once acquired 191-2

Reversal of Darwinism 14, 416

Rhamnaceae 165, 170

Rhamnus 107-8, 143, 174, 286, 353, 428

Rhododendron omitted in list on p. 193

Rhodomyrtus sea and mts. 460

Ridley, H. N. 127

Ritigala mountain (Ceylon) 67, 318, 356

Riviera flora 418-20

Rondeletieae (Rubi.) 468

Rubiaceae 85-9, terat 102, 159, 161-2, *Galium* 181-2, 189, 209, 212, 311, 314, 328, 350, 361, 364, 372, 420, 426, 434, by size 468, 469-70; as coast plants 426; common on islands 469-73

Rubus 21, 107, 183-4

Ruellia 195-6, 202, 204

Rutaceae 162

St. Hilaire G. 125, 331, 364

Salix dispersal 21, 308

Salvia (Labi.) 36, 329, 334

Same char. in genus and sub-genus 132; in widely separated places 146-7, 149-50, 170, 192-6, 209-10, 272-3, 435-6; wide with different endcs. in different places 106; selection cannot explain, due to sudden mutation 149

Sameness of leading fams. in tropics 466

Santalaceae 90

Sapindaceae 173

Sapotaceae 174

Saprophytes 422

Sardinian flora 80-2, 475

Sarraceniaceae 252

Satellitic genera 35

Saxifraga(ceae) 189, 70-1; generic sub-leaders 428

Sayers, Miss Dorothy 313

Scandicineae (Umb.) 457

Scaphocalyx (Flac.) 166

Schellenberg 142

Schiedea (Caryo.) e.g. 166

Schismatoclada isl. e.g. 337

Schizandraceae 327

Schizopetalon (Cruc.) 104-5, 365

Schröter C. 461-3

Schumacheria (Dill.) 145-8, 150, 153, 156, 192, 201, 432

Sciaphila (Triur.) 126, 437

Scilly Is. 466

Scirpus, leaders sub-gen. 427

Scolopia (Flac.) 122, 125

Scott, D. H. 169, 257
 Scottish flora a reduced English 48
Scrophulariaceae 25, 42, 69, 81, 84, 173
Scyphanthus (Loas.) div. 166
 Seabeach and mountain conditions, similarity 460
 Second genus or species usually about half the size of first and in another subdivision 191
Securidaca (Polygal.) div. 119
 Selection (including natural) 11-17, 48-9, 63-4, 72, 88, 92-4, 96, 98-9, 103-4, 114, 128, 149-50, 232, 251, 258, 286, 292-3, 298-300, 317-22, 335, 388, 418, 453, 457, 466, 493; a very unsatisfactory explanation 98, 172, 298-303, 388, 457; action of, test on *Cruciferae* and *Umbelliferae* 457; assumptions of 418; based on total efficiency of all chars. 388, 396; by adaptl. value drops out as explanation of dispersal other than local 114, 395; too hard worked 115; cannot control divergent mutation 14-15, nor expect structural perfection 98-9; as "appointments board" 103-4; by age (size) 448; destruction 99, 232; worked too hard 114-5; eliminated 298-9; individual 388; rarely destroys whole sp. 406; involves destruction of transitions 98; little to do with distribution 182-192, 457; or classification 340 &c.; more difficult by it to explain generic and fam. differences 95-9; not responsible for structural differences 96; on combined efficiency of all chars. 388; out of court 211; picks out individuals, not spp. 99, 304, most efficient combination of chars. 105, 388; populations, not types?

258, 254; too local to produce an ordered world 448
Senecio (Comp.) 80, 98, 101, 173-7, 188, 265, 364, 415, 420, 423-4, 430, 433, 438-41; *Senecioneae* by size 177
 Separation of plants with close structural affinity 90-9, 141-2
 Sex ensures rapid divergence 190
 Seychelles 469-74; *Palms* 473-4, *Rubiaceae* 469-70
 Shetland Is. 23-4, 466
Shoreae (Diptero.) 153-5; plants that happened to show a particular set of chars. 155, 366
Shortia (Diapens.) 89
 Shuffling of chars. 190
 Similar conditions, similar results with similar plants 170, 270-3, 283, 289, 485
 Single mutation may cover any gap parent/child 96, 152
 Sinnott, E. W. 202
Siparuna (Monim.) Ch. X. p. 222, esp. 223, 233, 241, 243, 245-6, 224 (map), 353
 Six leading gen. in all fams. down to size of 50, 512
 Size and space 19-22, 25, 50-2, 76-9, 81, 83, 173-90, 201, 327; hitherto neglected 232, 190, 307, 446; of first and second gen. in fam. 191; of gen. for *Dictionary* 190-1
 Small, J. 416
 Small area goes with small size 130; small dispersal goes with late arrival 11; small families 306-7, (§) 341, 391; small genera 129, 165; increasing in number towards centre 49; largely in mountns. and islands 197; mutatns. may be frequent 227-8; tend to be left behind at barriers 46, 402; usually satellites, not relict 32
 Smaller the fam., smaller the proportn. of endm. 74; smaller the genus, the more

likely to be left behind at barriers 46, 402

Soil changes 8, 223

Solanum 35, 174, 329

Solitary divergent child may grow into new sp. 258

Solanum 35, 174, 329

Southern endm. and Brit. flora 68, 82; southern invasion of New Zealand 407-8; southern zone of vegetation 454

Spanish flora 44-49, 57-60, 68, 70-80, 83, 87, 453

Specialisation and localisation 391, 398

Species 20-6, 32-4, 51-4, 58, 61, 65, 70-87, 91-4, 96-100, 105-7, Chs. V, VI, VII, 95-163, 178, 187, 191, 211, 228, 237, (§) 242, 276, 297, 395, 398-9, 404-12, 415; do not compete as units 282, 99, 304; begin with local adaptation 32, 394, 405; formation of new ones seems regularly to involve divergence 107; in first and second gen. closely 2/1 335; Hurst on mutation of specific chars. 105; Komarov on origin 404; lists &c. of endemics 70-3, 74, 77-9, 81, 110-3, Chs. V, VI, VII, 95-163, 318-9, 389, 407-8, 444-6, 466-79; mutations of less than specific rank; new usually within or near to the parental area 201; older the sp. the more rapid its spread 398; selection picks out individuals, not spp. 99, 258, 304; spread of (§) 403, (§) 404, 187-92, 331, 344, 386, 415-29 and cf. Large, larger, largest

Speculation 42, 45, 163, 225-6, 231, 249, 256-7, 261, 295, 331, 344, 386, 460

Splitting into leaders of sub-groups 170-2, 211, 235; does not exorcise the difficulties 108, 140, 146, 150, 279

Spread of fam. 215; of gen. and spp. 9, 57, 403-4

Stachys (Labi.) 452

Standard for chars (none) 144

Stearn, W. T. 191, 432

Stebbins 317, 393

Stemonoporus (Diptero.) e.g. 153

Sterculia WE 114

Stewart, C. B. 130, 164

Stewart I. New Zd. 465

Stilpnophytum (Comp.) div. 132

Stratiotes (Hydroch.) 345, 468

Stress in *Podostem.* always in direction of dorsiventrality 277

Strobilanthes (Acan.) 204-5

Structural alterations indifferent to selection 105; discontinuity 90; not necessarily explicable by same rules as geographical (real) 143, 281, 287; evolution little to do with 99; selection little to do with dispersal 182, 246; taxonomy will not harmonise with dispersal, or with divergent mutation 148

Struggle for existence 493

Study of problems in geographical distribution 424-5, 437, 443-6, 469

Styliadiaceae 361

Styracaceae 130-1

Subconscious, use of 248-9, 482

Subdivisions largely from leaders 423; early formed 142; rarely headed by later mutations 236, 334

Subgenera 181

Success 44, 167, 257; "successful" genera 321

Suggested origin of *Araceae* 287

Summerhayes, V. S. 469

Superior ovary in *Gaertnera* (Rubi.) 361

Super-Mendelism 103, 122-4, 134, 263, 297, 311, 392, 485

Supporters of selection driven to refuge of destruction of intermediates 249

Surname distribution *Evol* pp. 35-9
 Survival of parent at mutation ensures production of hollow curve, 170, 395; an important new conception 304
 Survivors must be locally adapted 38
 Swainson, W. 294-5
 Swamping 202, 478, 482
 Symplocos, dispersal 117, 131
 System of classification in use largely artificial 155, 158, 175

Tamaricaceae 83

Tansley, A. G. 7, 26-8, 37, 410-5
 Taxonomy 103, 107-8, 115, 125-7, 149-50, 170 seq., 183-214, 225-6, 246, 266-80, 303, 307, 316, 322-72, (§) 352, 498; a necessary corollary of DDM 340, 250; age and size ignored 313; and geographical discrepancy 352; based on structure only cannot readily be harmonised with geographical distribution 150-1; breaking up of head, but with its survival, into subheads 143-4, 149-50, 170-87, 193-214, 245-6, 250-1, 267-72, and cf. Leaders; breaking up tends to produce more confusion 91; changes partly at any rate due to changes of conditions 105; often cannot be directly structurally united to original 115-6; chars. 128; complexes 211; difficult to reconcile with DDM or dispersal 150-1; classification has too narrow a base 197; classification of individual fams. see fams.; complexes 211; discrepancy with geographical relations (§) 352, 285; division depends on divergence 322; early marked out 309, 316, 423; tends to destroy arithmetical relations 214, 316; great taxo-

nomic divisions of early appearance (§) 353, 266-89, 423, 315, 322, and cf. *Comp.* 174 seq.; importance of char. dependent on age 392-3; improvement of methods makes disc. incidence of chars. more marked 115; maxims (§) 351; subdivision much used in taxonomy 354; will not give proper genetic classification 340-1, nor always harmonise with geographical distribution 334-5; taxonomic classification 205-6; value of a char. 126-7 &c.
Teratology 100-05, 118, 362, 392, 435-6; evidence of incomplete evolution 104; proves that a char. not shown by parent may be given complete and perfect to offspring 104, 435-6; seems to show that any genus may carry almost any char. 104; suggests super-Mendelism and kaleido. mutation 104
 Termination of area of dispersal 243-4
 Tertiary flora, see Canaries, and 443, 476-7
Tetracera (Dillen.) artificial grouping with structural discontinuity 148, more satisfactory result by throwing all together 148 seq.
Tetrachondraceae 158
Tetraplasandra (Aral. e.g.) 145
Thalictrum (Ran.) 49, 50, 55, 100
 The early bird gets the worm 8 and others
 Theory of natural selection trying to work evolution backwards 15; of upward change 386, of downward 391, 396-7
 There is plenty of room at the top, 98, 410
 Things left at a barrier smaller and more local 45-6
 Thompson H. S. 418-9

Thraulococcus (Sapind., div.) 132
 Thunbergia (oideae) 216, 307-8
 Tiliaceae 135, 157, 366
 Tillandsia (Brom.) 348
 Time all-important 66, 405
 Tinospora (Meni., div.) 127
 Tithonia (Comp.) 406-8
 To him that hath shall be given 8, 13, 38, 82, 167, 297, 307-9, 314, 321, 412, 488 (law 15)
 Top Medit. gen. all in Brit. 79
 Tops of fams. for early sub-groups 172, 250, 310
 Tournefortia (Borag.) 111
 Transition forms, destruction of 253, 314, 88-9; very hard worked 109, 115; inherent in selection 98; need no longer be sought 167; killing-out illogical 206; seem rather appearances of other chars. by kaleido. mutn. 308
 Travel 8, 9, 48, 397, 405
 Treatment of problems in distribution, e.g. Inuleae in Australia 443-7, 468-91
 Trees, shrubs, herbs 420
 Triangle of distribution 23
 Tribes of Acanthoideae 216, 327, and cf. Leaders
 Trichadenia (Flac., e.g.) 151-2
 Trifolium, disp. in Brit. 20-1
 Trigonostemon (Euph., div.) 112
 Trimen, H. 289
 Trimenia (Monim.) 229-30
 Tristegineae (Gram.) 39
Tristichaceae 240
 Triumfetta (Tili.) WE 114
Triuridaceae by size 437
 Trollius (Ran.) 10, 48
 Trop. fams. in Brit. 193
 Tubuliflorae (Comp.) 259
 Two chars. however different may have to coalesce somewhere in ancestry 228
 Tyndall on glaciers 266, 480
 Type of vegetation as barrier *AA* 50-1
 Types of receptacle in *Monimiaceae* 241
 Typha in Brit. 28
 Ultimate factors of dispersal 449
Umbelliferae 10, 25, 51-2, 57-8, 69-71, 81, 84, 86, 172-3, 186-8 (parallelism with *Comp.* 186-7), (with *Crucif.*) 457
 Unexpected appearance (change) of char. 134-6, 151-3, 435-7
 Unimportant chars. 424
 Union of divergent chars. must occur in ancestry 297
 "Unsuccessful" fams. 11, 19
 Uppermost tribes tend to be smaller in large fams. 327
 "Upward" replaced by "downward" evolution 386
 Upward step, family to order 323, 357
 Use of inductive methods 231
 Useless chars. *AA* 209
 "Usually" and similar terms, much used in taxonomy 128, 302, 355, 363; neutral or indifferent, 303
 Value of char. 128, 133, 145, 168-70, 185, 188, 197, 217-9, 341; no standard 144, 170; of given char. in fam. etc. 126 seq.
 Vancouveria, development 432
 Variation, each may bring others 264; in size of tribes 327, 447; at each birth 345
 Varieties in Brit. flora 186, in Indian 351; in older genera 186; in *Hieracium* &c. 181-4; of situations to which *Ranunculus* suited 12-13
 Verbal anodynes 12, 118
 Verification of all predictions makes work hold together 256-7
 Very difficult to make a genetic group 236-7
 Very early genera probably smaller 344; few really widely

dispersed spp. at head of a genus 117
Viable mutations 100, 103
Vice-countries 9
Violaceae 160
Vital factors 45, 50-5, 61, 81, 184-5, 207-303, 448
Vries, H. de 96, 125, 254-5

W usually parent in most W and E combinations 106-8, 110-1
Water carriage 48-9, 202; plants 36, 48-9, 202, 240, 289-90, 349-50, 360, 388, 417, 421, 434-5; and see *Podostemaceae*
Watson, H. C. 9, 25
WE and WEE combinations, Ceylon &c. 106, 110-3, 118-9, 121-6
Wegener separation 202, 456
Went, F. A. F. C. 124, 298
West Indies, *Acanthaceae* 203, *Araceae* 466-7
Wet and dry regions, Ceylon, Madras 402
What really matters is age of char. 170, 226
Wheeler, L.R. 490
Whole tree of a fam. may now exist on earth 303, 343
Whymper 463
Wide dispersal abroad goes with wide at home 16-7, 168
separation of plants with close structural relationship, 88, 90, 225, 327-33

Wides; common parent carries potentialities of both char. or neither 93; usually occur near margin(s) of genus 320; parents of endemics 106-8, 110-1
Wiesner, J., 374
Willis, J. C. *Anemone* 113; *Canaries* 82; *Cardamine* 92; *Dilleniaceae* 96; *Maldives* 49; *Overrider* 283, 363; *Podostemaceae* see P.; *Special Creation* 124; *Tristichaceae* 240; *Working Hypothesis* 49, 96, 87, 97, 192, 225, 277, 316, 388, 431 &c.
Work in this book hangs together well 144, 256-7, 259, 263, 319-20, 388
Working hypothesis, see Willis
Writer has always stipulated for possible complete specific or generic change at one mutation 228, 481

Xerophytism 418

Young beginners 64-5, 67, 95, 387
Yule, G. Udny 16, 100, 124, 185, 219 (cataclysm), 300, 320, 387, 392, 431, 452
Yule and Willis 16, 124, 185, 300, 320, 392

Zeuner 386
Zingiberaceae 112, 174
Zones in New Zealand 477-8;
Zones of vegetation 454