

**Zeitschrift:** Boissiera : mémoires de botanique systématique  
**Herausgeber:** Conservatoire et Jardin Botaniques de la Ville de Genève  
**Band:** 70 (2016)

**Artikel:** A taxonomic revision of the genus *Noronhia* Stadtm. ex Thouars (Oleaceae) in Madagascar and the Comoro Islands  
**Autor:** Hong-Wa, Cynthia  
**Kapitel:** 1: *Noronhia alleizettei*  
**DOI:** <https://doi.org/10.5169/seals-1036126>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 27.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

**1. *Noronhia alleizettei*** Dubard in Bull. Mus. Hist. Nat. 13: 551. 1907 (Fig. 2A).

**Typus:** MADAGASCAR. **Prov. Mahajanga:** Ouest, Morondava (Menabe), [20°17'S 44°17'E], s.d., *Grevé 217* (holo-: P [P00418079]!; iso-: K [K000233201, K000233202, K000233203] images seen, P [P00418078, P04046978, P04046979, P04046981, P04046982, P04046983]!).

= *Noronhia ecoronulata* H. Perrier in Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 2: 287. 1949. **Typus:** MADAGASCAR. **Prov. Mahajanga:** Ouest, lopy entre Andranomavo et Sitampiky (Ambongo), IX.1903, *Perrier de la Bâthie 1602* (holo-: P [P00418115]!; iso-: P [P00418116]!).

**Description** *Tree* to 15 m tall, trunk to 12 cm diameter; young twigs cylindrical, 0.7-1.3 mm diameter, glabrous; bark dark to medium gray, smooth to rugose. *Leaves* opposite, deciduous; bud scales persistent; blades medium green above, lighter below, narrowly elliptic, 4-8.5 × 0.8-2.5 cm, coriaceous, glabrous, domatia absent, base acute, margin flat to slightly revolute, apex retuse to sometimes acuminate, the acumen 0-7 mm long, midrib slightly sunken above, raised below, secondary veins barely visible, 7-12 per side, 4-13 mm apart, looping 0.7-3 mm from the margin; petiole medium gray, 3-8 × 0.6-1.4 mm, entirely woody, glabrous. *Flowers* fasciculate; pedicel 1.5-6 mm long, sparsely pubescent; calyx moderately pubescent outside, glabrous inside, lobes triangular, 1-2.2 × 0.3-1.2 mm; corolla white, urceolate, 2-5 mm long, glabrous on both sides, the tube 1.5-4.5 mm long, lobes ovate, apex rounded; corona absent; stamens 1.4-2 mm long, anthers oblate to orbicular, 1.2-1.6 mm long; pistil 1.3-3.2 mm long, stigma capitate. *Fruiting* pedicel 3-12 × 0.6-1.5 mm; young fruits green, reddish black when mature, subglobose, 8.5-13.5 × 7-13.5 mm, surface smooth, sometimes covered with white dots, apex apiculate; dry pericarp 0.2-0.8 mm thick; endocarp woody; seed 5.5-8 × 4-6.5 mm.

**Distribution, ecology and phenology** *Noronhia alleizettei* occurs in low-elevation dry forests in the west, from Ankarafantsika to Morondava (Fig. 3). It produces flowers and fruits all year except in September, with a peak between October and February.

**Conservation status** There were 35 collections representing 25 localities available for analysis, which resulted in an Extent of Occurrence (EOO) of 61,385 km<sup>2</sup> and an Area of Occupancy (AOO) of 84 km<sup>2</sup> that is certainly much larger given the presence of uncollected areas of suitable habitat. Sixteen subpopulations representing 13 locations were estimated, of which six occur within the network of protected areas (Ankarafantsika, Beanka, Bemaraha, Corridor Bongolava, Menabe-Antanimena, and Namoroka). Therefore, *N. alleizettei* is assigned a preliminary status of “Least Concern”.



**Fig. 2.** Photographs of *Noronhia* Stadtm. ex Thouars. **A.** *Noronhia alleizettei* Dubard [Hong-Wa 622]; **B.** *Noronhia aminae* Hong-Wa [Hong-Wa 555]; **C.** *Noronhia ankaranensis* (H. Perrier) Hong-Wa [Hong-Wa 551]; **D.** *Noronhia capuronii* Bosser [Trigui 536].

Photos : taken by respective collectors

**Notes** *Noronhia alleizettei* can be recognized by its narrow leaf blades, fasciculate, white flowers and small subglobose fruits. PERRIER DE LA BATHIE (1949, 1952) distinguished *N. ecoronulata* H. Perrier from *N. alleizettei* by the absence of a corona. However, none of the examined flowers of *N. alleizettei* had a corona and the original description mentioned only a simple rim at the base of the corolla (DUBARD, 1907) that may actually correspond to the thickening of the base of the corolla when the corona is absent (PERRIER DE LA BATHIE 1949, 1952). The two species, as recognized by PERRIER DE LA BATHIE (1949, 1952), are similar in all other aspects and are therefore synonymized, with *N. alleizettei* having priority. The specimen labeled as *Alleizette 4* (P [P00418078]) should be regarded as an isotype of *N. alleizettei* rather than a syntype since it is a duplicate of *Grevé 217* as noted by H. Perrier de la Bâthie: “part du précédent indiquée par erreur comme provenant de la forêt de la Mandraka (près de Tananarive), c’est à dire d’une forêt ombrophile du versant oriental, située à 1200-1400 m d’altitude, alors qu’elle provient en réalité du sous-bois, de forêt tropophile, située sur le versant occidental, au voisinage de la mer, dans la partie sud du secteur «Menabe» du domaine occidental (18 Juillet 1948)”. The specimen *Decary s.n.* (P [P03558887]) includes four fragments, of which three are assigned to *N. alleizettei* and the fourth to *N. tropophylla* (H. Perrier) Hong-Wa & Besnard.

#### Additional specimens examined

**MADAGASCAR. Prov. Mahajanga:** Beanka, Sarodrano, 18°03'34"S 44°31'00"E, 394 m, 23.II.2012, *Bolliger et al. 245* (G, MO, TEF); Beanka, Andalaposa, 18°00'54"S 44°29'53"E, 237 m, 18.III.2012, *Bolliger et al. 307* (BR, G, K, MO, P, TEF); Domaine occidental, s.d., *Decary s.n.* (P); Ampijoroa STF, c. 30 km N d'Andranofasika, 16°20'S 46°51'E, 200 m, 13.IV.1984, *Dorr & Koenders 3026* (K, MO, P, TAN); Beanka, Sarodrano, 18°03'01"S 44°31'09"E, 450 m, 5.III.2012, *Hanitrarivo et al. 263* (BR, G, K, MO, P, TEF); Ankarafantsika PN, 16°18'54"S 46°48'46"E, 154 m, 13.III.2009, *Hong-Wa 622* (MO, P, TAN); *ibid. loc.*, 16°17'45"S 46°49'20"E, 160 m, 14.III.2009, *Hong-Wa 628* (MO, P, TAN); *ibid. loc.*, 16°19'07"S 46°48'13"E, 198 m, 14.III.2009, *Hong-Wa 631* (MO, P, TAN); Tsingy de Bemaraha, S of the Manambolo river, 19°09'S 44°49'E, 50 m, 13.XII.1996, *Jongkind et al. 3524* (G, MO, P, TAN, WAG); Antsalova, Bevitiky, Kinanaoka, Tsingy de Bemaraha RN9, 18°38'44"S 44°42'24"E, 180 m, 6.III.1995, *Labat 2667* (G, K, MO, P, TAN); Ampijoroa STF, 16°19'S 46°49'E, 7.IV.1988, *Leeuwenberg 13864* (MO, P, TAN); *ibid. loc.*, 16°19'S 46°49'E, 150 m, 20.VI.1987, *Phillipson 1911* (MO, P, TAN); Beanka, Kinahango, 18°01'34"S 44°30'27"E, 306 m, 14.III.2012, *Rakotozafy et al. 80* (G, MO); Port Bergé, Bongolava, forêt de Marosely, 15°38'58"S 47°35'03"E, 217 m, 17.XI.2004, *Razakamalala et al. 1716* (MO, P, TEF); Ambondromamy, forêt de Befotaka, 16°22'39"S 47°10'28"E, 17.IV.2007, *Razakamalala et al. 3388* (MO, P, TAN); Maintirano, Ambinda, 18°04'S 44°30'E, 203 m, 19.X.2009, *Razakamalala et al. 4426* (MO, P, TAN); Tsaramandroso, Bevazaha, [Ankarafantsika PN], [15°59'S 46°56'E], 5.X.1948, *Réserves Naturelles 1671* (MO, P); Ankarafantsika, [16°11'S 47°06'E], 3.XI.1950, *Réserves Naturelles 2020* (G, MO, P); Soalala, Andranomavo, [16°21'S 45°17'E], 71-227 m, 27.III.1954, *Réserves Naturelles 6142* (P, TEF); Ankarafantsika, [16°11'S 47°06'E], *Service Forestier 5bis* (P); *ibid. loc.*, *Service Forestier 55* (K, P); Route Morondava-Belo, Andranomena, 20°10'30"S 44°25'30"E, 10 m, 18.X.1962, *Service Forestier 21073* (TEF).






Fig. 3. Distribution maps of species of *Noronhia* Stadtm. ex Thouars: *N. alleizettei* Dubard to *N. capuronii* Bosser.