

Zeitschrift: Boissiera : mémoires de botanique systématique
Herausgeber: Conservatoire et Jardin Botaniques de la Ville de Genève
Band: 70 (2016)

Artikel: A taxonomic revision of the genus Noronhia Stadtm. ex Thouars (Oleaceae) in Madagascar and the Comoro Islands
Autor: Hong-Wa, Cynthia
Kapitel: 4: Noronhia armandiana
DOI: <https://doi.org/10.5169/seals-1036126>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 11.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

4. *Noronhia armandiana* Hong-Wa, spec. nova (Fig. 5).

Typus: MADAGASCAR. Prov. Fianarantsoa: Atsimo-Atsinanana, Farafangana, Maha-bo-Mananivo, forêt d'Agnalazaha, 23°08'42"S 47°42'18"E, 31 m, 22.IV.2004, Ludovic 759 (holo- : MO-6615572!; iso- : G, P, TAN).

Diagnosis *Noronhia armandiana* Hong-Wa can be distinguished from its congeners by its dense branching, its whitish bark, its oblong leaf blades and its rounded fruits.

Description Trees to 13 m tall, trunk to 15 cm diameter; young twigs cylindrical, 1-1.5 mm diameter, glabrous; bark light gray, smooth. Leaves opposite, persistent; bud scales persistent; blades dark green above, lighter below, oblong, 5.5-10.5 × 1.5-3.5 cm, coriaceous, glabrous, domatia rare, base attenuate, margin revolute, apex shortly cuspidate, the cusp 5-10 mm long, midrib slightly sunken above, distinctly raised below, secondary veins barely visible, 7-12 per side, 6-12 mm apart, looping 1-4 mm from the margin; petiole light gray, 4-8 × 1-2 mm, entirely woody, glabrous. Flowers fasciculate; pedicel 3-5 mm long, moderately pubescent; calyx moderately pubescent outside, glabrous inside, lobes triangular, 1 × 1.8 mm; corolla reddish brown, urceolate, 4.8 mm long, glabrous on both sides, the tube 3 mm long, lobes ovate, apex rounded; corona present, 1.5 mm long, undivided; stamens 1.8 mm long, anthers oblong, 1.2 mm long; pistil 1.8 mm long, stigma capitate. Fruiting pedicel 6-8 × 1.2-1.8 mm; young fruits green, dark brown when mature, globose, 12-15 × 12-15 mm, surface covered with white dots, apex flat to bluntly pointed; dry pericarp 0.9-1.6 mm thick; endocarp crustaceous; seed 7-8.7 × 5.5-8.3 mm.

Etymology This name honors Armand Randrianasolo, Curator at the Missouri Botanical Garden in St. Louis, for his dedication to the conservation of the Malagasy flora and particularly for his role in the conservation of Agnalazaha forest, where the type specimen was collected.

Distribution, ecology and phenology *Noronhia armandiana* occurs in littoral to low-elevation humid forests at Manombo in the east (Fig. 3). It produces flowers and fruits from November to April.

Conservation status *Noronhia armandiana* is currently known only from four collections representing four localities. With an EOO of 42 km², an AOO of 12 km², and three subpopulations representing three locations, none of which occurs in a protected area, although one is located just outside Manombo RS, *N. armandiana* is assigned a preliminary status of “Endangered” [EN B1ab(i,ii,iii,iv,v)+2ab(i,ii,iii,iv,v)]. Areas outside of Manombo RS have been gradually converted into agricultural land, reducing the suitable habitat

available to this species. In addition, wood harvesting and sporadic wildfire such as that of 2005, which burned 200 ha of forest (BIRKINSHAW et al., 2005), also constitute a substantial threat.

Notes *Noronhia armandiana* most closely resembles *N. marinae* Hong-Wa, from which it differs by its oblong (vs. linear to lanceolate) and shortly cuspidate (vs. acuminate) leaf blades. Its flowers are fasciculate, urceolate and reddish brown (vs. thyrsoid, cupuliform and purplish pink) and its fruits have a flat (vs. apiculate) apex and crustaceous (vs. woody) endocarp. This species can be recognized by its dense branching, whitish bark, oblong leaf blades, and rounded fruits.

Paratypi **MADAGASCAR. Prov. Fianarantsoa:** Forêt Classée near Manombo RS, 23°03'16"S 47°40'28"E, 100 m, 15.XI.2001, McPherson & Rabenantoandro 18419 (MO, TEF); Farafangana, Mananivo, labomary, 23°03'34"S 47°40'23"E, 0-50 m, 6.II.2001, Rabenantoandro et al. 442 (G, MO, P, TAN); Mahatsinjoriaka, [23°02'10"S 47°44'50"E], 0-10 m, 28.II.1956, Service Forestier 16264 (P, TEF).