

<b>Zeitschrift:</b>	Boissiera : mémoires de botanique systématique
<b>Herausgeber:</b>	Conservatoire et Jardin Botaniques de la Ville de Genève
<b>Band:</b>	70 (2016)
<b>Artikel:</b>	A taxonomic revision of the genus Noronhia Stadtm. ex Thouars (Oleaceae) in Madagascar and the Comoro Islands
<b>Autor:</b>	Hong-Wa, Cynthia
<b>Kapitel:</b>	7: Noronhia brevituba
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-1036126">https://doi.org/10.5169/seals-1036126</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 11.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

**7. *Noronhia brevituba*** H. Perrier in Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 2 : 290. 1949 (Fig. 6A).

**Typus:** MADAGASCAR. Prov. Toamasina: Forêt d'Analamazaotra, [18°55'48"S 48°25'48"E], II.1912, Perrier de la Bâthie 8820 (holo-: P [P00573414]!; iso-: K [K000233197] image seen, P [P00573415, P00573416]!).

= *Noronhia ambrensis* H. Perrier in Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 2 : 297. 1949. **Lectotypus** (designated here): MADAGASCAR. Prov. Antsiranana: Ouest (Nord): camp d'Ambre, [12°30'S 49°14'E], 900 m, XI.1932, Perrier de la Bâthie 18839 (P [P00418074]!).

**Description** Trees to 11 m tall, trunk to 15 cm diameter; young twigs cylindrical, 0.6-1.5 mm diameter, glabrous; bark medium to light gray, smooth, with scattered lenticels. Leaves opposite, persistent; bud scales rarely persistent; blades dark green above, lighter below, elliptic to obovate, 3-7.5 × 1.4-4.1 cm, chartaceous, glabrous, domatia absent, base acute to attenuate, margin flat to slightly undulate, apex cuspidate, the cusp 3-19 mm long, midrib flat above, raised below, secondary veins conspicuous mostly below, 5-10 per side, 5-16 mm apart, looping 1-4.5 mm from the margin; petiole yellow, 3-10 × 0.6-1.8 mm, not woody, glabrous. Thyrse fasciculate, pauciflorous, diffuse; peduncle 5-15 mm long, glabrous; pedicel 6-30 mm long, glabrous; calyx glabrous on both sides, lobes triangular, 1-1.5 × 0.7-2 mm; corolla yellow to pale lime green, sometimes tinged orange-brown with age, campanulate to subrotate, 3.5-7.5 mm long, glabrous on both sides, the tube 0.5-2.5 mm long, lobes ovate, apex slightly acute; corona present, 1-2 mm long, undivided; stamens 1.4-3 mm long, anthers widely oblong to almost square, 0.9-1.8 mm long; pistil 1.4-2.3 mm long, stigma capitate. Fruiting pedicel 3-19 × 0.5-1.7 mm; young fruits green, dark red when mature ovoid, 13-16.5 × 8-13 mm, surface smooth, apex flat to apiculate; dry pericarp 0.2-0.6 mm thick; endocarp woody; seed 9-12 × 5-7.5 mm.

**Distribution, ecology and phenology** *Noronhia brevituba* occurs in mid- to high-elevation humid forests from Montagne d'Ambre in the north to Ranomafana in the south (Fig. 3). It produces flowers and fruits from September to May.

**Conservation status** The assessment was based on 28 collections representing 26 localities and yielded an EOO of 133,270 km<sup>2</sup>, an AOO of 92 km<sup>2</sup>, and 20 subpopulations representing 15 locations, of which 10 occur within protected areas (Analamazaotra, Anjanaharibe-Sud, Corridor Marojejy, Loky-Manambato, Makira, Manongarivo, Masoala, Montagne d'Ambre, Ranomafana, and Zahamena). Given these results and the likelihood that the AOO is much greater than the estimated 2,000 km<sup>2</sup>, *N. brevituba* is assigned a preliminary status of "Least Concern".

**Notes** As currently circumscribed, *Noronhia brevituba* can be recognized by its non-woody petioles, chartaceous leaf blades and short-tubed, yellowish corolla. It mainly differs from *N. linoceroides* H. Perrier by its cuspidate (vs. acuminate) leaves, campanulate (vs. subrotate) flowers, capitate (vs. bilobed) stigma and apiculate (vs. bluntly pointed to rostellate) fruits. These two species belong to the same clade and form a complex (HONG-WA & BESNARD, 2013, 2014), for which additional information (e.g. anatomical data) and broader molecular sampling (e.g. individuals from each population across their range) may be required to disentangle their patterns of variation and relationships.

PERRIER DE LA BÂTHIE (1949, 1952) distinguished *Noronhia ambrensis* H. Perrier from *N. brevituba* based mainly on the length of the corolla tube, placing the first in the section *Noronhiae* and the latter in section *Linoceriae*. However, their range of morphological variation overlaps and material assignable to these two entities is interspersed within a single clade (HONG-WA & BESNARD, 2014), which precludes the recognition of two species. It should be noted that the accessions labeled as *N. ambrensis* on the phylogenetic tree of HONG-WA & BESNARD (2014) were incorrectly assigned a homonym, whose nomenclature is addressed later in this treatment under *N. olearia* Hong-Wa. Specimens assigned to *N. ambrensis* by PERRIER DE LA BÂTHIE (1949) were labeled as *N. brevituba* in the phylogenetic tree. *Perrier de la Bâthie* 18839 is chosen as the lectotype of *N. ambrensis* because it represents more complete material than the other syntype (Richard 192), which should be considered as an excluded syntype and is assigned to *N. olearia* (see p. 95).

#### Additional specimens examined

**MADAGASCAR.** **Prov. Antananarivo:** Andranofeno-Sud, forêt d'Ankafobe, 18°06'11"S 47°11'10"E, 1480 m, 14.XII.1999, Lowry *et al.* 5166 (MO, P); Anjozorobe, forêt de Saridrivotra, 18°16'S 47°54'E, 1570 m, 26.XII.2003, Randrianaivo *et al.* 1005 (CNARP, MO, P, TAN); Tampoketsa d'Ankazobe, forêt d'Antokomaro au PK 166/500 de la route de Majunga (au NO de Manerinerina), [17°57'S 47°07'E], 23.XII.1964, Service Forestier 24004 (G, MO, P, TEF). **Prov. Antsiranana:** Anjanaharibe-Sud RS, 10 km SW of Befingotra, 14°45'S 49°29'E, 1100-1300 m, 7.IX.1997, Birkinshaw *et al.* 487 (G, MO, P, TAN); Montagne d'Ambre PN, 12°32'19"S 49°09'32"E, 1219 m, 23.I.2009, Hong-Wa 579 (G, MO, P, TAN); Daraina, forêt d'Antsahabe, 13°12'37"S 49°33'27"E, 900 m, 17.I.2004, Nusbaumer 1023 (Daraina, G, K, MO, P, TEF); Manongarivo RS, Antsatrotro, 14°04'S 48°17'E, 1700-1846 m, 9-10.V.1993, Rakotomalaza *et al.* 79 (G, K, MO, P). **Prov. Fianarantsoa:** Ranomafana PN, 21°15'S 47°27'E, 1100 m, 11-15.XI.1991, Malcomber *et al.* 1060 (MO, P, TAN). **Prov. Mahajanga:** Befandriana Nord, Matsoandakana, Andranomena, 15°08'02"S 49°20'49"E, 1134 m, 12.II.2008, Bernard 789 (G, MO, P, TAN). **Prov. Toamasina:** Forêt au N de la route de Nickelville, 850 m, 29.XII.1994, Cours 2089, (P); Analamazaotra RS, 18°56'07"S 48°25'06"E, 910 m, 2.V.2010, Hong-Wa & Ortiz 638 (MO, P, TAN); Manakambahiny Est, Zahamena RNI, 17°40'34"S 48°45'31"E, 1100 m, 17.IX.2002, Rakotondrajaona *et al.* 233 (CNARP, MO, P, TEF); Ambatondradama (env. 10 km au NE d'Ankovona), Navana, 15°17'15"S, 50°01'07"E, 500 m, 1-6.X.1997, Ralimanana *et al.* 115 (G, K, MO, P, TAN); Ambatondrazaka, Antanandava, Antenina, 17°30'09"S 48°46'19"E, 917 m, 1.II.2002, Randrianjanaka 697 (CNARP, G, MO, P, TEF).



**Fig. 6.** Photographs of *Noronhia* Stadtm. ex Thouars.

**A.** *Noronhia brevituba* H. Perrier [Nusbaumer 1023]; **B.** *Noronhia candicans* H. Perrier [Ranirison 1066].

Photos: taken by respective collectors