

Zeitschrift: Boissiera : mémoires de botanique systématique
Herausgeber: Conservatoire et Jardin Botaniques de la Ville de Genève
Band: 70 (2016)

Artikel: A taxonomic revision of the genus *Noronhia* Stadtm. ex Thouars (Oleaceae) in Madagascar and the Comoro Islands
Autor: Hong-Wa, Cynthia
Kapitel: 10: *Noronhia candicans*
DOI: <https://doi.org/10.5169/seals-1036126>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 18.05.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

10. *Noronhia candicans* H. Perrier in Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 2: 301. 1949 (Fig. 6B).

Lectotypus (designated here): **MADAGASCAR. Prov. Mahajanga**: Bassin moyen du Bemarivo (Boina), [15°27'S 47°40'E], IX.1902, *Perrier de la Bâthie* 8829 (P [P00418071]!); isolecto-: P [P00418072]!). **Syntypus**: Bassin moyen du Bemarivo (Boina), [15°27'S 47°40'E], 500 m, IX.1902, *Perrier de la Bâthie* 8828 (P [P00418073]!).

Description *Shrubs* to small trees to 5 m tall, trunk to 10 cm diameter; young twigs cylindrical, 0.7-1.8 mm diameter, glabrous; bark medium gray, smooth to slightly rugose. *Leaves* opposite, persistent; bud scales persistent; blades dark green above, lighter below, oblong to broadly elliptic, 4-12 × 1.6-5.2 cm, very coriaceous, glabrous, domatia absent, base acute to truncate, margin flat to slightly undulate, apex acute to acuminate, the acumen 1-16 mm long, midrib sunken above, distinctly raised below, secondary veins barely visible, 7-12 per side, 5-15 mm apart, looping 1-6 mm from the margin; petiole medium to light gray, 3-10 × 1.1-2.7 mm, entirely woody, glabrous. *Flowers* fasciculate; pedicel 3-20 mm long, sparsely pubescent; calyx sparsely pubescent outside, glabrous inside, lobes triangular, 2-2.6 × 1.5-2 mm; corolla purplish brown, urceolate, 4.5-7 mm long, glabrous on both sides, the tube 3-5 mm long, lobes widely ovate, apex obtuse; corona present, 1.2-1.7 mm long, undivided; stamens 1.8-2 mm long, anthers oblong, 1.3-1.8 mm long; pistil 2.2-2.5 mm long, stigma capitate. *Fruiting* pedicel 4-25 × 1-1.5 mm; young fruits green, reddish when mature, ovoid, 10-17.5 × 6-10 mm, surface smooth, apex apiculate to rostellate, the rostellum flattened, rounded, style persistent; dry pericarp 0.3-1.3 mm thick; endocarp woody; seed 7.5-10.5 × 5-6 mm.

Distribution, ecology and phenology *Noronhia candicans* occurs in low- to mid-elevation dry forests from Montagne des Français and Daraina in the north to Sofia in the west (Fig. 3). It produces flowers and fruits from September to March.

Conservation status There were 28 collections representing 21 localities available for analysis, which resulted in an EOO of 27,250 km², an AOO of 68 km², and 14 subpopulations representing 10 locations, of which seven occur within protected areas (Andava-koera-Andrafiarena-Ambohipiraka, Ankarana, Loky-Manambato, and Montagne d'Ambre). The actual AOO is likely larger but would not exceed 2,000 km². In addition, the locations that are not encompassed within protected areas are affected by fire, land conversion, and illegal and/or artisanal mining resulting in habitat degradation and loss and reduction in the number of mature individuals. Therefore, *N. candicans* is assigned a preliminary status of "Vulnerable" [VU B1ab(i,ii,iii,iv,v)+2ab(i,ii,iii,iv,v)].

Notes *Noronhia candicans* can be recognized by its very coriaceous, oblong to elliptic leaf blades, with barely visible venation and its fasciculate, purplish-brown flowers. It differs from *N. crassinodis* H. Perrier by its oblong to elliptic (vs. ovate) leaf blades with an acute to acuminate (vs. mucronulate) apex, barely visible (vs. conspicuous) venation, and purplish brown (vs. reddish) flowers. This species was described based on two syntypes, *Perrier de la Bâthie* 8828 and 8829, of which a specimen of the latter was chosen as the lectotype as it has more reproductive organs.

Additional specimens examined

MADAGASCAR. Prov. Antsiranana: Ankarana RS, 12°54'43"S 49°06'39"E, 180 m, 19.II.1994, *Andrianarisata* 45 (MO, P); *ibid. loc.*, 12°54'36"S 49°06'25"E, 130 m, 16.X.1997, *Bardot-Vaucoulon* 775 (K, MO, P, TAN); *ibid. loc.*, tsingy close to Campement des Anglais, 12°50'47"S 49°06'18"E, 82 m, 25.V.1999, *De Block* 1013 (BR, MO, P, TAN); *ibid. loc.*, from Campement des Anglais towards Campement des Américains, 12°50'47"S 49°06'18"E, 82 m, 13.I.2002, *De Block* 1193 (BR, G, MO, P, TAN); Andrafiarana, Anjahankely, 12°55'00"S 49°20'25"E, 390 m, 27.XII.2010, *Burivalova et al.* 140 (G, MO); Daraina, forêt d'Antsahabe, 13°12'58"S 49°32'55"E, 830 m, 4.XII.2004, *Gautier* 4825 (Daraina, G, K, MO, TEF); Andrafiarana, Anjahankely, 12°54'46"S 49°19'37"E, 410 m, 10.XI.2010, *Gautier & Ranirison* 5382 (G, MO); Marivorahona, Betsimiranja, 12°56'27"S 49°07'21"E, 136 m, 18.I.2009, *Hong-Wa* 566 (MO, P, TAN); Sakaramy, Montagne d'Ambre PN, 12°26'41"S 49°13'45"E, 367 m, 30.V.2010, *Hong-Wa* 699 (MO, P, TAN); Andrafiabe, Mont Antsahabe, 12°54'50"S 49°19'41"E, 408 m, 3.VII.2010, *Hong-Wa* 723 (MO, TAN); Ankarana RS, near Campement des Anglais, 12°54'42"S 49°06'42"E, 240-260 m, 22.V.1993, *Jongkind & Rapanarivo* 964 (MO, P, TAN); Tanambao-Marivorahona, 13°02'54"S 49°09'30"E, 154 m, 4.VII.2005, *Leopold et al.* 119 (CNARP, G, MO, P, TAN); Daraina, forêt de Bekaraoka, 13°06'32"S 49°42'28"E, 216 m, 12.II.2004, *Nusbaumer & Ranirison* 1154 (Daraina, G, K, MO, TEF); *ibid. loc.*, Andranovaka, Ankijomantsina, 13°07'56"S 49°28'20"E, 75 m, 29.X.2005, *Rakotonandrasana et al.* 946 (CNARP, G, MO, P, TAN); *ibid. loc.*, forêt de Binara, 13°14'14"S 49°37'19"E, 310 m, 28.III.2004, *Ranirison* 506 (Daraina, G, K, MO, P, TEF); *ibid. loc.*, forêt d'Ambohitsitondroina, 13°07'56"S 49°28'08"E, 165 m, 7.I.2006, *Ranirison & Nusbaumer* 1057 (Daraina, G, K, MO, TEF); *ibid. loc.*, 13°08'19"S 49°27'58"E, 200 m, 8.I.2006, *Ranirison & Nusbaumer* 1066 (Daraina, G, K, MO, P, TEF); Analabe, [14°03'S 49°43'E], 200 m, 13.IX.1955, *Service Forestier* 15497 (TEF). **Prov. Mahajanga:** Antsohihy, Antonibe, Anjajavy, 15°01'26"S 47°15'34"E, 24 m, 19.VIII.2007, *Rasoafaranaivo et al.* 251 (MO, P, TAN).