

Zeitschrift: Boissiera : mémoires de botanique systématique
Herausgeber: Conservatoire et Jardin Botaniques de la Ville de Genève
Band: 70 (2016)

Artikel: A taxonomic revision of the genus *Noronhia* Stadtm. ex Thouars (Oleaceae) in Madagascar and the Comoro Islands
Autor: Hong-Wa, Cynthia
Kapitel: 16: *Noronhia cordifolia*
DOI: <https://doi.org/10.5169/seals-1036126>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 08.10.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

16. *Noronhia cordifolia* (Labat, M. Pignal & O. Pascal) Hong-Wa & Besnard in Mol. Phylogenet. Evol. 67: 376. 2013 (Fig. 7E).

≡ *Chionanthus cordifolius* Labat, M. Pignal & O. Pascal in Novon 9: 68. 1999.

Typus: MAYOTTE: Mlima Choungi, [12°57'S 45°07'E], 29.XII.1995, Pascal 288 (holo-: P [P00075021]!; iso-: K [K000691510] image seen).

Description *Trees* to 6 m tall; young twigs 1-2.2 mm diameter, glabrous; bark medium gray, slightly rugose. *Leaves* opposite, persistent; blades dark green above, lighter below, ovate, 4.5-9 × 2-3.5 cm, coriaceous, glabrous, domatia absent, base truncate to cordate, margin flat, apex acuminate, the acumen 3-7 mm long, midrib flat above, flat to slightly raised below, secondary veins conspicuous, 8-10 per side, 6-14 mm apart, looping 1-3 mm from the margin; petiole medium gray to brownish, 0.8-5 × 1-2.3 mm, usually entirely woody, glabrous. *Thyrse*s geminate, pauciflorous, diffuse; peduncle 3-8 mm long, glabrous; pedicel 4-12 mm long, glabrous; calyx lobes ovate, 0.7-0.9 × 0.7-1 mm, glabrous on both sides; corolla yellow, subrotate to cupuliform, 2.2-2.5 mm long, glabrous on both sides, the tube 1.2-1.3 mm long; corona absent; stamens 1.8-1.9 mm long; pistil 1.6-1.8 mm long. *Fruiting* pedicel 5-7 × 1.5-2 mm; young fruits green, unseen mature, ovoid, 16.5-23 × 13.5-18 mm, surface smooth, apex flat to bluntly pointed, style persistent; dry pericarp 0.8 mm thick; endocarp woody; seed 10.7 × 10.3 mm.

Distribution, ecology and phenology *Noronhia cordifolia* occurs in mid-elevation humid forest in northwestern Madagascar and on Mayotte in the Comoro Islands (Fig. 9). It produces flowers and fruits from November to March.

Conservation status With only four collections representing two localities, the EOO cannot be calculated. However, it was estimated to have an AOO of 8 km² and two subpopulations representing two locations, of which only one occurs within a protected area (Réserve forestière des Crêtes du Sud, Mayotte). The subpopulation found in Madagascar is currently known only from three collections made in the 1950s-1960s in an area located between Manongarivo RS and Tsaratanana RNI in heavily degraded habitat. Therefore, *N. cordifolia* is assigned a preliminary status of "Endangered" [EN B2ab(iii)].

Notes *Noronhia cordifolia* can be recognized by its short-petiolate leaves, cordate to truncate leaf base, and diffuse inflorescences. It differs mainly from *N. boivinii* by its ovate (vs. oblong to elliptic), acuminate (vs. cuspidate to mucronate) leaf blades, thyrseid inflorescences (vs. fasciculate flowers), and subrotate (vs. urceolate to

campanulate), yellowish (vs. pink to reddish) corolla, and by the absence (vs. presence) of a corona. This species was previously known only from the Comoro Islands (Mayotte). The Malagasy specimens are somewhat different from the type by having slightly shorter petioles that are seldom woody as well as relatively longer inflorescences. They may actually represent a distinct entity but exhaustive comparison with the Comorian specimens is currently unfeasible since the former include only fruits and the latter only flowers. Therefore, a broad *N. cordifolia* is retained here to accommodate this range of variations, pending additional material, from which molecular data could also be obtained.

**Additional specimens
examined**

MAYOTTE: Grande Terre, sommet du Mt Choungi, [12°57'20"S 45°07'55"E], 21.III.2002, *Barthelat & Labarthe* 788 (P). **MADAGASCAR. Prov. Antsiranana:** DIANA, Ambanja, Beankany, [14°04'S 48°42'E], 12.XII.1963, *Rakotozafy* 337 (P, TAN); *ibid. loc.*, 23.VI.1950, *Réserves Naturelles* 38 (TAN); *ibid. loc.*, *Saboureau* 41 (P).