

Zeitschrift: Boissiera : mémoires de botanique systématique

Herausgeber: Conservatoire et Jardin Botaniques de la Ville de Genève

Band: 70 (2016)

Artikel: A taxonomic revision of the genus Noronhia Stadtm. ex Thouars (Oleaceae) in Madagascar and the Comoro Islands

Autor: Hong-Wa, Cynthia

Kapitel: 23: Noronhia densiflora

DOI: <https://doi.org/10.5169/seals-1036126>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 11.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

23. *Noronhia densiflora* Bosser in *Adansonia* ser. 2, 13: 461. 1973 (Fig. 15).

Typus: MADAGASCAR. Prov. Fianarantsoa: Route Farafangana-Vangaindrano, E de Madagascar, [22°49'S 47°49'E], XII.1963, Bosser 18739 (holo-: P [P00413225]!; iso-: MO-4407063!, P [P00413226, P00413227]!, TAN!).

Description Shrubs to trees to 8 m tall, trunk to 12 cm diameter; young twigs cylindrical to subquadangular, 3.5-6.5 mm diameter, glabrous; bark light brown to whitish, flaky. Leaves opposite, persistent; bud scales persistent; blades dark green above, lighter below, oblong, 22.5-62 × 7-17 cm, extremely coriaceous, glabrous, domatia absent, base rounded to cordate, margin revolute, apex acute to mucronate, sometimes rounded, the mucro (0-)3-18 mm long, midrib sunken above, distinctly raised below, secondary veins very conspicuous, 11-18 per side, 25-64 mm apart, looping 7-17 mm from the margin; petiole light brown to white, 8-27 × 4-8 mm, entirely woody, with flaky bark, glabrous. Thyruses fasciculate, pauciflorous, compact; peduncle 7-12 mm long, sparsely pubescent; pedicel 7-15 mm long, sparsely pubescent; calyx sparsely pubescent outside, glabrous inside, lobes widely ovate, 1-1.5 × 1.2-1.7 mm; corolla purple-red, globose, 4-5 mm long, glabrous on both sides, the tube 3.8-4.5 mm long, lobes acute to obtuse, sometimes dentate; corona absent; stamens 2-2.5 mm long, anthers oblong, 1.5 mm long; pistil 3-3.5 mm long, stigma bilobed. Fruiting pedicel 1.5-1.8 × 1.8-2.5 mm; young fruits green, blackish when mature, ovoid, 14-17 × 10-11.5 mm, smooth to rugose, apex apiculate; dry pericarp 0.5-0.9 mm thick; endocarp woody; seed 9 × 6 mm.

Distribution, ecology and phenology *Noronhia densiflora* occurs in littoral to low-elevation humid forests in the east, from Mananjary to Vangaindrano (Fig. 9). It produces flowers and fruits from June to February.

Conservation status The assessment, which used the 12 available collections representing 10 localities, indicated an EOO of 1,220 km², an AOO of 40 km², and seven subpopulations representing six locations, of which three occur within protected areas (Ankarabolava-Agnakatrika and Manombo). In recent years, *Noronhia densiflora* has been mostly collected from around Farafangana and Vangaindrano, but it is known to occur around Mananjary as well. However, the latter subpopulation was only sampled once in 1967 and the area is currently highly degraded as a result of slash-and-burn agriculture and other forms of exploitation. With continuing decline in habitat quality and habitat loss, which also leads to loss of mature individuals, *N. densiflora* is therefore assigned a preliminary status of “Vulnerable” [VU Blab(i,ii,ii i,iv,v)+2ab(i,ii,iii,iv,v)].

Notes *Noronhia densiflora* differs from all other members of the genus by its somewhat lianescent habit when young, flaky bark, thickly coriaceous leaves measuring up to 75 cm long, compact intrapetiolar inflorescences, and globose red flowers.

Additional specimens examined **MADAGASCAR. Prov. Fianarantsoa:** Farafangana, Anankara, Manombo RS, 23°22'47"S 47°35'59"E, 100-200 m, 16.IV.2000, *Andrianjafy et al.* 40 (CNARP, MO, P); *ibid. loc.*, [23°02'S 47°44'E], *Friedmann* 164 (TAN); Anankara, Manombo AP, parcelle 2, 23°00'S 47°42'E, 5.III.2009, *Hong-Wa* 611 (MO, P, TAN); Vangaindrano, Antavikazaha, Androkabe forest, 23°30'47"S 47°30'23"E, 77 m, 14.VI.2011, *Letsara et al.* 1122 (CAS, MO); S of Farafangana, Manombo RS, 23°04'02"S 47°40'14"E, 50 m, 17.XI.2001, *McPherson & Rabenantoandro* 18442 (MO, P); Marofototra, Ankarabolava, Letomanga, 23°30'15"S 47°29'10"E, 50 m, 5.IX.2010, *Randrianarivony et al.* 178 (MO, P, TAN); Tsianofana, Tsararano, 23°26'47"S 47°30'31"E, 135 m, 19.X.2010, *Razafitsalama et al.* 1398 (MO, P, TAN); S de Farafangana, route de Manombo, PK 20-21, 14-17.X.1964, *Service Forestier* 23602 (P, TEF); *ibid. loc.*, 17.X.1964, *Service Forestier* 23643 (P, TEF); Mananjary, Ambodikijy, [21°13'S 48°21'E], 50 m, 25.II.1967, *Service Forestier* 26294 (G, P, TEF); Manombo RS, 23°00'51"S 47°43'41"E, 30 m, 4.II.2006, *Tosh* 339 (BR, MO, P, TAN).



Fig. 15. *Noronhia densiflora* Bosser [Razafitsalama 1398].

Photo: C. Birkinshaw



Fig. 16. *Noronhia disjuncta* Hong-Wa.

A. Flowering branch; B. Abaxial side of leaf blade; C. Inflorescence; D. Flower; E. Inner side of corolla; F-G. Pistil; H. Stamen.

[Randria 16, MO] Drawings : R.L. Andriamiarisoa