

**Zeitschrift:** Boissiera : mémoires de botanique systématique

**Herausgeber:** Conservatoire et Jardin Botaniques de la Ville de Genève

**Band:** 70 (2016)

**Artikel:** A taxonomic revision of the genus Noronhia Stadtm. ex Thouars (Oleaceae) in Madagascar and the Comoro Islands

**Autor:** Hong-Wa, Cynthia

**Kapitel:** 27: Noronhia edentata

**DOI:** <https://doi.org/10.5169/seals-1036126>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

**Download PDF:** 16.05.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

**27. *Noronhia edentata*** (H. Perrier) Hong-Wa, comb. & stat. nov. (Fig. 18A).

≡ *Noronhia emarginata* var. *edentata* H. Perrier in Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 2 : 296. 1949.


**Lectotypus** (designated here): **MADAGASCAR. Prov. Toamasina:** Forêt d'Analamaotra, [18°55'48"S 48°25'48"E], 800 m, s.d., *Perrier de la Bathie* 8809 (P [P00791237]!); isolecto-: K [K000233192, K000233193] image seen; P [P03558849, P03558850]!).

**Syntypi:** **MADAGASCAR. Prov. Toamasina:** Analamaotra RS, 2.IV.1905, *Thouvenot* 58 (K [K000233191] image seen, P [P00701239, P00701240]!); *ibid. loc.*, comm. 3.XII.1934, *Ursch* 82 (P [P03558851]!).

**Description** Trees to 25 m tall, trunk to 35 cm diameter; young twigs cylindrical, 1-6 mm diameter, glabrous; bark medium to light gray, smooth to rugose. Leaves opposite, persistent; bud scales persistent; blades dark green above, medium green below, elliptic to obovate, 5-17 × 2-7.5 cm, coriaceous, glabrous, domatia absent, base acute to attenuate, margin slightly revolute, apex rounded to slightly acuminate, the acumen 0-5(-15) mm long, midrib sunken above, raised below, secondary veins conspicuous, 7-17 per side, 7-25 mm apart, looping 1.5-7 mm from the margin; petiole medium gray, 6-38 × 1.3-4 mm, entirely woody, glabrous. Thyrse fasciculate, pauciflorous, compact; peduncle 8-9 mm long, moderately pubescent; pedicel 1.5-8 mm long, moderately pubescent; calyx moderately pubescent outside, glabrous inside, lobes triangular, 1-1.5 × 0.8-2 mm; corolla white sometimes tinged greenish, urceolate, 3-5.5 mm long, glabrous on both sides, the tube 1.2-3 mm long, lobes widely ovate, apex obtuse; corona present, 1.4-2.2 mm long, undivided; stamens 1.8-2.5 mm long, anthers widely oblong, 1.1-1.9 mm long; pistil 1.4-2.5 mm long, stigma capitate. Fruiting pedicel 2-9 × 1.8-3.6 mm; young fruits green, reddish black when mature, ovoid to subglobose, 12-31 × 8.5-26 mm, surface smooth, sometimes covered with a white pellicle, apex flat to bluntly pointed or apiculate; dry pericarp 1.3-4.2 mm thick; endocarp woody; seed 11.5-21.5 × 7-18 mm.


**Distribution, ecology and phenology** *Noronhia edentata* occurs in low- to high-elevation humid forests in the east, from Sambava to Ambondrombe (Fig. 19). It produces flowers and fruits from September to May.

**Conservation status** The assessment was based on 27 collections representing 19 localities, resulting in an EOO of 38,642 km<sup>2</sup>, an AOO of 68 km<sup>2</sup>, and 13 subpopulations representing 12 locations, of which six occur within protected areas (Analamaotra, Betampona, Makira, Makirovana, Masoala, and Zahamena). Being widespread and present in many protected areas, *Noronhia edentata* is assigned a preliminary status of “Least Concern”.



**Fig. 18.** Photographs of *Noronhia* Stadtm. ex Thouars. **A.** *Noronhia edentata* (H. Perrier) Hong-Wa [Razanatzima 266]; **B.** *Noronhia crassiramosa* (H. Perrier) Hong-Wa [A. Randrianasolo s.n.].

Photos: F. Rakotoarivony



**Fig. 19.** Distribution maps of species of *Noronhia* Stadtm. ex Thouars. *N. edentata* (H. Perrier) Hong-Wa to *N. insularis* (Labat, M. Pignal & O. Pascal) Hong-Wa & Besnard.

**Notes** *Noronhia edentata* can be recognized by its leaves, which have broadly elliptic blades and long, woody petioles, its white greenish flowers and its large fruits, with thick pericarp. There is a slight leaf heteroblasty in this species. In particular, the leaf base is narrower and the acumen longer (up to 15 mm long) in seedlings and saplings than in adult plants. This taxon was previously recognized as a variety of *N. emarginata*, from which it differs by its habit, leaf shape, and flower color and shape. Moreover, these two entities belong to separate clades (HONG-WA & BESNARD, 2013, 2014). PERRIER DE LA BÂTHIE (1949) described *N. emarginata* var. *edentata* based on the following syntypes: *Perrier de la Bâthie* 8809, *Thouvenot* 58 and *Ursch* 820; the latter is a typographical mistake for *Ursch* 82. The overall quality and completeness of *Perrier de la Bâthie* 8809 justify the selection of one of the specimens at P as the lectotype.

**Additional specimens examined**

**MADAGASCAR.** **Prov. Antsiranana:** Ambatobiribiry, [14°11'S 50°05'E], 50-345 m, 29-30.XI.1950, Humbert & Capuron 24423 (G, MO, P). **Prov. Toamasina:** Maroantssetra, Ambinanitelo, Ankirindro forest, 15°17'35"S 49°23'59"E, 664 m, 16.I.2003, Antilahimena et al. 1741 (G, MO, P); Manambolo, Amparihibe forest, 15°02'26"S 49°34'59"E, 800 m, 22.II.2003, Antilahimena 1919 (MO); Moramanga, Sahaivo, 18°50'38"S 48°16'54"E, 1006 m, 11.XI.2006, Antilahimena & Edmond 4944 (G, MO, P, TAN, TEF); Ambatovy, 18°51'43"S 48°17'39"E, 1049 m, 9.XII.2006, Antilahimena & Edmond 5092 (MO, P, TAN, TEF); ibid. loc., 18°50'28"S 48°18'16"E, 1085 m, 13.V.2010, Bernard 1566 (MO, P, TAN); Analamazaotra RS, 18°56'12"S 48°25'09"E, 953 m, 2.V.2010, Hong-Wa & Ortiz 650 (MO, P, TAN); Betampona RNI, 17°51'S 49°12'E, 275-650 m, 30.IX.1993, Lewis 711 (K, MO, P, TAN); Analamazaotra RS, [18°56'S 48°26'E], 25.VII, Louvel 88 (P); Andasibe, 18°55'50"S 48°25'04"E, 963 m, 5.XII.2008, Rakotondrafara et al. 932 (MO, P, TAN); Moango, Zahamena RNI, 17°33'32"S 48°53'55"E, 800 m, 4.V.2003, Rakotondrajaona et al. 274 (CNARP, G, MO, P, TEF); Lakato, Ambodigavo, forêt d'Analanjahana, 19°07'57"S 48°24'00"E, 794 m, 31.V.2007, Razanatsima et al. 266 (G, MO, P, TAN); 32 km E of Moramanga, 18°57'S 48°27"E, 870 m, 29.I.1993, Schatz 3452 (MO, P, TAN); Maroantssetra, Farankaraina, [15°25'S 49°52'E], 20-25 m, 16.IX.1952, Service Forestier 6151 (P, TEF); Périnet, [18°56'S 48°26'E], 1000 m, 1.VI.1954, Service Forestier 10374 (P, TEF); Masoala Peninsula, Ambanizana, 15°40'24"S 49°57'51"E, 110-260 m, 28.X.1994, Vasey & Velo 83 (MO, P, TAN).