

Zeitschrift: Boissiera : mémoires de botanique systématique
Herausgeber: Conservatoire et Jardin Botaniques de la Ville de Genève
Band: 70 (2016)

Artikel: A taxonomic revision of the genus *Noronhia* Stadtm. ex Thouars (Oleaceae) in Madagascar and the Comoro Islands
Autor: Hong-Wa, Cynthia
Kapitel: 50: *Noronhia maculata*
DOI: <https://doi.org/10.5169/seals-1036126>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

50. *Noronhia maculata* Hong-Wa, *spec. nova* (Fig. 33C, 35).

Typus: MADAGASCAR. **Prov. Antsiranana:** DIANA, Diégo II, Ramena, env. 2 km à l'O d'Andavakoera, près de la grotte et du campement la Casa Aventura, 12°19'41"S 49°20'26"E, 76 m, 3.VIII.2007, *Hong-Wa et al.* 514 (holo-: MO-6386603!; iso-: CNARP!, G [G00341577]!, P!, TAN!).

Diagnosis *Noronhia maculata* Hong-Wa can be distinguished from other members of the genus by its partially woody petioles, its maculate leaf blades, its short pedicels, and its subrotate, yellow-green flowers.

Description Trees to 7 m tall, trunk to 10 cm diameter; young twigs cylindrical, 0.9-1.8 mm diameter, glabrous; bark medium gray, smooth. Leaves opposite, persistent; bud scales rarely persistent; blades medium green above, lighter below, obovate, 3.5-7 × 1.5-3 cm, subcoriaceous, glabrous, domatia casual, base attenuate, margin flat, apex retuse, midrib slightly sunken above, distinctly raised below, secondary veins conspicuous only below, 6-9 per side, 5-11 mm apart, looping 1-3 mm from the margin; petiole medium gray, 3-7 × 0.9-1.5 mm, partially to entirely woody, glabrous. Thyrses solitary to geminate, multiflorous, compact; peduncle 5-7 mm long, moderately pubescent; pedicel 2-3 mm long, moderately pubescent; calyx moderately to sparsely pubescent outside, glabrous inside, lobes deltate, 1-1.5 × 1-1.5 mm; corolla yellow-green, subrotate, 3-5 mm long, glabrous on both sides, the tube 1.4-1.5 mm long, lobes ovate, apex acute; corona present, 1.3-1.5 mm long, undivided; stamens 1.4-1.5 mm long, anthers obovate, 1.1-1.3 mm long; pistil 2.4-2.5 mm long, stigma capitate. Fruiting pedicel 2-10 × 0.8-1.7 mm; young fruits green, brownish to blackish when mature, ovoid, 11-17 × 7.5-11.5 mm, smooth, often covered with a pellicle, apex flat to rostellate, the rostellum flattened, slightly ridged, rounded, with the persistent style; dry pericarp 0.4-0.8 mm thick; endocarp woody.

Etymology *Noronhia maculata* has distinctively spotted to mottled leaves, for which the Latin word is used to form the specific epithet.

Distribution, ecology and phenology *Noronhia maculata* occurs in low-elevation dry forests from Montagne des Français to Ankarana in the north (Fig. 36). It produces flowers and fruits from December to August.

Conservation status There were 12 collections representing 12 localities available for analysis, which resulted in an EOO of 835 km², an AOO of 44 km², and six subpopulations representing five locations, of which three occur within protected areas (Ankarana, Mon-

tagne d'Ambre, and Montagne des Français). Land conversion, charcoal production, and forest exploitation currently affect non-protected areas, whereas those benefiting from protection suffer from encroachment and illegal logging, all of which will lead to continuing decline in habitat quality and habitat loss as well as loss of mature individuals. Therefore, *N. maculata* is assigned a preliminary status of "Endangered" [EN B1ab(i,ii,iii,iv,v)+2ab(i,ii,iii,iv,v)].

Notes *Noronhia maculata* is similar to *N. ankaranensis*, but can be distinguished by its woody (vs. non-woody) petioles, short (less than 3 mm vs. 3-8 mm long) pedicels, and subrotate (vs. cupuliform) flowers. Its distinctive features include partially woody petioles, maculate leaf blades, short pedicels and subrotate, yellow-green flowers.

The labels accompanying specimens of *De Block 1200* mention flowers that are reddish-brown outside and whitish inside, but the specimen is otherwise morphologically similar to the other specimens cited here. It is not clear whether the distinctive color of *De Block 1200* is related to the age of its flowers, as is sometimes the case in other species, or the result of hybridization. It might also indicate that this collection represents a different entity that is otherwise identical to *N. maculata* in all other macromorphological features. These specimens are best accommodated in *N. maculata* pending further studies of micromorphology, anatomy and/or genetics.

Paratypes **MADAGASCAR. Prov. Antsiranana:** Ambilobe, Marivorahona, Andakoroa, Mahamasina, Ankarana RS, 12°56'34"S 49°07'01"E, 133 m, 7.XII.2006, *Andriamihajarivo et al. 1044* (MO, P, TAN); *ibid. loc.*, Campement des Anglais, 12°54'59"S 49°06'24"E, 180 m, 6.VII.1994, *Andrianantoanina & Rabeharinosy 718* (G, K, MO, P, TAN); *ibid. loc.*, from Campement des Anglais towards Campement des Américains (not beyond first savanna), 12°50'47"S 49°06'18"E, 82 m, 13.I.2002, *De Block 1200* (MO, B, BR, G, P, TAN, US); Marivorahona, Betsimiranja, 12°56'27"S 49°07'21"E, 136 m, 18.I.2009, *Hong-Wa 570* (MO, P, TAN); Sakaramy, Montagne d'Ambre PN, 12°26'40"S 49°13'56"E, 300 m, 29.V.2010, *Hong-Wa 698* (MO, TAN); Mahavanona, Andranomanitra, Ampitiliantsambo, 12°22'49"S 49°22'55"E, 418 m, 15.I.2005, *Rakotonandrasana et al. 906* (CNARP, MO, P, TAN); Montagne des Français, 12°23'13"S 49°22'50"E, 291 m, 17.VI.2004, *Ramananjanahary et al. 24* (CNARP, MO, P, TAN); *ibid. loc.*, 12°21'06"S 49°21'49"E, 170 m, 7.X.2004, *Randriambololomamonjy 28* (MO, P, TAN); *ibid. loc.*, 12°22'05"S 49°19'37"E, 382 m, 8.II.2005, *Randrianarivelo et al. 215* (G, MO, P, TAN); Marivorahona, Betsimiranjana, Ampondrabe, 5 km au SE de Mahamasina, Ankarana, 12°58'57"S 49°09'50"E, 127 m, 10.XII.2007, *S. Randrianasolo et al. 633* (CNARP, G, MO, P, TAN); Ramena, Andavakoera, forêt d'Andranonakomba, 12°21'03"S 49°21'29"E, 50 m, 5.XII.2007, *Ratovoson et al. 1436* (CNARP, G, MO, P, TAN).




Fig. 35. *Noronhia maculata* Hong-Wa.
A. Fruiting branch; **B.** Abaxial side of leaf blade; **C** Flower;
D. Lateral view of corona; **E.** Inner side of corolla; **F.** Pistil; **G.** Fruit.

[Hong-Wa 514, TAN] Drawings: R.L. Andriamiarisoa