

Zeitschrift:	Boissiera : mémoires de botanique systématique
Herausgeber:	Conservatoire et Jardin Botaniques de la Ville de Genève
Band:	70 (2016)
Artikel:	A taxonomic revision of the genus Noronhia Stadtm. ex Thouars (Oleaceae) in Madagascar and the Comoro Islands
Autor:	Hong-Wa, Cynthia
Kapitel:	54: Noronhia martiniana
DOI:	https://doi.org/10.5169/seals-1036126

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 08.10.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

54. *Noronhia martiniana* Hong-Wa, spec. nova (Fig. 39).

- *Linociera obtusifolia* var. *minoriflora* H. Perrier in Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 2: 279. 1949 [nom. inval.].
- *Noronhia obtusifolia* var. *minoriflora* (H. Perrier) Hong-Wa & Besnard in Mol. Phylogen. Evol. 67: 377. 2013 [nom. inval.].

Typus: MADAGASCAR. Prov. Fianarantsoa: Mahabo-Mananivo, forêt de Mahabo, 23°10'34"S 47°42'12"E, 24 m, 4.XI.2001, Rabenantoandro & McPherson 683 (holo-: MO-5587144!; iso-: K, P [P03559153]!, TEF).

Diagnosis *Noronhia martiniana* Hong-Wa can be distinguished from other species of the genus by its long, non-woody petioles, its oblanceolate to obovate leaf blades, its white, campanulate flowers lacking a corona, and its ovoid to oblong fruits.

Description Trees to 12 m tall, trunk to 28 cm diameter; young twigs cylindrical, 1.4-3.4 mm diameter, glabrous; bark medium gray to brown, smooth to rugose. Leaves opposite, persistent; bud scales rarely persistent; blades medium green above, darker below, oblanceolate to obovate, 5.5-16 × 2-7.5 cm, coriaceous, glabrous, domatia absent, base attenuate, margin revolute and slightly undulate, apex retuse to obcordate, midrib sunken above, raised below, secondary veins conspicuous, 9-15 per side, 6-21 mm apart, looping 1-4 mm from the margin; petiole yellowish to reddish, 4-28 × 1-4 mm, not woody, glabrous. Thyruses geminate, multiflorous, compact; peduncle 2-8 mm long, glabrous; pedicel 1-5 mm long, glabrous; calyx glabrous on both sides, lobes triangular, 1-1.7 × 0.9-1.5 mm; corolla white, campanulate, petals almost free, 4.5-6 mm long, glabrous on both sides, the tube 1-3 mm long, lobes ovate, apex obtuse; corona absent; stamens 2-2.5 mm long, anthers oblong, slightly apiculate, 1.7-2 mm long; pistil 1.5-2 mm long, stigma bilobed. Fruiting pedicel 2-9 × 0.9-2.4 mm; young fruits green, reddish black when mature, ovoid to oblong, 13-23.5 × 7.5-19.5 mm, smooth, sometimes covered with a white pellicle, apex flat to rostellate, the rostellum circular, truncate, with the persistent style; dry pericarp 0.2-1.2 mm thick; endocarp woody; seed 10.5-15 × 6-11.5 mm.

Etymology This species is named in honor of Martin Callmander, Head Curator of the Library and publications at the Conservatoire et Jardin botaniques de la Ville de Genève and former Technical Advisor for the Conservation and Research program of the Missouri Botanical Garden in Antananarivo, who played the detective to find hidden material of *Noronhia* at G and P, and whose help in nomenclature and historical collections greatly enhanced the quality of this revision.

Distribution, ecology and phenology

Noronhia martiniana occurs in littoral to low-elevation humid forests in the east, from the Masoala peninsula in the north to around Ste Luce in the south (Fig. 36). It produces flowers and fruits throughout the year except in April and May.

Conservation status


There were 22 collections representing 21 localities available for analysis, which yielded an EOO of 21,396 km², an AOO of 72 km², and 12 subpopulations representing 10 locations, of which four occur within the network of protected area (Agnalazaha, Ambohidena, Masoala, and Tampolo). *Noronhia martiniana* occurs mostly in the narrow band of littoral forests along the east coast of Madagascar and may have an actual AOO of less than 2,000 km². Decline in habitat quality and habitat loss will also very likely to continue as a result of forest exploitation, land conversion, and possibly also gradual sea level rise. Therefore, *N. martiniana* is assigned a preliminary status of “Vulnerable” [VU B2ab(i,ii,iii,iv,v)].

Notes

The taxon described here as *Noronhia martiniana* was first mentioned by PERRIER DE LA BATHIE (1949) as a variety of *N. obtusifolia*, but was not validly published as he failed to provide a Latin description or diagnosis, in violation of Art. 39.1 of the Code (MCNEILL et al., 2012). *Noronhia martiniana* differs from *N. obtusifolia* by having large leaf blades (up to 16 cm vs. 10 cm long), long petioles (up to 28 mm vs. 20 mm long), small flowers (less than 6 mm long vs. more than 6 mm), and stamens that are oblong (vs. ovate). Molecular studies also placed these species in two different clades (HONG-WA & BESNARD, 2014). Therefore, the taxon initially recognized by PERRIER DE LA BATHIE (1949) is described here as a new species using a different epithet and a type specimen that is more widely distributed in various herbaria. The new species can be recognized by its long, non-woody petioles, oblanceolate to obovate leaf blades, white, campanulate flowers lacking a corona, and ovoid to oblong fruits.

Additional specimens examined

MADAGASCAR. **Prov. Antsiranana:** Ampanavoana, Vinanivao, Masoala PN, 15°41'33"S 50°13'30"E, 12-100 m, 15.IX.1996, Bernard 334 (K, MO, P, TAN); Masoala PN, Beankoraka, 15°57'S 50°13'E, 14.XI.1994, Rahajasoa et al. 880 (K, MO, TAN). **Prov. Fianarantsoa:** Farafangana, Mahabo Mananivo, Nosiala, forêt d'Agnalazaha, 23°11'08"S 47°42'48"E, 50 m, 5.III.2009, Hong-Wa et al. 619 (MO, P, TAN); ibid. loc., 5.III. 2009, Hong-Wa et al. 620 (MO, P, TAN); Nosy Varika, Ambahy, 20°48'07"S 48°28'56"E, 9 m, 15.II.2004, Rabehevitra 881 (K, MO, P, TEF); Mahabo-Mananivo, forêt de Mahabo, 23°10'38"S 47°42'35"E, 20 m, 8.XI.2001, Rabenantoandro 717 (G, MO, P, TEF); Nosy Varika., Ampasimaneva, forêt d'Ambolo, 20°43'59"S 48°27'09"E, 13 m, 15.VI.2004, Razakamalala et al. 1452 (G, K, MO, P, TEF); Ambahy Forest, c. 20 km S of Nosy Varika, 20°46'09"S 48°28'46"E, 3 m, 22.IV.2004, Rogers et al. 300 (MO, P, TAN). **Prov. Toamasina:** Forêt de Tampina, [18°31'S 49°17'E], X.1923, Louvel 126 (P); Mahanoro, Ambalavontaka, forêt de Nankinana, 20°22'21"S 48°33'15"E, 41 m, 8.XI.2003, Rabehevitra et al. 685 (G, K, MO, P, TEF); Sainte Marie, Lokintsy, forêt d'Ambohidena,



16°51'11"S 49°57'10"E, 1.VI.2004, *Rabehevitra et al.* 1179 (K, MO, P, TEF); Fénérive-Est, Tampolo STF, 17°16'52"S 49°24'44"E, 0-150 m, 24.I.1995, *Raholivelolo* 212 (MO); Mahanoro, Ambalavontaka, forêt d'Antaimby, 20°22'22"S 48°33'56"E, 13 m, 17.IV.2004, *Ranaivojaona et al.* 587 (K, MO, P, TEF); Mahanoro, Ambalavontaka, forêt d'Ampanamara, 20°22'48"S 48°33'23"E, 13 m, 12.VI.2004, *Ranaivojaona et al.* 707 (MO, P, TEF); Ambalavontaka, Nankinana Forest, c. 20 km N of Nosy Varika, 20°21'34"S 48°36'20"E, 10 m, 18.IV.2004, *Rogers et al.* 195 (MO, P, TAN). **Prov. Toliara:** Fort-Dauphin, Sainte Luce, 24°46'47"S 47°10'06"E, 14 m, 17.X.2012, *Ramanjanahary et al.* 784 (MO, P, TAN); Iaboakoho, Ambanihazo, forêt d'Analandrasambo, 24°40'55"S 47°11'56"E, 22 m, 22.X.2012, *Randrianaivo et al.* 2036 (MO, P, TAN).