

Zeitschrift: Boissiera : mémoires de botanique systématique

Herausgeber: Conservatoire et Jardin Botaniques de la Ville de Genève

Band: 70 (2016)

Artikel: A taxonomic revision of the genus Noronhia Stadtm. ex Thouars (Oleaceae) in Madagascar and the Comoro Islands

Autor: Hong-Wa, Cynthia

Kapitel: 57: Noronhia oblanceolata

DOI: <https://doi.org/10.5169/seals-1036126>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 11.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

57. *Noronhia oblanceolata* H. Perrier in Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 2: 284. 1949 (Fig. 40B).

Lectotypus (designated here): **MADAGASCAR. Prov. Antsiranana:** près de la cime du Mt Mahabenofo (N), bassin de la Loky, [13°16'S 49°26'E], 1932, *Perrier de la Bâthie* 2316 (P [P00701217]!). **Syntypus:** **MADAGASCAR. Prov. Mahajanga:** Boeny, 7e Réserve [Ankarafantsika PN], chemin Ankorika-Sainte Marie, [16°09'S 47°04'E], 150 m, *Service Forestier* 82 (P [P00413237]!).

Description Shrubs to small trees to 6 m tall, trunk to 11 cm diameter; young twigs cylindrical, 0.5-1.2 mm diameter, glabrous; bark medium gray, smooth to slightly rugose. Leaves opposite, verticillate at the tips of the branches, semi-deciduous; bud scales persistent; blades dark green above, lighter below, oblanceolate, 1.5-5 × 0.5-1.5 cm, coriaceous, glabrous, domatia absent, base acute to attenuate, margin flat to slightly revolute, apex retuse, midrib sunken above, slightly raised below, secondary veins inconspicuous, 5-11 per side, 2-6 mm apart, looping 0.4-1.5 mm from the margin; petiole yellow to brownish, 2-5 × 0.4-1.1 mm, not woody, glabrous. Thyrse fasciculate, pauciflorous, compact to diffuse; peduncle 2-5 mm long, moderately pubescent; pedicel 1.5-4 mm long, moderately pubescent; calyx moderately pubescent outside, glabrous inside, lobes triangular, 1.5-2.2 × 0.9-1.2 mm; corolla white, oblong, 3.3-5.5 mm long, glabrous on both sides, the tube 1.5-2 mm long, lobes deltate, apex obtuse; corona absent; stamens 1.6-2.1 mm long, anthers ovate to lanceolate, 1.3-1.6 mm long; pistil 1.4-2 mm long, stigma slightly bilobed. Fruiting pedicel 4-11 × 1.1-1.9 mm; young fruits green, dark red when mature, globose to subglobose, 11.5-23 × 9.5-17 mm, smooth, apex apiculate to rostellate, the rostellum circular, apiculate; dry pericarp 0.7-2.2 mm thick; endocarp woody; seed 9-12.5 × 7-10 mm.

Distribution, ecology and phenology *Noronhia oblanceolata* occurs in low-elevation dry forests in the north (Analamerina to Daraina) and the west (Ankarafantsika) (Fig. 36). It produces flowers and fruits from October to March.

Conservation status The assessment used 12 collections representing 12 localities and resulted in an EOO of 12,991 km², an AOO of 40 km², and eight subpopulations representing seven locations, of which five occur within protected areas (Analamerana, Ankarafantsika, Ankarana, and Loky-Manambato). Because of forest exploitation, land conversion and artisanal mining, habitat degradation and loss will likely persist outside of protected areas, although a reduction in EOO is not expected in the near future. Therefore, *N. oblanceolata* is assigned a preliminary status of “Vulnerable” [VU B1ab(ii,iii,iv,v) +2ab(ii,iii,iv,v)].

Notes *Noronhia ob lanceolata* can be recognized by its oblanceolate leaf blades that are verticillate at the tips of the branches and its globose to subglobose, apiculate fruits. It differs from *N. maculata* by its non-woody (vs. woody) petioles, oblanceolate (vs. obovate) leaf blades, oblong (vs. subrotate), white (vs. pale green) corolla lacking a corona, and globose to subglobose (vs. ovoid) fruits.

PERRIER DE LA BATHIE (1949, 1952) used *Service Forestier* 82 (from Ankrafantsika in the west) as the flower-bearing syntype and *Perrier de la Bâthie* 2316 (from Analamera in the north) as the fruit-bearing syntype. The flowers of *Service Forestier* 82 are clustered into diffuse inflorescences and were described as subcampanulate, the corolla being wider than long, with a thick corona as long as the corolla tube. Flowers of specimens recently collected in the north and corresponding to *Perrier de la Bâthie* 2316 are clustered into compact inflorescences, are oblong, have only a short corolla tube, and lack a corona. The northern and western plants may thus represent two different entities. However, because the western entity is known only from a single specimen that may simply represent an extreme in variation, it is preferable to await more collections from this area to determine their respective identities. In the meantime, a broad definition of *N. ob lanceolata* is retained but in order to avoid ambiguity regarding the use of the name *N. ob lanceolata* and to associate it permanently with populations from the north, *Perrier de la Bâthie* 2316 has been selected here as the lectotype.

Additional specimens examined

MADAGASCAR. Prov. Antsiranana: Daraina, forêt d'Antsahabe, 13°12'55"S 49°33'00"E, 750 m, 3.XII.2004, Gautier 4820 (Daraina, G, MO, P, TEF); Analamera, [12°42'S 49°20'E], 50-400 m, 1.I.1938, Humbert 19137 (P); Ambohipiraka, [13°11'S 49°06'E], 1.X.1932, *Perrier de la Bâthie* 18757 (P); 3 km E d'Ambilobe, 13°11'57"S 49°04'44"E, 72 m, 23.I.2007, Rakotonandrasana et al. 1069 (CNARP, G, MO, P, TAN); Daraina, forêt d'Antsahabe, 13°10'51"S 49°33'13"E, 415 m, 1.V.2004, Ranirison 756 (Daraina, G, K, MO, P, TEF); Daraina, forêt d'Ambohitsitondroina, 13°07'56"S 49°27'16"E, 230 m, 6.I.2006, Ranirison & Nusbaumer 1053 (Daraina, G, K, MO, P, TEF); Daraina, forêt d'Antsahabe, 13°12'32"S 49°33'15"E, 560 m, 3.XI.2005, Razafitsalama et al. 757 (CNARP, MO, P, TAN); *ibid. loc.*, 13°12'32"S 49°33'15"E, 560 m, 3.XI.2005, Razafitsalama et al. 760 (CNARP, MO, P, TAN); *ibid. loc.*, 13°12'32"S 49°33'15"E, 560 m, 3.XI.2005, Razafitsalama et al. 765 (CNARP, MO, P, TAN); Ankarana RS, [12°49'S 49°01'E], 50-409 m, *Service Forestier* 10421 (P, TEF).