

**Zeitschrift:** Boissiera : mémoires de botanique systématique  
**Herausgeber:** Conservatoire et Jardin Botaniques de la Ville de Genève  
**Band:** 70 (2016)

**Artikel:** A taxonomic revision of the genus *Noronhia* Stadtm. ex Thouars (Oleaceae) in Madagascar and the Comoro Islands  
**Autor:** Hong-Wa, Cynthia  
**Kapitel:** 58: *Noronhia obtusifolia*  
**DOI:** <https://doi.org/10.5169/seals-1036126>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 11.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

**58. *Noronhia obtusifolia*** (Lam.) Hong-Wa & Besnard in Mol. Phylogenet. Evol. 67: 376. 2013 (Fig. 42).

- ≡ *Olea obtusifolia* Lam., Tabl. Encycl. Méthod. 1: 28. 1791.
- ≡ *Linociera obtusifolia* (Lam.) H. Perrier in Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 2: 279. 1949.
- ≡ *Chionanthus obtusifolius* (Lam.) Stearn in Bot. J. Linn. Soc. 80: 203. 1980.

**Lectotypus** (designated here): **MADAGASCAR**: *sine loc.*, s.d., *Commerson s.n.* (P-JU n° 4900 [P00657336] image seen; isolecto-: G [G00007660] image seen, P [P00413214]!, P-LA [P00356887] image seen).

– *Linociera obtusifolia* var. *thouarsii* H. Perrier in Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 2: 279. 1949 [nom. inval.] ≡ *Noronhia obtusifolia* var. *thouarsii* (H. Perrier) Hong-Wa & Besnard in Mol. Phylogenet. Evol. 67: 376. 2013 [nom. inval.].

**Description** *Trees* to 20 m tall, trunk to 25 cm diameter; young twigs cylindrical, 1-2.5 mm diameter, glabrous; bark dark to medium gray, smooth to slightly rugose. *Leaves* opposite, persistent; bud scales rarely persistent; blades dark green above, lighter below, obovate, 3.5-10.5 × 1.5-4 cm, coriaceous, glabrous, domatia absent, base attenuate, margin slightly revolute, apex obtuse to rounded or obcordate, midrib slightly sunken above, raised below, secondary veins distinct to barely visible, 7-13 per side, 5-10 mm apart, looping 1-2 mm from the margin; petiole yellow to brownish, 6-20 × 0.5-2.3 mm, not woody, sometimes only partially, glabrous. *Thyraxes* geminate, multiflorous, compact; peduncle 8-14 mm long, glabrous; pedicel 1.5-4 mm long, glabrous; calyx glabrous on both sides, lobes triangular, 1-1.3 × 0.8-1.2 mm; corolla white, campanulate, petals almost free, 5-7 mm long, glabrous on both sides, the tube 1-4 mm long, lobes ovate, apex obtuse; corona absent; stamens 2-2.8 mm long, anthers ovate, apiculate, 1.5-1.8 mm long; pistil 1.5-2.5 mm long, stigma bilobed. *Fruiting* pedicel 1-1.7 × 1.3-1.6 mm; young fruits green, reddish black when mature, ovoid to subglobose, 16-21 × 6-16.5 mm, smooth to ribbed, apex flat to rostellate, with the persistent style; dry pericarp 0.2-1.2 mm thick; endocarp woody; seed 14-15 × 5-12 mm.

**Distribution, ecology and phenology** *Noronhia obtusifolia* occurs in low- to high-elevation humid forests in eastern Madagascar, from around Anjanaharibe-Sud in the north to around Fort-Dauphin in the south (Fig. 36). It produces flowers and fruits all year.

**Conservation status** Based on 52 collections representing 43 localities, the analyses indicated an EOO of 97,820 km<sup>2</sup>, an AOO of 132 km<sup>2</sup>, and 26 subpopulations representing 26 locations, of which 12 occur within protected areas (Agnalazaha, Ambohidena, Andohahela, Anjanaharibe-Sud, Corridor Ankeniheny-Zahamena, Makira, Mandena, Marojejy,

Ste Luce, Tampolo, Tsitongambarika, and Vohibola). With a large EOO and an actual AOO likely exceeding 2,000 km<sup>2</sup>, *N. obtusifolia* is assigned a preliminary status of “Least Concern”.

**Notes** *Noronhia obtusifolia* can be recognized by its elliptic to narrowly or very widely obovate leaf blades terminated by an obtuse to obcordate apex, compact inflorescences, white flowers with barely connate petals, and ovoid, almost elongate, smooth to ribbed fruits. The variation in the shape of the leaf blades and fruits as well as fruit ornamentation may be sufficient to distinguish different entities within this species. Indeed, PERRIER DE LA BATHIE (1949, 1952) recognized three varieties, of which two were, however, invalidly published due to a lack of Latin diagnosis (Art. 39.1; McNEILL et al., 2012). Consequently, the combinations *N. obtusifolia* var. *minoriflora* (H. Perrier) Hong-Wa & Besnard and *N. obtusifolia* var. *thouarsii* (H. Perrier) Hong-Wa & Besnard are also invalid as they were based on these names.

The taxon referred to as *N. obtusifolia* var. *minoriflora* is here recognized as distinct at the species level and described as new (see p. 179) using the name *N. martiniana*. By contrast, *N. obtusifolia* var. *thouarsii*, based on a very poor material, is better treated as *N. obtusifolia* (without recognition at an infraspecific level) until additional specimens become available. The collection in P-JUS is here designated as lectotype as it is more complete than the material in P-LAM. Furthermore, the latter collection bears the indication “ex-Juss” meaning that Lamarck had received the collection from Jussieu. LAMARCK (1791) indicated Mauritius as the origin of the type by error; it is in fact a Commerson collection from Madagascar.

#### Additional specimens examined

**MADAGASCAR. Prov. Antsiranana:** Ambatosoratra, [14°34'S 49°44'E], 400 m, 5.I.1949, *Cours 3292* (MO, P); Mt Ambatosoratra, [14°25'48"S 49°43'48"E], 1500-1540 m, 4-8.I.1949, *Humbert & Cours 22941* (P); Anjanaharibe-Sud RS, 14°48'15"S 49°26'45"E, 1000-1100 m, 6.VIII.1997, *McPherson 17254* (K, MO). **Prov. Fianarantsoa:** Nosy Varika, Ambahy, 20°48'07"S 48°28'56"E, 22 m, 13.XI.2003, *Rabehevitra et al. 741* (K, MO, P, TEF). **Prov. Mahajanga:** Befandriana-Nord, Matsoandakana, Andranomena, forêt d'Anjiahe, 15°08'10"S 49°21'08"E, 1085 m, 11.II.2008, *Ravelonarivo et al. 2775* (G, MO, P, TAN). **Prov. Toamasina:** Maroantsetra, Ambinanitelo, Marovovonana, Ankirindro forest, 15°17'34"S 49°32'46"E, 690 m, 13.I.2003, *Antilahimena 1711* (G, MO, P); Antsirabe Sahatany, Anjiahely, 15°24'45"S 49°22'29"E, 1201 m, 19.VI.2004, *Antilahimena 2529* (MO); Ambinanitelo, Marovovonana, 15°19'03"S 49°24'06"E, 800 m, 4.IX.2004, *Antilahimena 2767* (MO); Mananara, Vanono, Lohan'i Ambitsy forest, 15°53'27"S 49°28'57"E, 846 m, 26.VII.2007, *Antilahimena 5706* (MO, P, TAN); Mahanoro, Ambodibonara, Ambalavontaka, forêt de Nankinana, 20°22'21"S 48°33'15"E, 16 m, 12.II.2004, *Rabehevitra 801* (MO, P, TEF); Brickaville, Ambila-Lemaitso, 18°49'S 49°08'E, 0-5 m, 11.XI.1999, *A. Randrianasolo & Ranaivojoana 650* (G, MO, P); Brickaville, Maroseranana, Ambodilendemy, 18°24'49"S 48°47'32"E, 1175 m, 24.III.2011, *Ravelonarivo et al. 3848* (G, MO, P, TAN); Sainte Marie, Lokintsy, Sahasifotra, forêt d'Ambohidena, 16°50'25"S 49°57'09"E, 10 m, 19.II.2004, *Razakamalala 943* (K, MO, P, TEF); Ambila-Lemaitso, [18°49'S 49°08'E], 30 m, 8-10.VIII.1957, *Service Forestier 18067* (MO). **Prov. Toliara:**

Fort-Dauphin, Mt Vohimavo, 24°19'32"S 47°08'11"E, 836-837 m, 4.XII.2001, *Hong-Wa* 29 (G, MO, P, TAN); Ampasy Nampoana, forêt de Mandena, 24°57'10"S 47°00'10"E, 9 m, 12.II.2009, *Hong-Wa* 599 (MO, P, TAN); Ambatoabo, forêt d'Ankoba, 24°46'44"S 46°43'17"E, 660 m, 19.VII.2011, *Randrianaivo et al.* 1888 (G, MO, P, TAN); Iabakoho, Antsofso, forêt de Bemangidy, 24°35'33"S 47°12'52"E, 22.V.2006, *Randriatafika et al.* 655 (G, MO, P, TEF); *ibid. loc.*, 24°34'10"S 47°12'01"E, 651 m, 2008, *Randriatafika* 903 (G, MO, P, TEF); Ampasy Nampoana, forêt de Mandena, 24°57'26"S 47°01'34"E, 90 m, 13-15.VI.1999, *Ratovoson et al.* 106 (MO, P, TAN); Iaboko, Antsofso Avaratra, 24°33'42"S 47°12'02"E, 662 m, 11.XII.2007, *Razakamalala et al.* 3989 (G, MO, P, TAN); Fort-Dauphin, Manafiafy, 24°47'S 47°10'E, 15 m, 13.XI.1990, *Schatz et al.* 3009 (G, MO, TAN); Vataza, au SE d'Imonty, [24°36'S 46°42'E], 16.I.1963, *Service Forestier* 22458 (G, MO). **Sine loc.**: s.d., *du Petit-Thouars* s.n. (P).



**Fig. 42.** *Noronhia obtusifolia* (Lam.) Hong-Wa & Besnard [Schatz 3009].

Photo: G. Schatz