

Zeitschrift:	Boissiera : mémoires de botanique systématique
Herausgeber:	Conservatoire et Jardin Botaniques de la Ville de Genève
Band:	70 (2016)
Artikel:	A taxonomic revision of the genus Noronhia Stadtm. ex Thouars (Oleaceae) in Madagascar and the Comoro Islands
Autor:	Hong-Wa, Cynthia
Kapitel:	60: Noronhia orientalis
DOI:	https://doi.org/10.5169/seals-1036126

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 11.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

60. *Noronhia orientalis* Hong-Wa, spec. nova (Fig. 43B, 44).

Typus: MADAGASCAR. Prov. Antsiranana: SAVA, Sambava, Anjangoveratra, Ambavala, forêt d'Antsahakasaka, 2 km au NO du village Ambavala, 14°05'39"S 50°02'42"E, 146 m, 16.II.2006, Andrianjafy et al.1599 (holo-: MO-6615571!; iso-: G [G00341629]!, P!, TAN!).

Diagnosis *Noronhia orientalis* Hong-Wa can be distinguished from other members of the genus by its terminally verticillate leaves, its obovate leaf blades, with very conspicuous venation and its compact inflorescences with white flowers.

Description Trees to 10 m tall, trunk to 10 cm diameter; young twigs cylindrical, 1.5-3.1 mm diameter, glabrous; bark dark gray, smooth to rugose, sometimes lenticellate. Leaves opposite, sometimes verticillate at the tips of the branches, persistent; bud scales persistent; blades medium green above, lighter below, oblong to obovate, 7-15 × 2.5-5 cm, subcoriaceous, glabrous, domatia casual to abundant, base acute to attenuate, sometimes truncate, margin flat, slightly revolute, apex retuse to shortly cuspidate, the cusp 1-6 mm long, midrib sunken above, distinctly raised below, secondary veins conspicuous, 10-15 per side, 8-16 mm apart, looping 2-5 mm from the margin; petiole light gray to brownish, 6-22 × 1.5-3 mm, entirely woody, glabrous. Thyrses geminate to fasciculate, pauciflorous, compact; peduncle 5-10 mm long, glabrous; pedicel 3-4 mm long, glabrous; calyx glabrous on both sides, lobes triangular, 0.7-1.2 × 1-1.2 mm; corolla white, cupuliform, 3-4.5 mm long, glabrous on both sides, the tube 1.5-3 mm long, lobes ovate, apex acute; corona present, 1.6-2 mm long, undivided; stamens 1.5-2 mm long, anthers oblong, 0.7-1.3 mm long; pistil 2-2.3 mm long, slightly bilobed. Fruiting pedicel 8-10 × 3-3.5 mm; young fruits green, unseen mature, ovoid to oblong, 18-21 × 14-15 mm, surface smooth, sometimes glaucous and slightly rugose, apex flat; dry pericarp 1mm thick; endocarp woody; seed 10-15 × 6-10 mm.

Etymology The specific epithet refers to its strictly eastern distribution, specifically northeastern Madagascar.

Distribution, ecology and phenology *Noronhia orientalis* occurs in littoral to low-elevation humid forests in the northeast, from Antsirabe Nord to the Masoala peninsula (Fig. 36). It has been collected in flowers and fruits in February.

Conservation status *Noronhia orientalis* is currently known from seven collections representing seven localities. With an EOO of 4,012 km², an AOO of 28 km², and five subpopulations representing five locations, of which four occur within protected areas (Makirovana-Tsi-

homanaomby, Marojejy, and Masoala), *N. orientalis* is assigned a preliminary status of “Endangered” [EN B1ab(ii,iii,v)+2ab(ii,iii,v)] because habitat loss as well as continuing decline in habitat quality and number of mature individuals is projected both outside and inside of protected areas due to wood harvesting, land conversion and illicit exploitation.

Notes *Noronhia orientalis* and *N. retusifolia* Hong-Wa are similar in appearance but can be distinguished from each other by the shape of their leaves (oblong to obovate in *N. orientalis* vs. oblanceolate to obovate in *N. retusifolia*), the length of their petioles (up to 22 mm vs. less than 11 mm), and the surface (smooth to sometimes slightly rugose vs. rugose) and apex (flat vs. bluntly pointed) of their fruits. Features such as terminally verticillate leaves, obovate leaf blades, with very conspicuous venation and compact inflorescences with white flowers characterize this species.

Paratypi **MADAGASCAR.** **Prov. Antsiranana:** Sambava, Anjangeratra, forêt d'Andalibe, 14°07'35"S 50°03'07"E, 115 m, 23.III.2009, Hong-Wa 636 (MO, P, TAN); Vohémar, Antsirabe Nord, Andravinambo, forêt d'Antsaolatra, 14°06'24"S 49°58'39"E, 328 m, 31.I.2014, Martial et al. 432 (DOV, MO, P, TAN); Sambava, Anjaggeratra, Ambavala, forêt de Tsihomanaomby, 14°06'36"S 50°02'41"E, 330 m, 15.II.2014, Martial et al. 498 (MO, P, TAN); Masoala PN, Beankoraka, 15°57'S 50°13'E, 14.XI.1994, Rahajaso et al. 888 (MO, TAN); Sambava, Andratamarina, 14°25'48"S 49°50'45"E, 451 m, 19.X.2010, Ravelonarivo & Raharivelo 3516 (MO, P, TAN). **Prov. Toamasina:** Masoala PN, Antalavia, 15°47'S 50°02'E, 0 m, 27.XI.1994, Rahajaso et al. 1078 (MO).




Fig. 44.

Noronhia orientalis Hong-Wa.

A. Flowering branch; **B.** Flower; **C.** Lateral view of corona; **D.** Inner side of corolla; **E.** Pistil.

[Andriananjafy 1599, TAN] Drawings: R.L. Andriamiarisoa