

Zeitschrift:	Boissiera : mémoires de botanique systématique
Herausgeber:	Conservatoire et Jardin Botaniques de la Ville de Genève
Band:	70 (2016)
Artikel:	A taxonomic revision of the genus Noronhia Stadtm. ex Thouars (Oleaceae) in Madagascar and the Comoro Islands
Autor:	Hong-Wa, Cynthia
Kapitel:	61: Noronhia ovalifolia
DOI:	https://doi.org/10.5169/seals-1036126

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 08.10.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

61. *Noronhia ovalifolia* H. Perrier in Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 2 : 301. 1949 (Fig. 45A).

Typus: MADAGASCAR. Prov. Toliara: bassin de réception de la Mananara, affluent du Mandrare, pentes occidentales des montagnes entre l'Andohahela et l'Elakelaka, Mont Apiky au-dessus de Mahamavo, 800-900 m, I-II.1934, Humbert 13822 (holo- : P [P00418097]!; iso- : K [K000233183] image seen, P [P00418095, P00418096]!, WAG [WAG0002495] image seen).

Description Shrubs to 3 m tall, trunk to 3 cm diameter; young twigs cylindrical, 0.8-1.6 mm diameter, glabrous; bark medium to light gray, smooth. Leaves opposite, persistent; bud scales persistent; blades dark green above, yellowish below, ovate, 3.5-9.5 × 2-4.5 cm, coriaceous, glabrous, domatia absent, base rounded, margin slightly revolute, apex acuminate, the acumen 1-9 mm long, midrib flat to slightly raised above, flat below, secondary veins conspicuous mostly below, 7-10 per side, 4-13 mm apart, looping 2-5.5 mm from the margin; petiole medium to light gray, 3-7 × 1-2 mm, entirely woody, glabrous. Flowers solitary; pedicel 3-12 mm long, glabrous; calyx glabrous on both sides, lobes deltate to obovate, 1.5-2 × 2.1-2.6 mm; corolla pink, sometimes tinged white outside, cupuliform, 6.5-10 mm long, glabrous on both sides, the tube 4.5-8.5 mm long, lobes widely ovate, apex obtuse; corona present, 2.5-3 mm long, slightly lobed; stamens 2.8-3 mm long, anthers obovate, 1.8-2.2 mm long; pistil 2.5-2.8 mm long, stigma capitate. Fruiting pedicel 12-16 × 1-1.5 mm; young fruits green, reddish black when mature, subglobose to ovoid, 19-19.5 × 15-17.5 mm, surface smooth, apex flat to apiculate; dry pericarp 0.6-0.9 mm thick; endocarp woody; seed 16 × 10.5 mm.

Distribution, ecology and phenology *Noronhia ovalifolia* occurs in low- to mid-elevation transitional forests on basement rocks and alluvial deposits in and around Andohahela PN in the south (Fig. 36). It produces flowers and fruits from October to March.

Conservation status The assessment included eight collections representing eight localities and resulted in an EOO of 93 km², an AOO of 20 km², and three subpopulations representing three locations, of which two occur within a protected area (Andohahela). The subpopulation outside of the protected area occurs in an area that is now heavily degraded and has been converted into agriculture fields and pastureland, and will therefore likely experience continuing decline, potentially leading to extirpation. Therefore, *N. ovalifolia* is assigned a preliminary status of “Endangered” [EN B1ab(i, ii, iii, iv, v)+2ab(i, ii, iii, iv, v)].

Notes *Noronhia ovalifolia* can be recognized by its short petioles, ovate leaf blades with light-colored venation, and solitary flowers with a pink to orangish corolla. It differs from *N. boivinii* and *N. mangorensis* by its ovate (vs. oblong to elliptic in *N. boivinii* and oblong to ovate in *N. mangorensis*), acuminate (vs. cuspidate to mucronate and cuspidate, respectively) leaf blades, solitary (vs. fasciculate) flowers, and subglobose (vs. ovoid) fruits.

Additional specimens examined **MADAGASCAR.** **Prov. Toliara:** Fort-Dauphin, Ambatoabo, Evasia, Andohahela PN, parcelle 1, 24°47'10"S 46°43'33"E, 681 m, 16.I.2007, *Andriamihajarivo et al.* 1097 (MO, P, TAN); Ankoba, near E edge of Andohahela PN Park, 24°46'46"S 46°43'11"E, 600 m, 22.I.2008, Lowry *et al.* 6955 (MO, P); *ibid. loc.*, 24°47'34"S 46°41'55"E, 200 m, 24.I.2008, Lowry *et al.* 6977 (MO, P); Fort-Dauphin, Ambatoabo, forêt d'Ankoba, 24°46'56"S 46°42'14"E, 228 m, 1.III.2008, Randrianaivo *et al.* 1548 (G, MO, P, TAN); *ibid. loc.*, 24°47'12"S 46°42'25"E, 300 m, 25.II.2009, Randrianaivo *et al.* 1760 (MO, P, TAN); Amboasary, Behara, [24°36'S 46°42'E], 11.X.1958, Réserves Naturelles 10060 (P, TEF); Italy (baie de Ranofotsy), [25°09'00"S 46°45'30"E], 0-25 m, 10.I.1963, Service Forestier 22371 (G, P, TEF).




Fig. 45. Photographs of *Noronhia* Stadtm. ex Thouars.

A. *Noronhia ovalifolia* H. Perrier [Lowry 6977]; **B.** *Noronhia peruviana* (Knobl.) H. Perrier [Bolliger 223].

Photos: taken by respective collectors