

**Zeitschrift:** Boissiera : mémoires de botanique systématique  
**Herausgeber:** Conservatoire et Jardin Botaniques de la Ville de Genève  
**Band:** 70 (2016)

**Artikel:** A taxonomic revision of the genus *Noronhia* Stadtm. ex Thouars (Oleaceae) in Madagascar and the Comoro Islands  
**Autor:** Hong-Wa, Cynthia  
**Kapitel:** 62: *Noronhia patricei*  
**DOI:** <https://doi.org/10.5169/seals-1036126>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

**Download PDF:** 16.05.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

**62. *Noronhia patricei*** Hong-Wa, *spec. nova* (Fig. 46).

**Typus:** MADAGASCAR. **Prov. Toamasina:** Analanjirofo, Ambanizana, along path between Ambanizana-Andranobe, 15°38'29"S 49°57'54"E, 7 m, 25.X.2004, *Antilahimena* 2959 (holo-: MO-6615553!; iso-: G [G00341616]!, K!, P [P03532773]!, TAN!).


**Diagnosis** *Noronhia patricei* Hong-Wa can be distinguished from its congeners by its three-verticillate, large leaves, its fasciculate infructescences, and its large fruits characterized by a thick pericarp.

**Description** Trees to 11 m tall, trunk to 16 cm diameter; young twigs subquadrangular to cylindrical, 5.5-8.6 mm diameter, glabrous; bark medium gray, rugose. Leaves verticillate, persistent; bud scales persistent; blades medium green above, lighter below, lanceolate, 26-36 × 4.5-9.5 cm, very coriaceous, glabrous, domatia absent, base acute, margin flat, apex acuminate, the acumen 5-12 mm long, midrib deeply sunken above, distinctly raised below, secondary veins conspicuous, 14-20 per side, 19-40 mm apart, looping 2-7 mm from the margin; petiole medium to dark gray, 16-20 × 4-6.5 mm, entirely woody, glabrous. Flowers unseen, but infructescence thyrsoid, fasciculate. Fruiting pedicel 5-11 × 3.5-5 mm; young fruits green, yellow when mature, ovoid to oblong, 25-38 × 19.5-24 mm, surface smooth, sometimes covered with white dots, apex flat to bluntly pointed; dry pericarp 2.2-3 mm thick; endocarp woody; seed 15 × 20 mm.

**Etymology** This species is dedicated to Patrice Antilahimena, a botanist at the Missouri Botanical Garden in Madagascar, for collecting the type specimen and for keeping an eye out for *Noronhia* during his extensive fieldwork.

**Distribution, ecology and phenology** *Noronhia patricei* occurs in low-elevation humid forests and basement rocks and unconsolidated sands in the Masoala region in northeastern Madagascar (Fig. 36). It fruits from October to June.

**Conservation status** *Noronhia patricei* is currently known only from five collections representing five localities. The assessment resulted in an EOO of 1,023 km<sup>2</sup>, an AOO of 20 km<sup>2</sup>, and four subpopulations representing four locations, of which two occur within a protected area (Masoala). With decline in habitat quality, habitat loss and reduction of the number of mature individuals likely to continue outside of the protected area as a result of wood harvesting and land conversion, the small EOO and AOO as well as the small number of locations, *N. patricei* is assigned a preliminary status of "Endangered" [EN B1ab(i,ii,iii,iv,v)+2ab(i,ii,iii,iv,v)].



**Fig. 46.**  
*Noronhia patricei* Hong-Wa.  
**A.** Branch; **B.** Fruit with pedicel; **C.** Fruit.

[Antilahimena 2959, TAN] Drawings: R.L. Andriamiarisoa

**Notes** *Noronhia patricei* is similar to *N. crassiramosa* but can be distinguished by its entirely verticillate (vs. verticillate only at the tips of the branches), lanceolate (vs. oblong) leaves, and its ovoid to oblong (vs. subglobose) fruits with a flat (vs. apiculate) apex. Its three-verticillate, large leaves, fasciculate infructescences, and large fruits characterized by a thick pericarp allow for the unambiguous identification of this species.

**Paratypes** **MADAGASCAR.** **Prov. Antsiranana:** Maroantsetra, Anjahana, Ambanizana, 15°37'03"S 49°58'36"E, 200 m, 11.VI.2002, *Antilahimena 1090* (G, MO, P); Antalaha, Ampanavoana, forêt de Faho, [15°41'S 50°21'E], 24.IV.1951, *Service Forestier 87R140* (TEF). **Prov. Toamasina:** Ambatondradama (env. 10 km au NE d'Ankovona), Navana, 15°17'13"S 50°01'10"E, 500 m, 27.IX-6.X.1997, *Ralimanana et al. 111* (MO, P, TAN); Maroantsetra, Anjahana, Ambanizana, Tampolo (Masoala), 15°43'45"S 49°57'38"E, 10 m, 19.X.2001, *Randrianaivo 710* (MO, P, TAN).