

Zeitschrift: Boissiera : mémoires de botanique systématique
Herausgeber: Conservatoire et Jardin Botaniques de la Ville de Genève
Band: 70 (2016)

Artikel: A taxonomic revision of the genus *Noronhia* Stadtm. ex Thouars (Oleaceae) in Madagascar and the Comoro Islands
Autor: Hong-Wa, Cynthia
Kapitel: 67: *Noronhia populifolia*
DOI: <https://doi.org/10.5169/seals-1036126>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 11.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

67. *Noronhia populifolia* H. Perrier in Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 2 : 298. 1949 (Fig. 51).

Typus: MADAGASCAR. **Prov. Antsiranana:** Sambirano, bord de l'Andranomalaza, [14°19'00"S 48°17'30"E], VIII.1907, *Perrier de la Bâthie* 8825 (holo-: P [P00418101]!).

Description *Trees* to 16 m tall; young twigs cylindrical, 2-2.5 mm diameter, glabrous; bark medium to light gray, smooth to slightly rugose. *Leaves* opposite, persistent; bud scales persistent; blades light green, oblong to ovate, 10.5-22 × 5.5-8 cm, subcoriaceous, glabrous, domatia absent, base rounded, margin flat to undulate, apex acuminate, the acumen 13-36 mm long, midrib slightly sunken above, raised below, secondary veins conspicuous, 9-14 per side, 13-16 mm apart, looping 3-4 mm from the margin; petiole yellowish to brown, 10-16 × 1.9-2.6 mm, not woody, glabrous. *Thyraxes* geminate to fasciculate, pauciflorous, compact; peduncle 8-11 mm long, moderately pubescent; pedicel 4-8 mm long, moderately pubescent; calyx moderately pubescent outside, glabrous inside, lobes triangular, 1-2.5 × 1.4-2 mm; corolla pink, cupuliform, 5-5.5 mm long, glabrous on both sides, the tube 2.5-3 mm long, lobes widely ovate; corona present, 1.7-2 mm long, undivided; stamens 2-2.5 mm long, stamens widely obovate, 1.5-1.6 mm long; pistil 2.2-2.5 mm long, stigma capitate. *Fruiting* pedicel 5 × 3 mm; young fruits green, blackish when mature, oblong to subglobose, 28.5-35 × 24-30 mm, surface verrucose, apex apiculate; dry pericarp 1.7-2.2 mm thick; endocarp woody; seed 25 × 23 mm.

Distribution, ecology and phenology *Noronhia populifolia* occurs in mid-elevation humid forests and sandstones in the Sambirano region in the northwest (Fig. 49). It has been collected in flowers and fruits in October.

Conservation status With just a single collection made in 1907 from an area that is not protected and that has been gradually degraded as a result of agriculture and wood harvesting, *N. populifolia* is assigned a preliminary status of "Critically Endangered", and could possibly be extinct [CR(PE) B2(iii)].

Notes *Noronhia populifolia* can be recognized by its non-woody petiole, broadly ovate leaf blades with a long apex, somewhat compact inflorescences bearing pink flowers, and large, verrucose fruits with a thick pericarp. It differs from *N. pervilleana* by its non-woody (vs. woody) petioles, cupuliform (vs. rotate), pink (vs. white) flowers, and oblong to subglobose (vs. ovoid), verrucose (vs. smooth to punctate) fruits. This species is known only from the type specimen, collected over a century ago close to a current protected area (Manongarivo). Despite numerous botanical explorations in the surrounding areas over the last five decades, no other material is available, suggesting perhaps a very small and restricted population or that the species is now extinct.




Fig. 51.
 Holotype of
Noronhia populifolia
 H. Perrier.

[Perrier de la Bâthie 8825, P]
 [© Muséum national d'Histoire naturelle,
 Paris. Reproduced with permission.]