

Zeitschrift: Boissiera : mémoires de botanique systématique
Herausgeber: Conservatoire et Jardin Botaniques de la Ville de Genève
Band: 70 (2016)

Artikel: A taxonomic revision of the genus Noronhia Stadtm. ex Thouars (Oleaceae) in Madagascar and the Comoro Islands
Autor: Hong-Wa, Cynthia
Kapitel: 69: Noronhia ratovosonii
DOI: <https://doi.org/10.5169/seals-1036126>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

Download PDF: 15.05.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

69. *Noronhia ratovosonii* Hong-Wa, spec. nova (Fig. 54).

Typus: MADAGASCAR. Prov. Antsiranana: DIANA, Andranovondronina, Anjabe, forêt de Belamoty, à 3 km au NE, 12°06'09"S 49°19'34"E, 85 m, 10.XI.2006, Ratovoson et al. 1153 (holo- : MO-6615566!; iso- : CNARP, G, P [P04254202] image seen, TAN).

Diagnosis *Noronhia ratovosonii* Hong-Wa can be distinguished from other members of the genus by its coriaceous, linear to obovate leaf blades, with a retuse to obcordate apex, and its ovoid, somewhat rugose fruits.

Description Shrubs to 2 m tall, trunk to 5 cm diameter; young twigs cylindrical, 1-1.5 mm diameter, glabrous; bark medium gray, smooth to slightly rugose. Leaves opposite, persistent; bud scales persistent; blades medium green above, lighter below, linear to obovate, 4-6 × 1-2.5 cm, coriaceous, glabrous, domatia absent, base rounded, margin revolute, apex retuse to obcordate, sometimes slightly acuminate, the acumen 0-3 mm long, midrib sunken above, raised below, secondary veins inconspicuous, 6-8 per side, 5-12 mm apart, looping 1-2 mm from the margin; petiole light gray to whitish, 2.5-5 × 1-1.3 mm, entirely woody, glabrous. Flowers unseen, but fruits solitary. Fruiting pedicel 4-8 × 1-1.5 mm; young fruits green, brownish when mature, ovoid, 8.5-17.5 × 7-10 mm, surface smooth to rugose, sometimes covered with a white pellicle, apex rostellate, the rostellum flattened, ridged, rounded, with the persistent style; dry pericarp 0.7 mm thick; endocarp woody.

Etymology This species is dedicated to Fidisoa “Fidy” Ratovoson, a botanist at the Missouri Botanical Garden in Madagascar, for collecting the type specimen and other numerous materials of *Noronhia*, for helping to locate specimens deposited at CNARP, TAN and TEF, and otherwise for assisting with various aspects of this work.

Distribution, ecology and phenology *Noronhia ratovosonii* occurs in low-elevation dry forests on limestones and unconsolidated sands in the north, around Montagne des Français (Fig. 49). Its flowering period is unknown, and it fruits from August to November.

Conservation status Based on four collections representing three localities, the assessment yielded an EOO of 74 km², an AOO of 12 km², and three subpopulations representing three locations, of which only one occurs within a protected area (Montagne des Français). Suitable habitat, both outside and inside of protected areas, will likely continue to experience some form of degradation and loss resulting from [illicit] forest exploitation, invasive species, and expansion of agricultural fields. Consequently, *N. ratovosonii* is assigned a preliminary status of “Endangered” [EN Blab(i,ii,iii,iv,v)+2ab(i,ii,iii,iv,v)].

Notes *Noronhia ratovosonii* is similar to *N. longipedicellata*, but can distinguished by its habit (shrubs vs. trees), retuse to obcordate (vs. retuse to mucronate) leaf blades, and shorter pedicels (less than 1 cm vs. 1-3 cm long). The new species can be recognized by its coriaceous, linear to obovate leaves and ovoid, somewhat rugose fruits.

Paratypi **MADAGASCAR. Prov. Antsiranana:** Andranovondronina, Anjabe, Antsaravy, forêt Antafian'i Tsirambaza, 12°08'53"S 49°20'31"E, 8 m, 17.III.2006, Guitou 318 (CNARP, MO, P, TAN); Ramena, Andavakoera, 12°19'57"S 49°21'19"E, 172 m, 4.VIII.2007, Hong-Wa et al. 525 (CNARP, MO, P, TAN); *ibid. loc.*, Hong-Wa et al. 548 (TAN).




Fig. 54.

Noronhia ratovosonii Hong-Wa.

A. Fruiting branch; **B.** Adaxial side of leaf blade; **C.** Abaxial side of leaf blade; **D.** Fruit.

[Hong-Wa 525, TAN] Drawings: R. L. Andriamariisoa