

Zeitschrift: Boissiera : mémoires de botanique systématique
Herausgeber: Conservatoire et Jardin Botaniques de la Ville de Genève
Band: 70 (2016)

Artikel: A taxonomic revision of the genus *Noronhia* Stadtm. ex Thouars (Oleaceae) in Madagascar and the Comoro Islands
Autor: Hong-Wa, Cynthia
Kapitel: 84: *Noronhia urceolata*
DOI: <https://doi.org/10.5169/seals-1036126>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 11.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

84. *Noronhia urceolata* H. Perrier in Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 2: 300. 1949.

Lectotypus (designated here): MADAGASCAR. **Prov. Mahajanga:** Ankaizina, [14°30'S 48°55'E], 1200 m, s.d., *Perrier de la Bâthie* 8827 (P [P00418103]!). **Syntypus:** Massif du Manongarivo, [14°00'S 48°23'E], 800 m, V.1909, *Perrier de la Bâthie* 8824 (P [P03559253, P03559254]!).

Description *Shrubs* to small trees to 4 m tall, trunk to 6 cm diameter; young twigs cylindrical, 1-2 mm diameter, glabrous; bark light gray to whitish, smooth. *Leaves* opposite, persistent; bud scales persistent; blades medium green above, lighter below, oblong to elliptic, 8.5-18 × 2-4 cm, chartaceous, glabrous, domatia casual, base acute to attenuate, margin flat to slightly revolute, apex acuminate, the acumen 4-12 mm long, midrib slightly sunken above, distinctly raised below, secondary veins conspicuous, 8-14 per side, 11-17 mm apart, looping 1-4.5 mm from the margin; petiole light gray, 5-10 × 1.2-2 mm, entirely woody, glabrous. *Thyrses* fasciculate, multiflorous, diffuse; peduncle 2-17 mm long, very sparsely pubescent; pedicel 7-14 mm long, very sparsely pubescent; calyx very sparsely pubescent outside, glabrous inside, lobes deltate, 1.4-2.2 × 1.3-2 mm; corolla dark brown on dry material, urceolate, 3.5-5 mm long, glabrous on both sides, the tube 2-3.5 mm long, lobes widely ovate, apex rounded; corona absent; stamens 1.2-2.8 mm long, anthers widely ovate to cordiform; pistil 2-3 mm long, stigma capitate. *Fruiting* pedicel 8 × 1.3-1.4 mm; young fruits green, brown when mature, subglobose, 23.5-27 × 23-24.5 mm, surface smooth, sometimes covered with a white pellicle, apex flat to slightly apiculate; dry pericarp 0.8 mm thick; endocarp crustaceous; seed 11.5 × 9.5 mm.

Distribution, ecology and phenology *Noronhia urceolata* occurs in low- to mid-elevation transitional forests from Sambirano in the northwest to Antsalova in the west (Fig. 68). It produces flowers and fruits from October to May.

Conservation status Based on six collections representing five localities, the assessment indicated an EOO of 59,126 km², an AOO of 20 km², and five subpopulations representing five locations, of which only one occurs within a protected area (Bemaraha). Given that the actual AOO is probably larger than estimated but likely does not exceed 2,000 km², along with the small number of locations and projected continuing decline in habitat quality and in the number of mature individuals as well as habitat loss resulting from wildfire, grazing, expansion of agricultural fields, and forest exploitation, *N. urceolata* can be assigned a preliminary status of "Vulnerable" [VU B2ab(i,ii,iii,iv,v)].







Fig. 68. Distribution maps of species of *Noronhia* Stadtm. ex Thouars. *N. urceolata* H. Perrier to *Noronhia verticillata* H. Perrier.

Notes *Noronhia urceolata* can be recognized by its chartaceous, oblong to elliptic leaf blades, large inflorescences with urceolate flowers, and subglobose fruits with crustaceous endocarp. It differs from *N. leandriana* by its woody petioles, the shape of its flowers and fruits, and the absence of a corona. *Noronhia urceolata* was based on two syntypes (*Perrier de la Bâthie* 8824 and 8827), the latter of which has better material and is thus chosen as the lectotype.

**Additional specimens
examined**

MADAGASCAR. **Prov. Antsiranana:** Ambanja, Marovato, [13°57'S 48°33'E], 16.XII.1951, *Saboureau* 3275 (MO, P); Ambanja, forêt Bemanasy, [13°47'S 48°37'E], 14.XI.1953, *Service Forestier* 7900 (MO, P, TEF). **Prov. Mahajanga:** Antsingy, vers Bevary (E d'Antsalova), [18°37'S 44°48'E], 400-600 m, 27.I.1960-5.II.1960, *Leandri & Saboureau* 2837 (G, MO, P); vers Ambondro, Antsingy, [18°40'S 44°52'E], 400-600 m, 2.II.1960, *Leandri & Saboureau* 2912 (MO, P); Kimanambolo, Belit-saka, 18°02'09"S 44°19'27"E, 18.X.2009, *Rakotonasolo et al.* 1401 (MO, P, TAN).