

<b>Zeitschrift:</b>	Boissiera : mémoires de botanique systématique
<b>Herausgeber:</b>	Conservatoire et Jardin Botaniques de la Ville de Genève
<b>Band:</b>	71 (2017)
<b>Artikel:</b>	Taxonomic treatment of Abrahamia Randrian. & Lowr : a new genus of Anacardiaceae from Madagascar
<b>Autor:</b>	Randrianasolo, Armand / Lowry, Porter P. / Schatz, George E.
<b>Kapitel:</b>	4: Abrahamia capuronii
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-1036071">https://doi.org/10.5169/seals-1036071</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 11.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

**4. *Abrahamaia capuronii* Randrian. & Lowry, spec. nova** (Fig. 4).

**Typus:** MADAGASCAR. Prov. Antsiranana: Rég. SAVA, Dist. Sambava, S de l'Anjehabe, [14°18'30"S 49°44'40"E], c. 600 m, 7.XI.1950, fl., Service Forestier 767 (holo-: P [P06774827]!; iso-: G [G00341710]!, MO-6684132!, NY!, P [P06774828, P06774829]!, TEF [TEF000881]!).

**Diagnosis** *Abrahamaia capuronii* Randrian. & Lowry can be distinguished from all other species of *Abrahamaia* by a combination of its elliptic leaves, narrow corolla lobes ( $3 \times 1.2$  mm), and slender inflorescence axes.

**Description** Trees 10-15 m tall, 20 cm DBH, bark with white or translucent latex; young twigs only pubescent at the tips. Leaves alternate or sub-opposite; blade obovate to elliptic,  $4-7.8 \times 2-3.6$  cm, chartaceous to subcoriaceous, apex rounded or retuse, margin undulate, base cuneate, adaxial surface glabrous, abaxial surface with short appressed indument, mainly along the midvein and at the base, venation craspedodromous, midvein prominent abaxially, secondary veins 18-33 pairs, 1-4 mm apart, prominent abaxially, slightly raised adaxially, tertiary veins slightly visible on adaxial surface; petiole 10-25 mm long, c. 0.5-1 mm in diam., canaliculate on adaxial surface, glabrescent. Inflorescence axillary or terminal, a panicle, 2-8 cm long, axes slender, strigose. Male flowers 5-merous, small, c. 2-2.5 mm long, subtended by a very small, pubescent bract; pedicel short, c. 0.3-0.5 mm long, pubescent; calyx lobes small, widely ovate,  $1 \times 1$  mm, valvate, connate at the base, adaxial surface glabrous, abaxial surface sparsely covered with short indument, glabrescent; corolla lobes  $3 \times 1.2$  mm, lanceolate, imbricate, adaxial surface glabrous, abaxial surface pubescent along its longitudinal axis or glabrescent to glabrous, white (according to field notes); stamens 5, filament c. 1.5 mm, glabrous, anthers c. 1 mm long, obloid, glabrous; disk glabrous, c. 1.2-1.5 mm in diam.; ovary rudimentary. Mature and opened female flowers unknown, developing buds with calyx lobes triangular to deltate, c.  $1 \times 0.5-0.8$  mm, adaxial surface glabrous, abaxial surface with some indument; corolla lobes ovate, c.  $1.5 \times 1$  mm, mostly glabrous but abaxial surface with a few appressed short trichomes; stamens vestigial; disk glabrous; ovary depressed, ovoid,  $0.3-0.5 \times 0.5$  mm, glabrous; style c. 1 mm, glabrous, stigmas capitate. Young fruits ellipsoid, very slightly asymmetrical, glabrous, with longitudinal striations.

**Etymology** The species epithet honors René Capuron, whose extensive field work contributed so much to our knowledge of Madagascar's woody flora, and who personally collected material of 24 of the 34 species of *Abrahamaia*.

**Distribution, ecology and phenology**

*Abrahamia capuronii* is restricted to far northeastern Madagascar (Map 1), where it occurs in mid-elevation humid and transitional forests on rocky substrate up to an elevation of 700 m. Material with buds and flowers has been collected in October and early November, and specimens with young fruits have been gathered in late November.

**Conservation status**

With an EOO of 2,585 km<sup>2</sup> and a minimum AOO of 24 km<sup>2</sup>, *A. capuronii* would qualify as “Endangered” under Criteria B1 and B2 of the IUCN Red List (IUCN, 2012) if other qualifying criteria were met. However, it is known from a total of 6 locations, one of which is situated within a protected area (Loky Manambato) while the others are at sites potentially subjected to clearing and habitat degradation, and should thus be assigned a preliminary status of “Vulnerable” [VU B1ab(iii)+2ab(iii)].

**Notes**


*Abrahamia capuronii* resembles both *A. grandidieri* and *A. sericea* by its relatively closely spaced, parallel and craspedodromous secondary veins, but can be distinguished from these two species by its smaller number of secondary veins (10-33 in *A. capuronii* vs 35-40(-47) in *A. grandidieri* and 40-70 in *A. sericea*).

**Paratypi**

**MADAGASCAR. Prov. Antsiranana:** Mont Anjenabe, [14°17'S 49°46'E], 650 m, 3-7.XI.1950, fl., *Humbert* 24055 (G, K, MO, P [2 sheets], TAN); Vohémar, forêt de Binara, 13°15'S 49°37'E, 700 m, 8.XI.2001, fl., *Gautier et al.* 4125 (G, MO, TAN); Vohémar, Analamazava forest, 13°15'43"S 49°35'22"E, 25.XI.2004, y. fr., *Ranarivelo* 446 (MO); Daraina, Ambilobe, 13°44'10"S 49°23'07"E, 1299 m, 27.X.2007, bud, *Randriambololomamonjy et al.* 166 (MO, P, TAN); Vohémar, Daraina, forêt d'Analamazava, [13°15'58"S 49°36'14"E], 700 m, 29.X.2005, fl., *Randrianaivo* 1259 (CNARP, MO, P, TAN); Vohémar, Daraina, forêt d'Antsahabe, 13°12'37"S 49°33'29"E, 545 m, 31.X.2005, fl., *Razafitsalama* 741 (CNARP, MO, P, TAN); Vohémar, Andrafainkona, Ampisarahana, 13°38'28"S 49°31'51"E, 1009 m, 9.XI.2007, bud, *Razakamalala et al.* 3716 (G, MO, P, TAN).



**Fig. 4.** *Abrahamia capuronii* Randrian. & Lowry. **A.** Flowering branch; **B.** Flowers (one with a petal removed); **C.** Detail of leaf; **D.** Inflorescence. [Service Forestier 767, P] Drawings: R.L. Andriamiarisoa



**Fig. 5.** *Abrahamia darainensis* Randrian. & Lowry. **A.** Fruiting branch; **B.** Fruit.

[Ranirison & Nusbaumer 1118, MO] Drawings: R.L. Andriamiarisoa