

Zeitschrift:	Boissiera : mémoires de botanique systématique
Herausgeber:	Conservatoire et Jardin Botaniques de la Ville de Genève
Band:	71 (2017)
Artikel:	Taxonomic treatment of Abrahamia Randrian. & Lowr : a new genus of Anacardiaceae from Madagascar
Autor:	Randrianasolo, Armand / Lowry, Porter P. / Schatz, George E.
Kapitel:	9: Abrahamia ellipticarpa
DOI:	https://doi.org/10.5169/seals-1036071

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 11.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

9. *Abrahamia ellipticarpa* Randrian. & Lowry, spec. nova (Fig. 8).

Typus: MADAGASCAR. **Prov. Toliara:** Rég. Anosy, Vinanibe, à l'W de Fort-Dauphin, [25°03'S 46°56'E], 7.XII.1968, fr., Service Forestier 28611 (P [P00722641]!; iso-: G [G00341714]!, K!, MO-6684134!, P [P00722642]!, TEF [TEF000884]!).

Diagnosis *Abrahamia ellipticarpa* Randrian. & Lowry can be distinguished from other member of the genus by its large, glabrous, ellipsoid fruits measuring 3-3.5 × 1.5-1.8 cm.

Description Trees c. 10-12 tall, bark with white latex; young twigs with appressed indument. Leaves alternate, subopposite or opposite; blade oblanceolate to narrowly elliptic, 3-9.5 × 1.3-3 cm, subcoriaceous, apex acute or sometimes retuse or rarely emarginate, margin entire, base cuneate, adaxial surface glabrous, abaxial surface with appressed indument, more so along the midvein and margin, venation craspedodromous, midvein prominent abaxially, secondary veins 18-30 pairs, more or less parallel, 2-3 mm apart, slightly raised adaxially and prominent abaxially, tertiary veins very few, impressed and barely visible; petiole 5-10 mm long, c. 1 mm in diam., canaliculate on adaxial surface, pubescent. Inflorescences and flowers unknown. Fruits 3-3.5 × 1.5-1.8 cm, ellipsoid, glabrous, with longitudinal striations.

Distribution, ecology and phenology *Abrahamia ellipticarpa* has been recorded from sublittoral forest on sand at several sites in the Fort-Dauphin area, including Mandena, Vinanibe, and Petriky (Map 1). It has been collected in flower in September and in fruit in December.

Conservation status With an EOO of 28 km², an AOO of 12 km², and three known locations, 2 of which are situated within protected areas (Mandena and Petriky) whereas the third one is highly degraded and subjected to continued degradation from uncontrolled harvesting and possibly also fire leading to a projected decline in all 5 sub-criteria, *A. ellipticarpa* is assigned a preliminary conservation status of “Endangered” [EN B1ab(i,ii,iii,iv,v)+2ab(i,ii,iii,iv,v)] using the IUCN Red List Categories and Criteria (IUCN, 2012).

Notes *Abrahamia ellipticarpa* is vegetatively similar to *A. oblongifolia* but can be distinguished by its fewer secondary veins (18-30 vs 35-40), impressed and barely visible (vs slightly raised and very visible) tertiary venation, and large fruits (3-3.5 × 1.5-1.8 vs 2 × 1 cm) with evident (vs obscure) longitudinal striations.

Paratypi MADAGASCAR. **Prov. Toliara:** Petriky, N shore of Lake Andranany, 25°03'S 46°53'E, 0-10 m, 14.IV.1989, ster., Gereau 3376 (MO, P, TAN); env. de Fort-Dauphin, entre le Pic St. Louis et la mer, 1-25 m, 20.IX.-6.X.1928, fl., Humbert 5984 (G, P [2 sheets]); Man-

dena forestry station, 24°57'S 47°00'E, 0-10 m, 20.XI.1996, ster., A. Randrianasolo 578 (MO, P, TAN); *ibid. loco*, 3.IX.1953, bud, Service Forestier 8171 (MO, P, TEF); *ibid. loco*, 12.XII.1952, ster., Service Forestier 207-R-16 (P).

