

Zeitschrift:	Boissiera : mémoires de botanique systématique
Herausgeber:	Conservatoire et Jardin Botaniques de la Ville de Genève
Band:	71 (2017)
Artikel:	Taxonomic treatment of Abrahamia Randrian. & Lowr : a new genus of Anacardiaceae from Madagascar
Autor:	Randrianasolo, Armand / Lowry, Porter P. / Schatz, George E.
Kapitel:	25: Abrahamia patrickii
DOI:	https://doi.org/10.5169/seals-1036071

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 11.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

25. *Abrahamia patrickii* Randrian. & Lowry, spec. nova (Fig. 22).

Typus: MADAGASCAR. Prov. Antsiranana: Rég. SAVA, Vohémar, commune rurale de Daraina, forêt de Bobankora, partie N, 510 m, 13°13'15"S 49°45'42"E, 510 m, 29.I.2005, fr., Ranirison & Nusbaumer 919 (holo-: G [G00019652]!; iso-: MO-6671434!, P [P00580356]!, TEF [TEF000886]!).

Diagnosis *Abrahamia patrickii* Randrian. & Lowry can easily be distinguished from other members of the genus by its sub-verticillate to subopposite leaves, narrowly oblanceolate leaf blade with very closely spaced secondary veins, and the presence of circular orange-brown scales on the abaxial leaf surface.

Description Trees 13.5 m tall, 20 cm in diam.; young twigs lenticellate. Leaves sub-verticillate or subopposite; blade oblanceolate, 3-5.5 × 1-1.5 cm, subcoriaceous, apex emarginate or round, margin undulate, base cuneate to acute, adaxial surface glabrous, abaxial surface mostly glabrous but with scattered short reddish indument along midvein and margin, as well as circular, orange-brown scales, venation craspedodromous, midvein prominent abaxially, secondary veins 24-28 pairs, 1-3 mm apart, very often closely spaced, raised on both surfaces, tertiary veins not visible on either surface; petiole 6-13 mm long, c. 0.8-1 mm in diam., glabrous, canaliculate on adaxial surface. Inflorescence and flowers unknown. Fruits 2.3-2.6 × 1.1-1.3 cm, slightly asymmetric, ellipsoid, glabrous, with longitudinal striations.

Etymology The species epithet honors Patrick Ranirison for his dedication to the study of the Malagasy flora, most notably his work in the Daraina region of northeastern Madagascar.

Distribution, ecology and phenology *Abrahamia patrickii* is known only from the type collection from the Daraina area and a second gathering made N of Sambava (Map 6), in semi-deciduous and evergreen humid forest. It has been collected with fruits in January and February.

Conservation status With a minimum AOO of 8 km² and 2 localities, both of which situated are within protected areas (Loky Manambato and Makirovona), *A. patrickii* is assigned a preliminary conservation status of "Least Concern" [LC] using the IUCN Red List Categories and Criteria (IUCN, 2012).

Notes *Abrahamia patrickii* closely resembles *A. nitida* by its sub-verticillate leaves with an oblanceolate blade, similar number of craspedodromous secondary veins (24-28 in *A. patrickii* and 20-30 in *A. nitida*), similar petiole length (6-13 mm in *A. patrickii* and 5-15 mm in *A. nitida*), and glabrous fruit. However, the leaves of *A. patrickii* differ from those of *A. nitida* by their undulate (vs revolute) margins, orange-brown circular scales (vs glabrous) on the abaxial surface, narrowly spaced secondary veins (1-3 vs 3 mm apart), obscure (vs visible and raised) tertiary veins, and fruit with longitudinal striations (vs smooth) on the surface.

Paratype MADAGASCAR. Prov. Antsiranana: Rég. SAVA, Sambava, forêt d'Antsahaha Kasaka, 2 km NW du village Ambavala, 14°05'39"S 50°02'42"E, 146 m, 16.II.2006, fr., Andrianjafy et al. 1590 (MO, TAN, P).



Fig. 22.

Abrahamia patrickii Randrian. & Lowry.

A. Fruiting branch; **B.** Fruit. [Ranirison & Nusbaumer 919, MO] Drawings: R.L. Andriamiarisoa