

<b>Zeitschrift:</b>	Boissiera : mémoires de botanique systématique
<b>Herausgeber:</b>	Conservatoire et Jardin Botaniques de la Ville de Genève
<b>Band:</b>	71 (2017)
<b>Artikel:</b>	Taxonomic treatment of Abrahamia Randrian. & Lowr : a new genus of Anacardiaceae from Madagascar
<b>Autor:</b>	Randrianasolo, Armand / Lowry, Porter P. / Schatz, George E.
<b>Kapitel:</b>	29: Abrahamia sambiranensis
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-1036071">https://doi.org/10.5169/seals-1036071</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 11.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

**29. *Abrahamia sambiranensis*** Randrian. & Lowry, spec. nova (Fig. 26).

**Typus:** MADAGASCAR. Prov. Mahajanga: Rég. Diana, Ambato forêt classée, c. 40 km N d'Ambohaha, 13°26'S 48°33'E, 90-459 m, 23.XI.1996, fr., Randrianaivo et al. 29 (holo: MO-6671436!; iso: G!, MO-6671435!, P [P06775160]!, TAN [TAN001998]!).

**Diagnosis** *Abrahamia sambiranensis* Randrian. & Lowry can be distinguished from other members of the genus by the combination of large (10-19 × 2.5-5.6 cm), coriaceous leaves with numerous (50-60) distinctly raised secondary veins and a folded apex in dry material, its long petiole (30-65 mm), and its large, glabrous fruits (2.5-3 cm long).

**Description** Trees 10-15 m tall, 15-18 cm DBH, bark with white latex; young twigs pubescent towards the tips. Leaves alternate, clustered toward the branch ends; blade oblanceolate or narrowly elliptic to lanceolate, 10-19.4 × 3.2-5.5 cm, coriaceous, apex acuminate, folded in dry material, margin undulate, base cuneate to broadly cuneate, adaxial surface glabrous, often glaucous, abaxial surface sparsely sericeous, more so on midvein, venation craspedodromous, midvein prominent abaxially, secondary veins 50-60 pairs, more or less parallel, 2-4 mm apart, raised on both surfaces, tertiary veins not visible on both surfaces; petiole (20)-25-58 mm long, 2-3.5 mm in diam., canaliculate on adaxial surface, tomentose. Inflorescence axillary, a panicle, 2-9.5 cm long, very few flowers reaching maturity, axes pubescent. Male flowers 5-merous; pedicel c. 0.5-1 mm long, pubescent; calyx lobes triangular to ovate, 1 × 1 mm, connate at base, adaxial surface glabrous, abaxial surface pubescent; corolla lobes ovate, 2.5 × 1.5 mm, imbricate, adaxial surface glabrous, abaxial surface pubescent; stamens 5, filaments 1 mm long, anthers c. 1 × 1 mm, obloid to ovoid; disk cupuliform, glabrous; rudimentary ovary absent. Female flowers unknown. Fruits 2.5-3 × 1.5-1.8 cm, ellipsoid, pubescent when young, glabrescent, with fine longitudinal striations.

**Distribution, ecology and phenology** *Abrahamia sambiranensis* is restricted to humid forests in the Sambirano region of NW Madagascar and extends into the northern part of the island (Map 9). It flowers in July and August, and fruits have been collected in October and November.

**Conservation status** With an EOO of 1,717 km<sup>2</sup>, a minimum AOO of 28 km<sup>2</sup>, and seven localities, with only a single subpopulation within a protected area (Galoko), *A. sambiranensis* exists at 7 locations, six of which are subjected to pressure from forest exploitation and land clearing; it is therefore assigned a preliminary conservation status of "Vulnerable" [VU B1ab(iii)+2ab(iii)] using the IUCN Red List Categories and Criteria (IUCN, 2012).

**Notes** *Abrahamia sambiranensis* vegetatively resembles *A. elongata*, but differs by having leaves that lack tertiary venation (vs present and raised on both surfaces in *A. elongata*) and whose apex is usually folded in pressed material (vs flat), and by its fruits that are glabrescent when mature (vs glabrous in *A. elongata*).

**Paratypi** **MADAGASCAR.** Prov. Antsiranana: Rég. Diana, Ambilomangodra, 13°02'42"S 49°09'11"E, 50 m, 14.VII.2005, fl., Randrianaivo et al. 1201 (CNARP, MO, P, TAN); forêt de Kalabenono, 13°37'27"S 48°40'05"E, 369 m, 10.XI.2006, fr., Razafitsalama et al. 1064 (G, MO, P, TAN); Galoko, [13°33'45"S 48°45'00"E], 29.VII.1955, y. fr., Service Forestier 15748 (P, TEF); forêts à l'E d'Ampondrabe, 28.X.1954, fr., Service Forestier 18931 (MO, P [2 sheets]); massif de l'Andavakoera, N d'Angodromena, [13°06'30"S 49°18'30"E], 1.X.1958, fr., Service Forestier 18947 (G, MO, P [2 sheets], TEF); Ampandrikoera, 12.X.1951, ster., Service Forestier 181-R-64 (P).

