

Zeitschrift:	Boissiera : mémoires de botanique systématique
Herausgeber:	Conservatoire et Jardin Botaniques de la Ville de Genève
Band:	71 (2017)
Artikel:	Taxonomic treatment of Abrahamia Randrian. & Lowr : a new genus of Anacardiaceae from Madagascar
Autor:	Randrianasolo, Armand / Lowry, Porter P. / Schatz, George E.
Kapitel:	33: Abrahamia turkii
DOI:	https://doi.org/10.5169/seals-1036071

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.10.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

33. *Abrahamia turkii* Randrian. & Lowry, spec. nova (Fig. 31).

Typus: MADAGASCAR. Prov. Fianarantsoa: Rég. Vatovavy-Fitovinany, Ranomafana National Park, parcelle 3, S of National road 25 at 7 km W of Ranomafana, Talatakely trail system, 21°15'30"S 47°25'00"E, 950-1150 m, 20.X.1993, fl., Solo & J. Randrianasolo 35 (holo-: MO-6671437!; iso-: G, K, MO-6671438!, P [P06168911]!, TAN [TAN002002]!, WAG).

Diagnosis *Abrahamia turkii* Randrian. & Lowry can be distinguished from other members of the genus by its coriaceous leaves, inflorescence with thick axes and ferruginous indument, and fruits that are small in size (1.8-2 × 1 cm) and glabrous.

Description Trees 11-30 m tall, 18-100 cm DBH, bark with white latex; young twigs sericeous on tips, lenticellate farther from the tip. Leaves usually alternate, sometimes subopposite; blade obovate, 2.5-8 × 1.5-3.5 cm, coriaceous, apex usually emarginate or retuse, sometimes rounded, margin slightly revolute, sometimes undulate, base cuneate, adaxial surface sparsely pubescent, abaxial surface sericeous or strigose, venation craspedodromous, midvein prominent abaxially, secondary veins 14-24 pairs, 1-4 mm apart, raised on both surfaces, tertiary veins visible on both surfaces; petiole 0.5-1 cm long, 1-1.5 mm in diam., pubescent or scabrous, canaliculate on adaxial surface. Inflorescence axillary or terminal, a panicle, erect, axes thick, 1-5 cm long, reddish brown pubescent. Male flowers 5-merous, c. 3-4 mm long; pedicel c. 0.3-0.5 mm long, reddish brown pubescent; calyx lobes triangular, 1.5-2 × 1-2 mm, adaxial surface glabrous, abaxial surface, pubescent; corolla lobes ovate, 2-4 × 2 mm, adaxial surface glabrous, abaxial surface pubescent; stamens 5, filament 1-2 mm long, glabrous, anthers 1 mm long, ovoid to obloid, glabrous; disk slightly cupuliform, c. 2 mm in diam., glabrous; ovary very rudimentary or absent. Female flowers unknown. Fruits 1.8-2 × 1 cm, ovoid, glabrous, wrinkled when dry (longitudinal striations not as evident as in other species).

Distribution, ecology and phenology *Abrahamia turkii* is distributed mainly in south-central and southern Madagascar, although one collection has been made at the Ambohitantely forest, far to the north of the other recorded subpopulations (Map 10). Flowering material has been recorded in October and November, and fruits have been collected in January.

Conservation status With an EOO of 85,992 km², a minimum AOO of 40 km² but an estimated value well exceeding 2,000 km², and 9 localities, with subpopulations in 6 protected areas (Ambohitantely, Andohahela, Betampona, Kalambatritra, Pic d'Ivohibe, and Ranomafana), representing 9 locations, only some of which may be subjected to pressure from forest exploitation and land clearing, *A. turkii* is assigned a preliminary conservation status of "Least Concern" [LC] using the IUCN Red List Categories and Criteria (IUCN, 2012).

Notes *Abrahamia turkii* closely resembles *A. ditimena* by its medium sized, coriaceous leaves with craspedodromous venation, usually emarginate or retuse apex, and tertiary venation that is visible on both surfaces, and its paniculate inflorescence, but it can easily be distinguished by its short (1-5 vs 2-10 cm) inflorescence covered with reddish brown pubescence (vs glabrescent or tomentose), and its glabrous (vs velutinous) fruits.

Paratypi **MADAGASCAR.** **Prov. Antananarivo:** forêt d'Ambohitantely, 18 km NW of Ankazobe, [18°10'S 47°18'E], 1400-1646 m, 21.XI.1985, bud, Barnett et al. 533 (MO, P, TAN). **Prov. Fianarantsoa:** Ivohibe RS, piste II et III, 22°29'00"S 46°58'06"E, 1275 m, 20.X.1997, fl., Rakotomalaza et al. 1434 (G, MO, P, TAN); Ivohibe RS, 5.5 km SE d'Angondongodona, camp 4, 22°25'18"S 46°53'54"E, 1150 m, 4.XI.1997, fl., Rakotomalaza et al. 1494 (G, K, MO, P, TAN); Haute Matsiatra, Ialatsara forest, 21°03'07"S 47°12'42"E, 12.X.2005, fl., Ranarivelo et al. 719 (CAS, MO); Ranomasana PN, parcelle 3, Vatoharana trail system, 21°17'S 47°26"E, 1100-1250 m, 18.XI.1993, fl., Turk & J. Randrianasolo 585 (G, MO, P, TAN); Andrambovato, E de Fianarantsoa, [21°31'S 47°25"E], 878-1000 m, 24.I.1955, fr., Service Forestier 11590 (G, MO, P, TEF). **Prov. Toamasina:** RN1 [Betampona], s.d., fl. & y. fr., Anonymous s.n. (G, MO, P [2 sheets]). **Prov. Toliara:** Betroka, Kalambatritra RS, forêt d'Analamaro, 23°28'28"S 46°25'15"E, 1423 m, 9.I.2004, fl., Andrianjafy et al. 601 (MO, P, TAN); Andohahela PN, parcelle 1, 13.5 km NW d'Eminiminy, 24°35'S 46°44'1"E, 1200 m, 15.XI.1995, fl., Messmer et al. 105 (G, MO).