

Zeitschrift: Boissiera : mémoires de botanique systématique
Herausgeber: Conservatoire et Jardin Botaniques de la Ville de Genève
Band: 71 (2017)

Artikel: Taxonomic treatment of *Abrahamia* Randrian. & Lowr : a new genus of Anacardiaceae from Madagascar
Autor: Randrianasolo, Armand / Lowry, Porter P. / Schatz, George E.
Kapitel: 34: *Abrahamia viguieri*
DOI: <https://doi.org/10.5169/seals-1036071>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 25.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

34. *Abrahamia viguieri* (H. Perrier) Randrian. & Lowry, **comb. nova.**

≡ *Protorhus viguieri* H. Perrier in Mém. Mus. Natl. Hist. Nat. 18: 262. 1944.

Lectotypus (designated here): **MADAGASCAR. Prov. Toamasina:** Rég. Alaotra-Mangoro, Moramanga, forêt d'Analamazaotra, [18°56'S 48°26'E], 1000 m, 19.X.1912, fl., *Humbert & Viguier 815* (P [P00580348]!; iso-: K, MO-6684877!, P [P00580349, P02141537, P06242027]!).

Description

Trees 2-25 m tall, 10-15 cm DBH; young twigs lanate. *Leaves* alternate, sometimes verticillate; blades oblanceolate, 8-30 × 2-6 cm, chartaceous, apex acuminate, margin slightly revolute, base acute or cuneate, adaxial surface lanate, abaxial surface pubescent or lanate only on midvein, venation craspedodromous, midvein prominent abaxially, secondary veins 30-40 pairs, 2-15 mm apart, prominent abaxially, tertiary veins visible on both surfaces; petiole 5-20 mm long, canaliculate on adaxial surface, lanate. *Inflorescence* axillary, a panicle, 3-11 cm long, axes terete, villous or lanate. *Male and female flowers* 5-merous, glabrous; pedicel 0.2-0.3 mm long, villous; calyx lobes triangular, 0.5 × 0.5-0.7 mm, imbricate, adaxial surface glabrous, abaxial surface pubescent to lanate, persistent on fruit; corolla lobes ovate, 2 × 1 mm, imbricate or quincuncial, adaxial surface glabrous, abaxial surface pubescent on the longitudinal main axis; stamens 5; disk c. 1-1.2 mm in diam., pubescent. *Male flowers* with filaments 1.5 mm long, anthers 0.5 mm long, ovoid to subglobose, yellow, glabrous; ovary completely absent. *Female flowers* with filaments 0.5-0.7 mm long, anthers 0.3-0.5 mm long, ovoid, sterile, yellow, glabrous; ovary widely depressed ovate, 1-1.2 × 0.8-1 mm, glabrous, placentation subapical, style simple (or with a minute trace of division), apically inserted, 0.5-0.7 mm long, glabrous; stigma 4-lobed, capitate. *Fruits* 3 × 1.5 cm, ellipsoidal, symmetrical or very slightly asymmetrical, glabrous, with longitudinal striations.

Etymology

The species is dedicated to Dan Turk, a forest botanist who has contributed tremendously to our knowledge of the trees of Ranomafana National Park.

Vernacular names

“Lambafohala”, “Natoboka”, “Sefana”, “Tsitsebona”.

Distribution, ecology and phenology

Abrahamia viguieri ranges from east coast humid littoral forests on sand and laterite at low elevation to premontane forests along the eastern escarpment at an elevation of up to c. 900 m (Map 9). Material has been collected with flowers in October and November; very young fruits have been recorded in December and mature fruits in late February.

Conservation status With an EOO of 28,641 km², a minimum AOO of 60 km², and 17 localities, with subpopulations in six protected areas (Ambalabe, Analalava, Betampona, Makira, Analamazaotra, and Zahamena), *A. viguieri* is assigned a preliminary conservation status of “Least Concern” [LC] using the IUCN Red List Categories and Criteria (IUCN, 2012).

Notes *Abrahamia viguieri* is easily distinguished from other members of the genus by the lanate upper surface of its leaves and its male flowers with a pubescent disk and completely lacking even a rudimentary or vestigial ovary.

PERRIER DE LA BÂTHIE (1944) cited two collections in the protologue of *Protorhus viguieri*, one with male flowers (*Humbert & Viguier 815*) and the other with fruits (*Perrier de la Bâthie 4387*, erroneously cited as 4397). We have selected a specimen of *Humbert & Viguier 815* as the lectotype because it is the best preserved among the syntypes and because an isolectotype was available for distribution to MO.

Additional material examined **MADAGASCAR. Prov. Fianarantsoa:** Fort Carnot, exploitation Charlier, [21°51'30"S 47°26'30"E], 23.II.1954, fr., *Service Forestier 14652* (MO, P, TEF). **Prov. Toamasina:** Vatoman-dry, Ambalabe, Ambinanindrano II, forêt de Vohibe, Sahamanahirana, 19°09'32"S 48°34'39"E, 635 m, 28.X.2012, bud & fl., *Andriamiarinoro 304* (MO, P, TAN); Brickaville, Sahatsara, 18°26'23"S 48°49'40"E, 698 m, 28.X.2005, fl., *Andriamihajarivo et al. 620* (MO, P, TAN, TEF); Brickaville, Anivorano-Sud, Ankerana, 18°26'03"S 48°49'53"E, 575 m, 31.X.2005, bud, *Andriantiana 297* (P, MO, TAN); Maroantsetra, Rantabe, Behovana village, 15°47'34"S 49°32'45"E, 521 m, 31.X.2003, fl., *Antilahimena 2237* (G, K, MO, NY, P, TAN, WAG); Brickaville, Didy à Brickaville, [18°07'S 48°32'E], s.d., fl., *Cours 4823* (MO, P [3 sheets]); Moramanga, station forestière d'Andasibe, [18°56'S 48°26'E], 900 m, 8.XII.1989, bud & fl., *Evrard 11232* (P, TAN); Betampona RN, 17°31'S 49°07'E, 275-650 m, 26.IX.1993, bud, *Lewis et al. 619* (G, K, MO [2 sheets], NY, P, TAN, WAG); forêt d'Analamazaotra, [18°55'48"S 48°25'48"E], 800 m, s.d., fr., *Perrier de la Bâthie 4387* (P); Zahamena PN, NE à partir de la rivière de Moango, 17°33'49"S 48°54'01"E, 794 m, 20.X.2000, fl., *Rakotonandrasana et al. 441* (CNARP, MO, P, TAN); Vavatenina, Zahamena PN, Besakay, 17°39'20"S 48°54'22"E, 780 m, 7.X.2001, bud, *Rakotonandrasana et al. 573* (CNARP, MO, P, TAN); Analamazaotra RS, along the Indri trail, about 300 m from observation point, 18°56'12"S 48°25'10"E, 1000 m, 10.I.2003, ster., *A. Randrianasolo 776* (MO, TEF); Fotsimavo, canton Tamatave, [17°56'S 49°13'E], 10.XI.1955, bud & fl., *Réserves Naturelles 7712* (MO, P, TEF); Ambodiriana, [17°55'S 49°13'E], 100-200 m, 12.IV.1959, bud, *Réserves Naturelles 10847* (MO, P); Sahatavy, [Zahamena PN], [17°38'S 48°50'E], 750-1912 m, 18.X.1960, bud, *Réserves Naturelles 11360* (MO, P [2 sheets], TEF); Fénériver-Est, forêt d'Analalava, Foulpointe, [17°42'S 49°29'E], 0-50 m, 30.X.1963, bud & fl., *Service Forestier 22781* (P, TEF); Ambatonandrohina, Sahatavy, [17°27'S 49°00'E], 13.X.1966, bud & fl., *Service Forestier 26261* (P, TEF).