

Zeitschrift:	Boissiera : mémoires de botanique systématique
Herausgeber:	Conservatoire et Jardin Botaniques de la Ville de Genève
Band:	71 (2017)
Artikel:	Taxonomic treatment of Abrahamia Randrian. & Lowr : a new genus of Anacardiaceae from Madagascar
Autor:	Randrianasolo, Armand / Lowry, Porter P. / Schatz, George E.
Kapitel:	Taxonomic treatment
DOI:	https://doi.org/10.5169/seals-1036071

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 11.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Fruits	The fruits of <i>Abrahamia</i> are always drupes with a resiniferous mesocarp and thin papyraceous endocarp, but several characters are very useful for delimiting species.
Shape	Fruits in <i>Abrahamia</i> can vary from globose or ellipsoid to ovoid or obovoid, depending on the species.
Size	Fruit dimensions are also useful for delimiting species within <i>Abrahamia</i> , varying from $1-1.5 \times 0.5-0.8$ cm in <i>A. minutifolia</i> , which has the smallest fruits, to $3-3.5 \times 1.5-1.8$ cm in <i>A. ellipticarpa</i> Randrian. & Lowry, which has the largest fruits in the genus.
Surface indument	The fruits of <i>Abrahamia</i> are either glabrous or velutinous, and this character is very useful for distinguishing species. Eight species have fruits with indument, viz. <i>A. betamponensis</i> Randrian. & Lowry, <i>A. buxifolia</i> , <i>A. delphinensis</i> , <i>A. ditimena</i> , <i>A. itremoensis</i> , <i>A. lenticellata</i> , <i>A. minutifolia</i> , and <i>A. sericea</i> , while those of the remaining 25 species are glabrous (fruits of <i>A. pauciflora</i> are unknown).

Taxonomic treatment *Abrahamia* Randrian. & Lowry, **gen. nov.**

Typus: *Abrahamia ditimena* (H. Perrier) Randrian. & Lowry (\equiv *Protorhus ditimena* H. Perrier).

Diagnosis Dioecious trees or shrubs. Leaves simple, persistent, opposite, subopposite, verticillate or alternate, chartaceous, subcoriaceous or coriaceous, lateral veins parallel, arcuate, forming an obtuse to acute angle with the midrib toward the base of the blade. Inflorescence erect, a thyrs or panicle. Flowers (4)-5-(6)-merous; pedicel short, articulated; sepals much smaller than petals; stamens alternipetalous, filament broadened at the base (sterile in female flowers, forming staminodes), anthers dorsifix, introrse, glabrous; ovary pubescent or glabrous (rudimentary or absent in male flowers), with 1 locule containing a single subpendulous or pendulous ovule. Fruits drupaceous, ellipsoidal or ovoid, symmetrical or slightly asymmetrical, resiniferous; seeds ruminate; cotyledons usually inseparable (except in *Abrahamia ibityensis* (H. Perrier) Randrian. & Lowry, in which the resiniferous canals are only peripheral).

Description Dioecious trees or shrubs; bark with white, translucent or red resinous latex. Leaves alternate, opposite or subopposite (verticillate in one species), simple, coriaceous, subcoriaceous or chartaceous, persistent (deciduous in *A. humbertii*). Venation pinnate, craspedodromous or semicraspedodromous, midvein usually flat, rarely very slightly raised above, prominently raised below (rarely on both surfaces), secondary veins parallel or nearly so, straight or arcuate, prominent below or on both surfaces, forming an obtuse to acute angle with the midrib toward the base of the blade. Inflorescence terminal and/or axillary, an erect thyrs or panicle, or intermediate and forming a paniculiform thyrs. Flowers usually 5-merous (sometimes 4- or 6-merous); sepals much smaller than petals;

stamens 5 or 4 (rarely 6), alternipetalous, filaments straight, flattened, broadened at the base, inserted basally on the outer surface of the disk, anthers dorsifixed, introrse, dehiscent with longitudinal slits (stamens reduced in size and sterile in female flowers, forming staminodes); disk usually cupuliform and crenulate in *male flowers*, annular and cupuliform in *female flowers*; ovary with 1 locule containing a single pendulous or subpendulous or ovule on a short funiculus, placentation subapical (ovary rudimentary or absent in male flowers); style one, short, apically trifid or unbranched with 3 capitate stigmatic lobes. Fruits drupaceous, ellipsoidal or ovoid, symmetrical or slightly asymmetrical, exocarp usually with longitudinal striations on surface, glabrous or velutinous, mesocarp resinous, endocarp thin, papyraceous; seed solitary, with resiniferous canals throughout (only peripheral in *A. ibityensis*), ruminant; cotyledons usually inseparable (separable in *A. ibityensis*).

Etymology

The genus is named in honor of the late Jean Prosper Abraham (1936–1996), a well-known Malagasy forester with a remarkable knowledge of the woody plants of Madagascar and who contributed much to our knowledge of the island's rich and diverse flora.

Notes

Abrahamia includes a total of 34 species, all endemic to Madagascar, including 15 species formerly placed in *Protorhus* (e.g., by PERRIER DE LA BATHIE, 1944, 1946), to which 19 additional species are newly described here. With the recognition of *Abrahamia* as a separate genus, *Protorhus* is now restricted to a single species in Africa, *P. longifolia*. The two genera differ in several flower and fruit characters, as indicated by PELL (2004) and PELL et al. (2011), and summarized in Table 1.

Table 1.
Morphological features
distinguishing *Abrahamia*
Randrian. & Lowry
from *Protorhus* Engl.

	<i>Abrahamia</i>	<i>Protorhus</i>
Fruit shape	Ellipsoidal or ovoid, radially symmetrical	Ovoid, mango-shaped or transversely oblong, asymmetrical
Seed	Resiniferous canals present, making the seed ruminant; cotyledons inseparable	Resiniferous canals lacking, seed not ruminant; cotyledons separable
Number of styles	1 (apically trifid or with 3 capitate stigmas)	3, slightly connate at the base or completely free

Abrahamia is represented in all of Madagascar's major vegetation types (MOAT & SMITH, 2007) and broad bio-climatic zones (CORNÉT, 1974; SCHATZ, 2000). Species can be found in humid forest in the eastern and Sambiranano regions (*A. antongilensis*, *A. sambiranensis*), wooded grassland and bushland in the center of the island (*A. ibityensis*, *A. buxifolia*), western dry forest (*A. deflexa*, *A. humbertii*), and southwestern subarid spiny forest-thicket (*A. phillipsonii* Randrian. & Lowry). Members of the genus occur on many types of substrate and at a wide range of elevations, from littoral forest on sand at the sea level (*A. nitida* (Engl.) Randrian. & Lowry, *A. suarezensis* Randrian. & Lowry) to humid forest on lateritic soil at altitudes up to 2000 m (*A. buxifolia*, *A. itremoensis*).

Species of *Abrahamia* were described by PERRIER DE LA BÂTHIE (1946) as polygamo-dioecious, indicating that he interpreted them as having both unisexual and hermaphroditic flowers on the same individual. However, careful observation has shown that all members of the genus are functionally dioecious. Female flowers have rudimentary, sterile stamens that are reduced in size, whereas male flowers either have an ovary that is rudimentary and non-functional or lack an ovary altogether.