

Zeitschrift: Boissiera : mémoires de botanique systématique
Herausgeber: Conservatoire et Jardin Botaniques de la Ville de Genève
Band: 74 (2024)

Artikel: Taxonomic revision of the genus *Pentacalia* (Compositae, Senecioneae) in Ecuador
Autor: Calvo, Joel
Kapitel: Material and methods
DOI: <https://doi.org/10.5169/seals-1052964>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 13.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

although the support is generally low too. *Pentacalia trianae* from Colombia and *P. arborea* from Colombia and Ecuador are strongly related to each other, and found in a subclade along with species of *Monticalia* and *Scrobicaria* Cass. This latter genus contains three species restricted to Colombia and Venezuela and mainly differs from *Pentacalia* in having opposite leaves. The accession of *P. desiderabilis*, a species endemic to Brazil, is nested among species of *Dendrophorbium* and *Graphistylis* B. Nord. also from Brazil. These results, though weakly supported, suggest that *Pentacalia* as currently circumscribed is not monophyletic. Additional studies based on an exhaustive sampling including species of *Pentacalia* that represent the entire distribution area of the genus, as well as species of the aforementioned allied genera, are essential for drawing the evolutionary relationships of these plant groups and, if needed, make the taxonomic adjustments accordingly.

Material and methods

The present contribution is the result of an exhaustive review of the published bibliography, fieldwork in Ecuador carried out in 2017, 2018, and 2023, and the examination of 369 herbarium specimens (duplicates excluded) kept at the following herbaria: AAU, G, HA, HUTPL, LOJA, MA, P, Q, QAP, QCA, QCNE, QPLS, US, and USM. Additionally, digital images of specimens from B, COL, E, F, GH, HAL, K, LD, LP, MO, MT, NY, P, PR, PRC, S, UC, US, VALLE, and VEN were studied. The loans of specimens received from AAU, MA, P, and US were essential to achieve this revision.

The types of all accepted names and synonyms were studied, except for *Pentacalia pailasensis* H. Rob. & Cuatrec., the holotype of which appears to be lost after having been loaned by F to another institution (K. Hansen, pers. comm.). It is therefore treated as doubtful. The qualitative characters provided in the descriptions were studied, when needed, with the aid of the stereo microscope Leica M60, whereas quantitative characters were recorded using a Mitutoyo digital caliper, CD-15DC. The maps were generated using QGIS 3.16 Hannover. Information concerning the habitat, elevation, and flowering period of each species was obtained from the herbarium specimen labels and the information compiled in the field. Nineteen (19) out of the 27 accepted species were observed in Ecuador during the field trips.

Accepted species are presented in alphabetical order and are listed in Appendix 1. All studied exsiccatae can be found in Appendix 2. An index to all scientific names is provided at the end of this publication. The list of specimens examined given for each species includes only the Ecuadorian material, except for the species that citing the collections from the neighboring countries is relevant for completeness sake. If herbarium specimens from a province were not studied but the presence of the species is expected there, the name of that province is included in the distribution and followed by a question mark.

The leaf descriptions and measurements provided in this treatment correspond to the leaf lamina. General botanical terms largely follow BEENTJE (2012), whereas specific terms concerning the family *Compositae* are in line with those used by ROBINSON & CUATRECASAS (1993).

Taxonomic treatment

Pentacalia Cass. in Cuvier F., Dict. Sci. Nat., ed. 2, 48: 461. 1827.

Typus: *Cacalia arborea* Kunth [= *Pentacalia arborea* (Kunth) Cass.].

= *Senecio* sect. *Triana* Cuatrec. in Fieldiana, Bot. 27(2): 71. 1951. = *Pentacalia* Cass. subg. *Pentacalia*.

= *Senecio* sect. *Streptothamni* Greenm. in Bot. Jahrb. Syst. 32: 19. 1902. **Typus:** *Senecio streptothamnus* Greenm. [= *Pentacalia streptothamna* (Greenm.) H. Rob. & Cuatrec.].

Plants scandent, woody, with long dangling branches, sometimes rather suberect with branches leaning over adjacent vegetation; stems terete, often furrowed, glabrous or with indumentum, solid or fistulose. *Leaves* alternate (opposite in 3 Peruvian species), simple, petiolate; laminas lanceolate, oblanceolate, elliptic, ovate, or obovate, apex rounded to acuminate, base attenuate to cordate, margin entire to dentate (usually remotely mucronate-denticulate), glabrous to covered by different types of indumentum (trichomes usually simple, multicellular, eglandular, rarely T-shaped in few non-Ecuadorian species), chartaceous, coriaceous, or fleshy, concolorous or slightly discolored, secondary and tertiary veins conspicuous or inconspicuous. *Synflorescences* terminal or lateral (axillary), corymbiform, thyrsoide-paniculiform, or racemiform. *Capitula* radiate, disciform, or discoid, sessile to pedunculate; involucre cylindrical to campanulate, glabrous or covered by indumentum; receptacles epaleaceous, rather flat, smooth or somewhat irregularly alveolate; involucre bracts 1-seriate, usually 8–13, linear-oblong; supplementary bracts (calycle) (1–)3–5(–7), linear-subulate to ovate. *Ray florets* pistillate; corollas with reduced or well-developed limbs, patent or curved downward, subentire to 3-toothed at apex, yellow or white (becoming red burgundy as florets mature in few species). *Peripheral florets* of disciform capitula pistillate; corollas tubular, usually shorter than disc florets, deeply (2–)4–5-lobed, yellow or whitish. *Disc florets* hermaphroditic; corollas tubular, 5-lobed, yellow or whitish (becoming red burgundy as florets mature in few species); filament collars balustriform; anther bases caudate (sometimes only very shortly caudate), anther appendages 2–3 times longer than wide; style branches truncate with crown of sweeping trichomes. *Achenes* cylindrical, 5–10-ribbed, usually glabrous (pubescent in few non-Ecuadorian species); pappus usually 1-seriate, bristles capillary, barbellate, whitish.