

Zeitschrift: IABSE publications = Mémoires AIPC = IVBH Abhandlungen

Band: 28 (1968)

Artikel: Semi-infinite plate subjected to the concentrated load acting at its internal point

Autor: Övünç, Bülent

DOI: <https://doi.org/10.5169/seals-22173>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 05.09.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Semi-Infinite Plate Subjected to the Concentrated Load Acting at its Internal Point

Plaque semi-infinie soumise à une charge concentrée agissant en un point intérieur

Halbebenen-Platte unter Einzellast im Innern

BÜLENT ÖVÜNÇ *)

Ph.D., Academic Research Fellow, University of British Columbia, Vancouver, Canada

In this paper, the complex variable method of N. I. MUSKHELISHVILI is applied to the solution of semi-infinite plate subjected to the concentrated load acting at its internal point. The coefficients of the two functions of complex variable, whose combination gives the Airy Stress function, are taken as unknowns. Then they are determined by using the boundary conditions and considering the limits of these two functions while the loaded point approaches to the boundary or tends to the infinity. Two examples are given in order to visualize the stress pattern in the semi-infinite plate.




Fig. 1.

Consider the region S occupied by the semi-infinite plate as the "lower" half plane (Fig. 1) bounded by the horizontal line mn at a distance ρ from the origin O . Let the horizontal and vertical components of the load, P , applied at the origin O , be respectively X , Y and let the two independent functions of complex variable be:

*) Presently Associate Professor, University of Southwestern Louisiana, U.S.A.

$$\begin{aligned}\tau(z) &= -(A+iB)\log z - (C+iD)\log(z-2i\rho) + \frac{E+iF}{z-2i\rho}, \\ \psi(z) &= (C-iD)\log z + (A-iB)\log(z-2i\rho) + \frac{E+iF}{z-2i\rho},\end{aligned}\quad (1)$$

where the constants A, B, C, D, E, F , are the unknowns to be determined by the boundary conditions and from the limiting conditions corresponding to ρ , approaching zero or infinity.

The boundary condition is given by MUSKHELISHVILI¹⁾

$$\tau(t) + t\overline{\tau'(t)} + \overline{\psi(t)} = \text{constant}, \quad t = x + i\rho. \quad (2)$$

Substituting the expressions (1) into the boundary condition (2) one obtains the following relations between the constants

$$E = 2\rho B, \quad F = 2\rho A.$$

Then $\tau(z)$ and $\psi(z)$ become:

$$\begin{aligned}\tau(z) &= -(A+iB)\log z - (C+iD)\log(z-2i\rho) + 2i\rho \frac{A-iB}{z-2i\rho}, \\ \psi(z) &= (C-iD)\log z + (A-iB)\log(z-i\rho) + 2i\rho \frac{A-iB}{z-2i\rho}.\end{aligned}\quad (3)$$

The first derivatives of the functions $\tau(z)$ and $\psi(z)$ are:

$$\begin{aligned}\Phi(z) &= -\frac{A+iB}{z} - \frac{C+iD}{z-2i\rho} - 2i\rho \frac{A-iB}{(z-2i\rho)^2}, \\ \Psi(z) &= \frac{C-iD}{z} + \frac{A-iB}{z-2i\rho} - 2i\rho \frac{A-iB}{(z-2i\rho)^2},\end{aligned}\quad (4)$$

where,

$$\Phi(z) = \tau'(z), \quad \Psi(z) = \psi'(z).$$

Now, consider the case when ρ approaches zero. Putting $\rho=0$ in the relations (4) one obtains

$$\Phi(z) = -\frac{A+iB}{z} - \frac{C+iD}{z}, \quad \Psi(z) = \frac{C-iD}{z} + \frac{A-iB}{z}. \quad (5)$$

Similarly, when ρ increases to infinity the relations (4) become

$$\Phi(z) = -\frac{A+iB}{z}, \quad \Psi(z) = \frac{C-iD}{z}. \quad (6)$$

But the solution of a semi-infinite plate under the concentrated load applied to its boundary is known.

¹⁾ N. I. MUSKHELISHVILI: Some Basic Problems of the Mathematical Theory of Elasticity, translated from the Russian by J. R. M. Radok, P. Noordhoff, Ltd., Groningen 1963.

$$\Phi(z) = -\frac{X+iY}{2\pi} \frac{1}{z}, \quad \Psi(z) = \frac{X-iY}{2\pi} \frac{1}{z}. \quad (7)$$

Similarly the known solution of the infinite plate under a concentrated load gives

$$\Phi(z) = -\frac{X+iY}{2\pi(1+\chi)} \frac{1}{z}, \quad \Psi(z) = \frac{\chi(X-iY)}{2\pi(1+\chi)} \frac{1}{z}, \quad (8)$$

where $\chi = 3-\nu$ for plane strain conditions

and $\chi = \frac{3-\nu}{1-\nu}$ for generalized plane stress conditions,

ν being the Poisson's ratio.

The remaining unknown constants A, B, C, D , can be determined from the relations (5), (6), (7), (8) as follows:

$$A = \frac{X}{2\pi(1+\chi)}, \quad B = \frac{Y}{2\pi(1+\chi)}, \quad C = \frac{\chi X}{2\pi(1+\chi)}, \quad D = \frac{\chi Y}{2\pi(1+\chi)}. \quad (9)$$

Finally, by substituting the expressions for the coefficients (9) in the relations (3), the functions $\tau(z)$ and $\psi(z)$ are found as follows.

$$\begin{aligned} \tau(z) &= \frac{1}{2\pi(1+\chi)} \left[-(X+iY)(\log z + \chi \log(z-2i\rho) + 2i\rho \frac{X-iY}{z-2i\rho}) \right], \\ \psi(z) &= \frac{X+iY}{2\pi(1+\chi)} \left[\chi \log z + \log(z-2i\rho) + \frac{2i\rho}{z-2i\rho} \right]. \end{aligned} \quad (10)$$




Fig. 2.

Example 1. Let the force P acting at the point 0 be horizontal, Fig. 2. MUSKHELISHVILI gives the following expressions for the stresses in polar coordinates²⁾.

$$\sigma_\rho + \sigma_\theta = 2[\phi(z) + \bar{\phi}(z)], \quad \sigma_\theta - \sigma_\rho + 2i\tau_{\rho\theta} = 2[\bar{z}\phi'(z) + \Psi(z)]e^{2i\theta}. \quad (11)$$

Taking the derivatives of the functions $\tau(z)$ and $\psi(z)$ in (10) and substituting them in (11), the stress components are obtained in polar coordinates as follows:

²⁾ Please refer to footnote 1.

$$\begin{aligned}\sigma_\theta &= \frac{P}{2\pi(1+\chi)} \left[\frac{(\chi-1)\cos\theta}{r} + \frac{(1-2\chi)r\cos\theta - 2\rho\sin 2\theta}{r^2 - 4\rho r\sin\theta + 4\rho^2} \right. \\ &\quad + \frac{[\chi r^2 + 4(6-\chi)\rho^2]r\cos\theta - 4(r^2 + 2\rho^2)\rho\sin 2\theta}{(r^2 - 4\rho r\sin\theta + 4\rho^2)^2} \\ &\quad \left. + 4\rho r \frac{r^3\sin 2\theta - 6\rho r^2\cos\theta + 8\rho^3\cos\theta}{(r^2 - 4\rho r\sin\theta + 4\rho^2)^3} \right], \\ \sigma_\rho &= \frac{P}{2\pi(1+\chi)} \left[(3+\chi) \frac{\cos\theta}{r} + \frac{(1+2\chi)r\cos\theta - 2\rho\sin 2\theta}{r^2 - 4\rho r\sin\theta + 4\rho^2} \right. \\ &\quad + \frac{[\chi r^2 - 4(\chi+2)\rho^2]r\cos\theta + 4(r^2 - 2\rho^2)\rho\sin 2\theta}{(r^2 - 4\rho r\sin\theta + 4\rho^2)^2} \\ &\quad \left. + 4\rho r \frac{r^3\sin 2\theta - 6\rho r^2\cos\theta + 8\rho^3\cos\theta}{(r^2 - 4\rho r\sin\theta + 4\rho^2)^3} \right], \\ \tau_{\rho\theta} &= \frac{P}{2\pi(1+\chi)} \left[(\chi-1) \frac{\sin\theta}{r} + \frac{r\sin\theta + 2\rho\cos 2\theta}{r^2 - 4\rho r\sin\theta + 4\rho^2} \right. \\ &\quad + \frac{[4\rho^2r(2-\chi) - \chi r^3]\sin\theta + 8\rho^3\cos 2\theta - 2\rho r^2(1-2\chi)}{(r^2 - 4\rho r\sin\theta + 4\rho^2)^2} \\ &\quad \left. + 4\rho r \frac{r^3\cos 2\theta + 6\rho r^2\sin\theta + 8\rho^3\sin\theta - 12\rho^2r}{(r^2 - 4\rho r\sin\theta + 4\rho^2)^3} \right].\end{aligned}$$

Example 2. The concentrated load P and the Poisson ratio ν are taken

$$P = 100\pi e^{\frac{i\pi}{6}}, \quad \nu = 0.3.$$

For a possible comparison with the results of photo-elastic experiments the calculation is made of isoclinic and isochromatic lines. The former (Fig. 3) keep constant angle with the directions of principal stresses at all points, and the latter (Fig. 4) correspond to lines along which the difference of principal stresses is constant.

The calculations have been made with the IBM 7040 computer.




Fig. 3. Isoclinics.




Fig. 4. Isochromatics.

Acknowledgement

The author wishes to express gratitude to Professor Dr. SEMIH TEZCAN and to the National Research Council for financial support.

Summary

An attempt has been made to obtain the solution of the presented problem by considering the solution of the semi-infinite plate loaded at the boundary and the solution of infinite plate loaded at the internal point are its limit cases.

Résumé

On a essayé d'obtenir la solution du problème présenté en examinant la solution d'une plaque semi-infinie chargée sur son bord, et la solution d'une plaque infinie chargée sur un point intérieur qui sont ses cas limites.

Zusammenfassung

Für das vorgelegte Problem ward der Versuch unternommen, eine Lösung zu finden, in Anbetracht dessen, daß die Lösungen der Halbebene für Lasten einmal am Rand und im Innern die Grenzfälle darstellen.

Leere Seite
Blank page
Page vide